VOL. 33. Ser. A. Title-page and Index. [Issued January, 1947]

THE REVIEW OF APPLIED ENTOMOLOGY

SERIES A: AGRICULTURAL

ISSUED BY THE IMPERIAL INSTITUTE OF ENTOMOLOGY

THE IMPERIAL INSTITUTE OF ENTOMOLOGY, 41, QUEEN'S GATE, S.W.7

Price 6s. net

All Rights Reserved

IMPERIAL INSTITUTE OF ENTOMOLOGY

Executive Council

G. H. Creasy, C.M.G., O.B.E., Chairman, Colonies, Protectorates and Mandated Territories.

Lieutenant-Colonel J. G. ROBERTSON, Vice-Chairman, Canada.

Sir Donald Fergusson, K.C.B., United Kingdom.

F. L. McDougall, C.M.G., Australia.

A. L. POOLE, New Zealand.

A. P. VAN DER POST, Union of South Africa.

J. W. DULANTY, C.B., C.B.E., High Commissioner, Eire.

D. James Davies, C.B.E., Newfoundland.

Sir DAVID B. MEEK, C.I.E., O.B.E., India:

B. O. BINNS, O.B.E., Burma.

W. C. ROBERTSON, Southern Rhodesia.

Sir David Chadwick, K.C.M.G., C.S.I., C.I.E., Secretary.

Director and Editor

S. A. NEAVE, C.M.G., O.B.E., D.Sc.

Assistant Director

W. J. HALL, M.C., D.Sc.

Assistant Editor

H. S. Bushell, M.A.

Head Office—British Museum (Natural History), Cromwell Road, London, S.W. 7.

Publication Office and Library-41, Queen's Gate, London, S.W. 7.

Director, Imperial Parasite Service

W. R. THOMPSON, D.Sc., F.R.S. 228, Dundas Street, Belleville, Ontario, Canada.

INDEX OF AUTHORS.

A reference in heavy type indicates that a paper by the author has been noticed.

Abbas, Manzoor, 296, 386. Abbott, W. S., 63. Abraham, R., 157. Acree jr., F., 239. Afzal Husain, Mohammad, 296, 362, 386, 388. Ahlberg, O., 35. Ahmad, Muzzafer, 389. Aitken, T. R., 402. Aiyar, K. S. Padmanabha, 387. Alexander, C. C., 272. Alexander, P., 156. Alfaro, A., 165, 166. Allen, H. W., 105. Allmendinger, D. F., 284. Ananda Rau, S., 246. Anderson, K. T., 228. Anderson, L. D., 51, 59, 214. Anderson, R. F., 271. André, M., 328, 356, 357. Andrewartha, H. G., 76, 136. Annand, P. N., 258. Anson, R. R., 87. Anwar, M. S., 391. Appanna, M., 164, 362. Apple, J. W., 278. Arbuthnot, K. D., 151. Armstrong, W. D., 315. Arnoldi, K. V., 159, 301. Arnott, D. A., 140. Arthur, D. R., 77, 393. Arthur, J. M., 245. Ashdown, D., 52, 180. Ass, M. J., 14. Atherton, D. O., 369. Atwood, C. E., 143, 354. Austin, L., 398. Avens, A. W., 58. Averill, A. W., 406. Ayyar, P. N. Krishna, 20, 164. Ayyar, T. V. Ramakrishna, 386.

Baeta Neves, C. M., 309, 310. Baker, A. C., 395. Baker, A. D., 141, 145. Baker, E. W., 52.

Azevedo e Silva, F., 310.

Baker, H., 275. Baker, M. R., 137. Balch, R. E., 154, 398. Bald, J. G., 33, 93, 378. Balzer, A. I., 235, 236. Barber, G. W., 337. Barber, H. S., 41, 45. Barbieri, N. A., 120, 252. Bare, C. O., 348. Barnes, H. F., **74**, **170**, **293**, **308**. Barnes, H. M., 244. Barnes, M. M., 179, 274. Barnes, O. L., 215, 341. Barthel, W. F., 189, 332, 368. Bartlett, K. A., 26, 312. Basu, A. C., 363. Bates, M., 289. Bawden, F. C., 192. Beall, G., 139. Beard, R. L., 280. Bebbington, A. G., 88. Becker, G., 18. Beckwith, C. S., **401**. Bedford, E. C. G., **324**. Bellod, M., 169. Benitez Morera, A., 170. Bennett, A. H., 233. Bergamin, J., 203, 205. Birch, L. C., 76. Bird, F. T., 398. Bishop, C. F., 173. Blair, R., 46. Blake, M. A., 315. Blatchley, W. S., 146. Blattny, C., 120, 252. Blauvelt, W. E., **180.** Blickle, R. L., 26. Blijdorp, P. A., 253. Blunck, H., 9. Boczkowska, M., 120, 298. Bodenheimer, F. S., 199, 225. Bodenstein, O. F., 212. Böhme, H. Müller-, 159. Bolwig, N., 380. Borden, A. D., 282. Börner, C., 118. Bose, M., 388.

Böttcher, F. K., 6.

Bourne, A. I., 56, 212, 272. Bovey, P., 394. Bovien, P., **12**, **252**, **380**, **381**, **382**. Böving, A. G., **163**. Bovingdon, H. H. S., 266, 295. Bowen, C. V., 54, 176, 214, 368. Box, H. E., 195. Boyce, A. M., 113. Bragg, A. N., 96. Bramble, W. C., **286.** Brammanis, L., 7. Brann jr., J. L., 148. Brasnett, N. V., 329. Bray, S. P. V., 92. Breakey, E. P., 23, 64. Brecheen, K. G., 398. Bredo, H. J., 366. Brett, C. H., 216. Briand, L. J., 142. Briejèr, C. J., 229. Brierley, P., 1, 2, 38, 70, 174. Briscoe, H. V. A., 156. Broadbent, B. M., 342. Broadhead, E., 181. Bromley, S. W., 43. Bronson, T. E., 214, 276. Brown, F. M., 81. Brown, L. R., 97. Brown, R. C., 238, 313. Bruel, W. E. van den, 120. Brunson, M. H., 105. Bua, G., **118.** Bugbee, R. E., 241. Burdick, H. L., 277. Busbey, R. L., 271. Butovitsch, V., 32.

Caldwell, N. E. H., 162. Callan, E. McC., 147. Calvery, H. O., **160.** Cameron, A. E., 233. Cameron, G. S., 87. Campau, E. J., 277, 282, 283. Campbell, G. A., **137.** Cañizo Gomez, J. del, 167, 328. Capoor, S. P., 392. Carlson, F. W., 56, 108, 274, 281. Carpenter, J. M., 70. Carter, Walter, 249, 345. Cartwright, W. B., 103. Casper, M., 315. Cassab, A., 226. Cassil, C. C., 128, 272, 281. Castillo, J. C., 264. Castro, A. Ruiz, 167, 298.

Chada, H. L., 79. Chamberlin, F. S., 54, 215. Chamberlin, T. R., 61. Chambers, V. H., 171. Chandler, S. C., 316. Chapin, E. A., 164. Chapman, P. J., **58.** Chapman, R. K., 140. Chatterjee, P. N., 390. Chauvin, R., 201. Cherian, M. C., 384. Childers, N. F., 314. Chisholm, R. D., 368. Chisholm II, J. J., 284. Chorley, J. K., 126. Christensen, J. R., 242. Clancy, D. W., 109, 208, 222. Clark, A. H., 81. Clemente, F. Gómez-, 168. Cockerham, K. L., 216. Collins, D. L., 180. Cook, C. W., 242. Cory, E. N., 49, 55, 284. Costa Lima, A. da, 401. Cotterell, G. S., 65. Cotton, R. T., 102, 235, 236. Cox, A. J., **190.** Coyne, F. P., 295. Craig, R., 347. Craighead, F. C., 287. Cressman, A. W., 342, 346. Crouzel, I. Santoro de, 401. Crowley, D. J., 23. Custodio P., T., 131. Cutright, C. R., 191, 208.

da Costa Lima, A., 401. Dadin, J. M., 125. da Fonseca, J. P., 311. Daguerre, J. B., **120.** Daines, R. H., 39, 315. Darrow, G. M., 401. Daviault, L., 223. Davidson, J., 129, 187. Davis, E. G., 176. Dean, F. P., 272. Dean, R. W., 148, 150. Decker, G. C., 57, 175. de Crouzel, I. Santoro, 401. de Figueiredo jr., E. R., 375. de Koning, M., 7. Delassus, M., **403.** del Cañizo Gomez, J., 167, 328. Delisle, R., 142. Denne, W., **369.**

De Santis, L., 117, 132. Dewey, J. E., 180. Dicker, G. H. L., 181. Dickinson, B. C., 60. Dickinson, R. C., 113. Dierick, G. F. E. M., 15. Dieter, C. E., 277. Dills, L. E., 53, 276. Dirks, C. O., 42, 47, 55. Ditman, L. P., 49. Dohanian, S. M., 341. Dolby, R. M., 130. Domínguez García-Tejero, F., 166. Dominick, C. B., 41: Donohoe, H. C., 58, 110. Doolittle, S. P., 39. Doudoroff, M., 81. Douglass, J. R., 281. Dow, D., 113. Downes, W., 334. Doyle, J. A., 139, 224. Draize, J. H., 160. Drake, C. J., 175. Driggers, B. F., 46, 213. Duck, L. G., 237. Dudley jr., J. E., 214, 276. Due, J. P. La, 113, 349. Duff, C. E., 247. Dugas, A. L., 25. Dumbleton, L. J., 183. Duncan, J., 138, 224. Dwarka Nath Nanda, 296. Dyk, A., 8.

Eastham, J. W., 154. Ebeling, W., 253, 349. Eckstein, F., 9. Eckstein, K., 6. Edelsten, H. M., 307. Emden, F. I. van, 296. Engel, H., 6. Engelhart, W., 182. Erickson, R. O., 180. Essig, E. O., 368. Evans, A. C., 294. Evans, J. A., 179. Evelyn, S. H., 87. Ewart, W. H., 52. Ewing, K. P., 210. Ext, W., 9. Eyer, J. R., 68.

Fadigas jr., M., 94. Faes, H., 124.

Falconer, D. S., 123. Farstad, C. W., 98. Faure, J. C., 246, 327. Fawcett, H. S., 128. Fedotov, D. M., 301. Felt, E. P., 43. Fennah, R. G., 313. Fenton, F. A., 151. Ferguson, G. R., 342. Feytaud, J., 157, 158. Figueiredo jr., E. R. de, 375. Filmer, R. S., 48, 213. Fisher, R. C., 181. Fitzgerald, J. S., 372. Fitzhugh, O. G., 160. Flanders, S. E., 59, 61, 105, 107, 214, 281, 374. Fleck, E. E., **347.** Fletcher, F. W., 343. Florov, D. N., 400. Fonseca, J. P. da, 311. Forsslund, K. H., 6. Fox, W. B., 153. Fox-Wilson, G. (see Wilson, G. F.). Frankenfeld, J. C., 235. Franzke, A., 228. Frazer, H. L., 357. Frazier, N. W., 172. Frear, D. E. H., 116, 275, 276. Frost, S. W., 53. Fryer, J. C. F., 265. Fullaway, D. T., 251. Fulmek, L., 117. Fulton, R. A., 271, 346. Funtikow, G. P., 14. Furniss, R. L., 40.

Gäbler, H., 252. Gadd, C. H., 294, 300. Gahan, A. B., 95. Gahan, J. B., 279. Gaines, J. C., 335, 336. García Mercet, R., 357. García Rada, G., 132. García-Tejero, F. Domínguez, 166. Gardner, J. C. M., 7, 264, 371. Garg, R. L., 406. Gauthier, G., 141. Gautier, C., 297. Geinitz, B., 6. Gentner, L. G., 220. Gerasimov, V., 304. Gertler, S. I., 107, 344. Giannotti, O., 205, 288. Gilvarov, M. S., 245.

Given, B. B., 247, 376. Glasgow, R. D., 46. Glass, E. H., 103. Gleissner, B. D., 27. Glendenning, R., 333. Goble, H. W., 140. Golding, F. D., 165, 383. Gomez, J. del Cañizo, 167, 328. Gómez-Clemente, F., 168. Gómez-Menor, J., 166. Gooden, E. L., 42. Goodhue, L. D., 43, 99, 264, 332. Goodwin, M. W., 221. Gösswald, K., 15, 16. Götz, B., 14, 253. Gough, H. C., 393. Gradojević, M., 5. Graham, A. R., 142. Graham, L. T., 271. Granger, A. M., 328. Granovsky, A. A., 207. Grayson, J. M., 45, 280, 281. Gries, G. A., 378. Grieve, B. J., 292. Griot, M., 310. Grison, P., 158. Griswold, G. H., 350. Guénaux, G., 328. Gunther, F. A., 349. Günther, O., 120. Gupta, B. D., 406. Györfi, J., 18. Gyrisko, G. G., 340.

Habib, A., 226. Haenseler, C. M., 239. Haeussler, G. J., 208, 348. Haller, H. L., 99, 106, 189, 239, 368. Hallock, H. C., 281. Hambleton, E. J., 278. Hamilton, A., 163, 184. Hamilton, D. W., 149. Hammond, G. H., 140, 180. Hamner, A. L., 25, 80. Hamnett, G. G., 405. Hansberry, T. R., 100, 179, 277. Hansen, E. L., 347. Hansen, J. W., 128, 347. Hansford, C. G., 329. Harman, S. W., 42. Harries, F. H., 220. Harris, W. V., 85, 245. Harrison, A., 128. Harrison, G. H., 73. Hartzell, A., 65, 245.

Hartzell, F. Z., 44. Hastings, E., 264. Hatch, M. B., 152. Hatfield, I., 69. Hattingh, C. C., 246. Hawboldt, L. S., **154**, **353**, **354**. Hayward, K. J., **4**, **132**, **133**, **134**, **287**. Heal, G. M., 155. Hebard, M., 146. Heidenreich, E., 7. Helson, G. A. H., 93, 323. Hely, P. C., **249**, **267**. Henderson, W. W., **174**. Hensill, G. S., 240. Herce, P., 169. Heriot, A. D., 332. Hespeler, O., 229. Heuberger, J. W., 173. Hey, G. L., **171, 233.** Hill, R. E., **216.** Hills, O. A., 279. Hinton, H. E., 135, 330. Hobby, B. M., 181. Hocking, B., 71, 72, 73. Hodson, W. E. H., 117. Hoerner, J. L., 219. Hogan, T. W., 377. Holdaway, F. G., 217, 251. Holst, E. C., 286. Hopperstead, S. L., 221. Horovitz, S., 67. Horsfall, J. G., **173.** Horsfall, J. L., **44.** Hoskins, W. M., **368.** Hosny, M., 227. Houston, B. R., 378. Hovanitz, W., 339. Hovasse, R., 356. Howard, N. F., 100. Hrisafi, C., 121. Huffaker, C. B., 79, 160. Hughes, J. H., 29. Husain, Mohammad Afzal, 296, 362, 386, 388. Hussein, M., 228. Hutson, R., 223.

Isely, D., **191.** Ivy, E. E., **57, 210.**

Jacob, F. H., **92**, **358**. Jameson, J. D., **88**. Jancke, O., **13**, **117**.

Janes, R. L., 277, 283.
Janisch, E., 7.
Jannone, G., 379.
Jarvis, H., 248.
Jenkins, C. F. H., 163.
Joffe, J. S., 63.
Johnson, A. C., 50.
Jones, E. W., 23.
Jones, J. R. E., 405.
Jones, L. K., 39, 70, 174.
Jones, M. G., 382.

Kadocsa, G., 11. Kadow, K. J., 221. Kalmus, H., 188. Kangas, E., 7. Kapur, A. P., 389. Karny, H. H., 146. Keenan, W. N., 142, 224. Keifer, H. H., 111. Kelsey, J. M., 234, 369. Kemper, H., 229. Kenaga, E. E., 343. Khvostova, V. B., 304. Kimmey, J. W., 40. King, H. E., 87. King, H. L., 275, 276. King, K. M., 144. Kirkor, S., 328. Kitchener, J. A., 156. Kloet, G. S., 92. Knechtel, W., 121. Knight, R. L., 87. Knowlton, G. F., 106, 206, 263. Koblitsky, L., 368. Köhler, P., 252. Kolesnik, I., 302. Komárek, J., 8. Koning, M. de, 7. Koppán, J., 6. Korschefsky, R., 11. Korsmeier, R. B., 113. Kosmachevsky, A. S., 159, 160. Kozikowski, A., 9. Krauss, N. L. H., 251. Krishna Ayyar, P. N., 20, 164. Krishnamurti, B., 164, 362. Kulash, W. M., 106. Kumar Mitra, Prabhas, 128, 393. Kunike, G., 230. Kuppuswami, 371. Küthe, K., 10, 119.

La Due, J. P., 113, 349. LaForge, F. B., 189. LaHue, D. W., 103. Lal, K. B., 300, 362. Lambert, R., 354. Lamerson, P. G., 117. Landaluze, P. Urquijo, 125. Lane, M. C., 23. Lange jr., W. H., 104, 337. Langford, G. S., 49, 55, 284. La Rivers, I., 146, 368. Lathrop, F. H., 42, 55, 222. Latta, R., 50. Leach, J. G., 173. Lechevalier, H., 71. Leech, H. B., 287, 354. Lefèvre, P. C., 36, 363, 367. Leius, K., 119. Lepage, H. S., **94**, **204**, **205**, **288**. Lepesme, P., **228**, **230**, **403**. Lepigre, A. L., 403. Lever, R. J. A. W., 136, 161, 201, 290, 291. Liégeois, P., 365. Lillie, R. D., 160. Lima, A. da Costa, 401. Lima, L., 375. Lindgren, D. L., 113. Linsley, E. G., 25, 102. List, G. M., 30. Liu, C. L., 369. Lloyd, D. C., 240. Lloyd, N. C., 184. Loewel, E. L., 6, 34. Logan, S. H., 179. Look, W. C., 251. Lopez, L. Rodriguez, 5, 95. Lord, F. T., 144. Lovàszy, P., 11, 12. Lubatti, O. F., 128. Ludwig, D., **406.** Lyle, C., 25.

McAfee, E. L., 251.
Macdonald, D., 87.
Macfie, J. W. S., 78.
MacGill, E. I., 135.
McGovran, E. R., 98.
McGregor, E. A., 132 (note), 397.
McGuffin, W. C., 180.
McHardy, J. W., 233.
Mackie, J. R., 370.
McKinney, K. B., 112, 279.
McKinstry, A. H., 89.
McLean, A., 266.
McLean, H. C., 63.
MacLeod, G. F., 30.

McLeod, W. S., 188. McNally, A. G., 141. McPhail, M., 395. Madden, A. H., 192. Madel, W., 230. Magie, R. O., 317. Makkus, W., 158. Mallis, A., 318, 343. Maltais, J. B., 142, 223. Manton, S. M., 135. Manzoor Abbas, 296, 386. Marchioni, A. H., 66, 67. Margabandhu, V., 164. Márquez, V. Moreno, 166. Martin, H., 197, 252, 253, 254. Martorell, L. F., 110, 243. Marvin, P. H., 218. Massee, A. M., 359, 361. Mathis, W., 211, 341. Mathur, C. B., 386, 388. Mathur, R. N., 388. May, A. W. S., 162. Mayer, E. L., 279, 344. Mayné, R., 157. Medler, J. T., 63. Meier, K., 124. Mendes, C. Teixeira, 375. Menor, J. Gómez-, 166. Menozzi, C., 166. Menzel, R., 124. Mercet, R. García, 357. Meyer, E., 10. Michelbacher, A. E., 25, 30, 40, 81, **241**, **282**. Milam, J., 344. Miles, H. W., 198, 199, 368. Miller, L. W., 129, 235, 292. Miner, F. D., 191, 270. Minkiewicz, S., 117. Mischon, 230. Mitchell, B. L., 91. Mitra, Prabhas Kumar, 128, 393. "Mjele", 327. Mohammad Afzal Husain, 296, 362, 386, 388. Molestina O., E., 95. Monro, H. A. U., 142. Monte, O., 95, 263, 287, 311. Moreno Márquez, V., 166. Morera, A. Benitez, 170. Morgan, W. L., 185. Morison, G. D., 121. Morley, B. D. W., 122. Mors, H., 8. Mote, D. C., 342. Mott, L. O., 53.

Moutia, A., 370.
Muesebeck, C. F. W., 116.
Muggeridge, J., 130, 375.
Mühlow, J., 36.
Muljee, 372.
Müller-Böhme, H., 159.
Muma, M. H., 55, 347.
Munger, F., 216, 346.
Munro, J. A., 96.
Munson, S. C., 64.
Muzzafer Ahmad, 389.

Nanda, Dwarka Nath, 296. Neiswander, C. R., 67. Neiswander, R. B., 262. Nel, R. G., 127. Nelson, A. A., 160. Nepveu, P., 5. Neu, W., 10. Neves, C. M. Baeta, 309, 310. Newcomer, E. J., **272.** Nicholas, J. E., 53. Nicholson, A. J., 254. Nickels, C. B., 209. Nishida, T., 217, 251. Nolte, H. W., 8. Norris, D. O., 33, 378. Norton, L. B., 179. Notley, F. B., 128. Nunberg, M., 7.

Oakley, R. G., 163.
Occhioni, P., 134.
O'Neal, E. J., 63.
Oosthuizen, M. J., 326, 327.
Orelli, O. Schneider-, 194, 379.
Örösi-Pál, Z., 6.
Ortega, J. J., 349.
Osburn, M. R., 211, 341.
Osnitzkaya, E., 304.
Ostapetz, A., 302.
Overholser, E. L., 284.
Overley, F. L., 284.
Owens, H. B., 49.

Padmanabha Aiyar, K. S., 387. Paillot, A., 328. Pál, Z. Örösi-, 6. Palm, C. E., 179, 180. Parker, J. R., 59. Parker, R. L., 53, 117. Parks, T. H., 191. Parrot, A. G., 357.

Paul, L. C., 144. Pavlakos, J. G., 16. Pearce, G. W., 58. Pearson, E. O., 89, 128. Peat, J. E., 88. Peay, W. E., 281. Peeters, G. A., 157. Peiris, J. W. L., 299. Pepper, B. B., 48, 221, 239. Pepper, J. H., 264. Perron, J. P., 141, 145. Persing, C. O., 97. Petch, T., 232. Peters, G., 252. Peterson, L., 354. Petherbridge, F. R., 297. Petkov (Petkoff), P., 119, 122. Pfeffer, A., 8. Picard, F., 166. Pierce, W. C., 209, 346. Planes, S., 169. Plank, H. K., 312, 338. Plumb, G. H., 378. Plummer, B. E., 55. Plummer, C. C., 190, 395. Poeteren, N. van, 12, 157. Polezhaev, V. G., 330. Ponomarenko, D., 305. Potts, S. F., 62, 239. Prabhaker Rao, V., 391. Prabhas Kumar Mitra, 128, 393. Pradhan, S., 390. Pratt jr., J. J., 206. Pravdin, F. N., 245. Prentice, A. N., 88. Prill, E. A., 245. Procter, C. H., 247. Pruthi, H. S., 391.

Rada, G. Garcia, 132.
Rahman, K. A., 300.
Rajarao, S. A., 393.
Ramakrishna Ayyar, T. V., 386.
Ramchandra Rao, Y., 384, 388.
Rao, V. Prabhaker, 391.
Rao, Y. Ramchandra, 384, 388.
Rašek, J. M., 8.
Rau, S. Ananda, 246.
Raucourt, M., 331.
Rawlins, W. A., 180, 206.
Recordati, A. Zappi, 6.
Records, E., 59.
Rehn, J. A. G., 146.
Reigel, A., 241.
Reinhard, H. J., 270.

Reynolds, J. M., 156. Rice, P. L., 78, 79. Richards, O. W., 361. Richardson, C. H., 278. Richmond, H. A., 354. Richter, B., 182. Ripper, W. E., 254. Risbec, J., 161, 165. Ritcher, P. O., 315. Rivers, I. La, 146, 368. Rivnay, E., 75, 225, 325, 384. Roark, R. C., 57. Robertson, F. W., 74. Robertson, P. L., 77. Robertson, W. H., 153. Robinson, R. H., 152. Rodriguez Lopez, L., 5, 95. Roebuck, A., 92. Romney, V. E., 279, 314. Roonwal, M. L., 136. Rosenberg; H. T., 354, 356. Rosenstiel, R. G., 342. Röszler, P., 16. Roth, R. W., 180. Rothgeb, R. G., 49. Ruiz Castro, A., 167, 298. Ruston, D. F., 87. Ruszkowski, J., 5. Ryan, W. St.-G., 224.

Sailer, R. I., **147.** St. George, R. A., 287, 406. Sakimura, K., 251. Sandstedt, R. M., 271. Santis, L. De, 117, 132. Santoro de Crouzel, I., 401. Sasscer, E. R., 100. Satterthwait, A. F., 219. Sauer, H. F. G., 202, 204. Schaerffenberg, B., 10. Schechter, M. S., 106, 368. Schedl, K. E., 8. Schilder, F. A., 118. Schimitschek, E., 7. Schmidt, E., 5. Schneider, G. W., 68. Schneider-Orelli, O., 194, 379. Schötmar, G. Wolff-, 14. Schulze, B., 228. Schütz, K., 15. Schwan, B., 36. Schwardt, H. H., 179. Schwartz, M., 157. Schweis, G. G., 59. Schwerdtfeger, F., 7, 9, 13, 14.

Schwitzgebel, R. B., 236. Scott, L. B., 344. Searls, E. M., 343. Secrest, H. C., 107. Sellke, K., 157. Semans, F. M., 296. Serebrovskii, A. S., 304. Severin, H. C., **193.** Severin, H. H. P., 292, **378.** Shaw, F. R., 212. Shaw, J. G., 340. Sheals, R. A., 238. Sherbakoff, C. D., 25. Shtakel'berg (Stackelberg), A. A., 328. Shull, W. E., 284. Siegler, E. H., 106, 107, 214, 344. Silva, F. Azevedo e, 310. Simmonds, F. J., 74, 354. Simmonds, H. W., 92. Singleton, H. P., 23. Slade, R., **256.** Slifer, E. H., 386. Smee, C., 85. Smith, C. C., 142. Smith, C. F., 96. Smith, C. L., 213. Smith, C. M., 332. Smith, D. N., 354. Smith, F. F., 1, 2, 38, 39, 43, 70, 174. Smith, H. S., 219. Smith, J. H., 369, 404. Smith, L. E., 214. Smith, L. G., 100. Smith, M. I., 160. Smith jr., R. B., 160. Smith, R. F., 30, 81, 241. Smith, R. H., 352, 373. Snyder, T. E., 176. Solomon, M. E., 368. Sontakay, K. R., 391. Sorenson, C. J., **101.** Speyer, E. R., **404.** Speyer, W., 119. Spiller, D., 93, 369. Springensguth, W., 13. Stackelberg (see Shtakel'berg). Stăl, C., 263. Stanley, W. W., 25. Stapel, C., 12, 380, 381, 382. Staples, R., 206. Stapley, J. H., 297. Stark, V., 306. Starostin, S., 303, 306. Starr, D. F., 216, 340, 345, 379. Steiner, L. F., 218. Stellwaag, F., 13.

Stepniewska, K., 117. Steudel, W., 15. Steyaert, R. L., 364, 365. Stiff jr., H. A., 264. Stitt, L. L., 280. Stoddard, E. M., 378. Stofberg, F. J., 293. Stohlman, E. F., 160. Stone, M. W., 55. Stone, W. E., 395. Strickland, A. H., 129. Stringer, A., 197. Strong, W. J., 64. Stubbs, L. L., 292. Subklew, W., 10. Sukhov, K. S., **160, 183.** Summerville, W. A. T., 268. Swain, R. B., 113. Sweetman, H. L., 56, 106, 272, 340. Swezey, O. H., 264. Swingle, M. C., 279, 344. Sylvén, E., 19. Szelényi, G. von, 122.

Tate, H. D., 25, 216, 271. Tauber, O. E., 175. Taylor, D. J., 116. Taylor, W. L., 404. Teixeira Mendes, C., 375. Tejero, F. Domínguez García-, 166. Telford, H. S., 96. Thalenhorst, W., 10. Thiem, H., 34. Thomas, I., 155. Thompson, R. C., 39. Thompson, R. W., 64, 137. Thompson, W. R., 128. Tiel, D., 11. Tooke, F. G. C., 83. Trägårdh, I., 7. Traill, D., 266. Trehan, K. N., 362, 391, 393. Trouvelot, B., 159. Tuft, P. H., 328. Tunblad, B., 35. Tupikov, V., 304. Turbott, E. G., 93. Turnbull, J., 96. Turner, N., 48. Twinn, C. R., 143.

Ullyett, G. C., 394. Upholt, W. M., 282. Uppal, B. N., 392. Urquijo Landaluze, P., 125. Ushatinskaya, R. S., 330.

van den Bruel, W. E., 120. van Emden, F. I., 296. van Poeteren, N., 12, 157. Vappula, N. A., 406. Veitch, R., 31. Venables, E. P., 334. Verma, P. M., 392. Voelcker, O. J., 91. Vollert, H. E., 157. von Szelényi, G., 122. Vrydagh; J. M., 127, 363, 366.

Wadley, F. M., 313. Wahlen, F. T., 158. Wain, R. L., **197.** Walkden, H. H., **236.** Walker jr., F. H., 82. Walker, H. G., 51, 59, 214. Wallace, P. P., 173. Wallis, R. L., **175.** Walton, R. R., 60, 70. Ward, I. J., 154. Waterston, J. M., 38, 172, 289. Watkins, T. C., 52, 179, 180. Watson, J. R., 116. Webb, J. E., 383. Weber, A. L., 63. Weber, N. A., 147. Webster, R. L., 21, 284, 343. Weddell, J. A., 291. Weil, J. W., 293. Weiss, H. B., 146. Welch, H., 59. Wellenstein, 8.

Wells, W. G., 87. Wene, G., 277. West, T. F., 137. Whelan, D. B., 25. Whitcomb, W. D., 177. Whitehead, F. E., 60, 69, 78, 151, 270. Wiesmann, R., 117, 124, 288. Wigglesworth, V. B., 188, 406. Wilcox, R. B., 401. Wilford, B. H., **53.** Will, H. C., 4. Wille, J. E., **5, 131, 132.** Williams, C. B., 296. Williams, F. X., **290.** Wilson, G. F., 118, 171. Wilson, H. F., 277, 282, 283. Winkler, E., 255. Winteringham, F. P. W., 264. Wishart, G., 138. Wolcott, G. N., 110, 243, 244. Wolfenbarger, D. O., 79, 173, 313, 345. Wolff-Schötmar, G., **14.** Wood, A. A., **140.** Woodard, G., **160.** Wressell, H. B., 139, 140. Wright, J. A., 187. Wylie, W. D., 115.

Yeager, J. F., **64.** Yothers, M. A., **56, 108, 274, 281.** Yuill, J. S., **398.**

Zacher, F., 231, 232. Zappi Recordati, A., 6. Zentmyer, G. A., 173. Zhukovskii, A., 302. Zoheiry, M. S. el, 5, 254. Zukel, J. W., 368.



GENERAL INDEX.

In the case of scientific names, the page reference is cited only under the heading of the generic name.

When a generic name is printed in brackets, it signifies that the name is not the one adopted.

A:

abbreviata, Aphis (see A. rhamni).

abbreviatus, Diaprepes.

abdominalis, Coeloides; Olla.

abdominalis, auct., Aulacophora (see A. foveicollis).

Abies, Chermes spp. on, in Central Europe, 11, 194.

Abies balsamea, pests of, in Canada and U.S.A., **143, 155, 313, 314.**

Abies cephalonica, Chermes prelli on, in Switzerland, 194.

Abies lasiocarpa, Harmologa fumiferana on, in U.S.A., 314.

Abies nordmanniana, Chermes prelli on, in Switzerland, 194.

abietella, Dioryctria.

abietis, Neodiprion. abnormis, Leptomastidea.

Acacia spp., pests of, in S. Africa, 83, 84; Icerya purchasi on, in France, 357; insects on, in India, 385, 391, 393; Coccus hesperidum on, in Palestine, 326; ash of, ineffective against Calandra in maize, 327.

acaciae, Colasposoma.

erythrocephala, Acantholyda bionomics and control of, on pine in Germany, 14. Acanthoscelides (see Bruchus).

Acaphylla steinwedeni, gen. et sp. n., on Camellia in U.S.A., 111.

Acarapis, infesting bees, 6, 328; investigations on forms of, 6. Acarapis woodi, 328.

acasta, Melittobia.

Acaulona peruviana, introduction of, against Dysdercus in Porto Rico, 312.

Acenaphthene, toxicity of, to Cydia pomonella, 215.

Acer (see Maple).

Acer negundo, Leptocoris trivittatus on, in Utah, 106.

aceris, Phenacoccus.

Acerophagus notativentris, utilisation of, against vine mealybug in California, 214. Acerophagus pallidus, bionomics and laboratory utilisation of, in California, 61; exported to Hawaii, 61.

Acetanilide, reducing toxicity of derrsi dust, 78.

Acetic Acid, and flour in sprays, 272. Acetone, as a solvent, 65, 197, 244, 379.

Acetophenone Oxime, action of, on Lepidopterous larvae, 344.

Achaea janata, parasites of, on Ricinus communis in India, 386.

Acherontia atropos, measures against, infesting beehives in Hungary, 6.

Achras sapota, Anastrepha serpentina on, 52. achyloessa, Glyphipteryx. aciculatus, Mastrus.

acinaciformis, Coptotermes.

acraea, Estigmene.

Acridiophaga caridei (see Sarcophaga). Acridotheres tristis, destroying Trichoptilus wahlbergi in Mauritius, 371.

Acrobasis caryae, measures against, on pecan in Texas, 210, 346, 347; cocooning habits of, 346.

Acrolepia assectella, on onion in Hawaii,

Acropyga (Rhizomyrma), revision and new species of, associated with Coccids in tropical America, 147.

Acrylonitrile, in aerosols, 44; in fumigant mixtures, 104.

acuminatus, Coccus (Lecanium).

acupunctatus, Scyphophorus. Adalia bipunctata, predacious on Aphids in Britain, 394; predacious on Saissetia nigra in California, 374.

adamsoni, Porotermes.

Adansonia digitata, Dysdercus spp. on, in Nyasaland, 91.

Adelphocoris lineolatus, on poppy in Hungary, 122; food-plants of, in Minnesota, 29, 207; on lucerne in Russia, 305, 306; bionomics of, 29; measures and experiments against, 29, 207, 305, 306.

Adelphocoris rapidus, measures and experiments against, on crops in U.S.A., 29, 207, 335; bionomics of, 29.

Adelphocoris superbus, on lucerne Utah, 101.

Adhesives, banding with, 320, 321; on trap boards, for catching Psylla pyricola, 22.

Adimonia tanaceti (see Galeruca).

Adonia variegata, parasite of, in Yunnan, 369.

adonidum, Pseudococcus.

Adoretus hirtellus (see A. umbrosus).

Adoretus sinicus, insecticides against, on beans in Hawaii, 217.

Adoretus umbrosus, on cacao in Gold Coast, 66.

Adoretus versutus, on rose in Fiji, 290.

Aegeria exitiosa, DDT against, on peach in New Jersey, 213.

Aegeria mellinipennis, bionomics and control of, on Platanus in U.S.A., 353.

Aegeria tipuliformis, on currant and gooseberry in Ontario, 64.

Aegocera rectilinea, on cassava in Belgian Congo, 368.

Congo, 308.

aegopodii, Cavariella. aenea, Halticoptera.

aeneus, Meligethes.

Aenoplex betulaecola (see Mastrus pilifrons).

Aenoplex carpocapsae (see Mastrus).

Aenoplex plesiotypus (see Mastrus pili-frons).

aequalis, Heterobostrychus; Pimpla (Ephi-

aites).

Aerosols, use of insecticides as, 43, 99, 142, 257, 261; increased concentration of solutions producing, during discharge, 99, 332; preparation of pyrethrum concentrate for use in, 189.

Aetorhinus angulatus (see Blepharidop-

terus).

affinis, Dibrachys; Pempherulus (Pem-

pheres).

Africa, French Equatorial (see Cameroons). Africa, French West, cacao pests in, 195, 196; miscellaneous pests in, 165, 196; parasite of Saissetia nigra in, 374.

Africa, South, cotton pests in, 87, 127; forest pests in, 83, 324; pests of stored products in, 326, 327; pests of vegetables in, 293, 326; beneficial insects in, 83, 107, 324, 374; Chelonus texanus imported into, from Montana, 75; Habrolepis rouxi introduced into California from, 107; plant quarantines in, 146; prohibition of importation of dahlias into Nyasaland from, 85; precautions against introduction of Cydia pomonella into S. Rhodesia from, 127; Pentatomid used as food in, 327.

Afromosia angolensis, resistance of timber of, to termites in N. Rhodesia, 247.

Afzelia, resistance of timber of, termites in N. Rhodesia, 247.

Agathis diversa, liberation of, against Cydia molesta in New Jersey, 105.

Agathis pumila, bionomics and establishment of, against Coleophora laricella in Canada, 143, 155.

Agathis stigmatera, parasite of Diatraea saccharalis in Florida, 260.

Agave fourcroides, pests of, in Mexico, 131.

agavium Ovaticoccus (Gymnococ

agavium, Ovaticoccus (Gymnococcus).
Agelastica alni (Elm Leaf Beetle), tests of
dusts against, in Germany, 231.

Agicide, **212.**

agilis, Diplazon (Syrphoctonus).

Aglaope infausta, measures against, on almond in Spain, 169.

Agrilus anxius (on birch), measures against, in New Brunswick, 154; factors affecting infestation by, in U.S.A., 43, 271; on aspen, 271.

Agrilus bilineatus, defoliation favouring infestation of oak by, in U.S.A., 43.

Agrilus hyperici, establishment of, against Hypericum perforatum in Australia, **321.** Agrilus rubicola, on currant and goose-

berry in Ontario, 64.

Agriotes, in Britain, 122, 123, 265, 294, 405; effect of soil and altitude on abundance of, 265, 405; studies on growth, movements and physiology of, 123, 294; toxicity of soil insecticides to, 393; predacious on Hylemyia floralis, 122.

Agriotes lineatus, in Britain, 265, 294, 405; in Germany, 11; larva of, 11.

Agriotes obscurus, in Britain, 265, 294, 405; in Germany, 11; bait for, 405; larva of, 11.

Agriotes sputator, in Britain, 232, 265, 294, 405; on poppy in Hungary, 122; new fungus infesting, 232.

Agromyza, on tea in Madras, 246.

Agromyza phaseoli, bionomics and control of, on beans in Belgian Congo, **363.**

Agromyza pinguis, hot water treatment of chicory against, in Belgium, 120.
Agromyza strigata, parasite of, on sweet-

potato in Fiji, 290.

Agromyza virens, parasites of, on Parthenium argentatum in California, 104.

Agropyrum repens, Hylemyia coarctata on, in Britain, 297; experiment with Phyllobius pyri and, 92.

Agropyrum trichophorum, Eurygaster integriceps on, in Uzbekistan, 301.

Agrothereutes slossonae, parasite of Gilpinia hercyniae in Quebec, 224.

Agrotis, on peas in Quebec, 223.

Agrotis fennica, plants attacked by, in Br. Columbia, 154.

Agrotis nigricans (see Euxoa).

Agrotis segetum, on melons in Greece, 16; experiment with *Trichogramma* against, in Russian Union, 306.

Agrotis ypsilon, on potato in Argentina, 4. aguti, Praon.

Aircraft, *Popillia japonica* intercepted in, in Canada, **137**; risk of introduction of

INDEX. 419

insects into Hawaii by, 290; use of, for distributing insecticides, 14, 238, 278, 303, 306, 338, 401; calcium cyanamide distributed from, 260.

Alabama, Coleoptera on crops in, 259; pests and disease of ornamental plants

in, 1, 111.

Alabama argillacea (on cotton), in Porto Rico, 97: measures and experiments against, in U.S.A., 57, 210, 260, 261, 289.

albatus, Plagiognathus.

Alberta, pests of field crops in, 98, 144; survey of forest pests in, 354. albicans, Monochaeta.

albifrons, Spilochalcis.

albipes, Technomyrmex (see T. detorquens). Albizzia, Collembola attacking seeds of, in Madras, 246.

Albizzia antunesi, resistance of timber of, to termites in N. Rhodesia, 247.

Albizzia procera, Bostrychids infesting timber of, in India, 371.

albofasciata, Leucopoecila; Pachypasa.

alboguttatus, Automalus.

albovittatus, Ceuthorrhynchus.

Alcohol, in bait for fruit-flies, 396; solvent, 135, 257, 340, 344, 379. Methylated Spirit.)

Alcohol, Sulphated, as wetting agent for

pyrethrum spray, 255.

Aleurites spp., pests of, in Nyasaland, 86, 87: Pseudaonidia trilobitiformis on, in Belgian Congo, 365: Aonidiella aurantii on, in Brazil, 204.

Aleurocanthus woglumi, Eretmocerus serius imported into Mexico against, 261.

Aleurodicus cardini, on Citrus and Citharexylum in Bermuda, 172, 290; measures against, 172.

Alfalfa (see Lucerne).

Alfalfa Butterfly (see Colias eurytheme). Alfalfa Plant Bug (see Adelphocoris lineolatus).

Alfalfa Snout Beetle (see Otiorrhynchus

ligustici).

Alfalfa Weevil (see Hypera variabilis). Algeria, Platyedra gossypiella in, .165; pests of stored grain in, 403.

algira, Parallelia.

alianta, Niphadolepis.

Allium spp., insects and virus diseases of, in U.S.A., 70, 71. (See Chives, Garlic and Onion.)

Allodorus semirugosus, parasite of Pissodes harcyniae in Finland, 12.

Allograpta exotica, parasites of, in Argentina, 117.

Allograpta obliqua, predacious on Cinara in Bermuda, 290.

Allotropa, attempted introduction of undescribed species of, against Pseudococcus comstocki in U.S.A., 209.

Allotropa burrelli, bionomics and establishment of, against Pseudococcus comstocki in U.S.A., 208, 209, 222.

Allotropa convexifrons, parasite of Pseudococcus comstocki in U.S.A., 208.

Allyl Bromide, effect of, as a fumigant, 229.

Allyl Chloride, effect of, as a fumigant,

Allyl Isothiocyanate, soil fumigation with, against Melolontha, 10.

Almond, new mite on, in California, 111; Myllocerus laetivirens on, in India, 388; Bryobia praetiosa on, in Morocco, 356; Capnodis spp. on, in Palestine, 75, 384; Aglaope infausta on, in Spain, 169. alni, Agelastica.

Aloe aculeata, leaf ash of, against Calandra

oryzae, 327.

Alomya debellator, attempted introduction of, into Australia against Oncopera, 322. alopecuri, Dasyneura.

Alophora subcoleoptrata (parasite of Eurygaster), bionomics and suggested utilisation of, in Russian Union, 301, 302.

Alorco, a form of cryolite, 280.

Alpine Ash (see Eucalyptus gigantea).

Alpine Fir (see Abies lasiocarpa).

Alsophila pometaria, rendering trees susceptible to other pests and disease in U.S.A., 43, 173.

Alternaria solani, 173.

alternatus, Nabis.

Althaea rosea (see Hollyhock).

althaeae, Tetranychus (Epitetranychus) (see T. telarius).

Altica (see Haltica). altiscuta, Pachyneuron.

Aluminium Oxide, against pests of stored grain, 157, 319; mode of action of, on insects, 157, 188, 189.

Aluminium Sulphate, as a safener for arsenicals, 62; effects of, in nicotinebentonite sprays, 273, 274.

amabilis, Ipobracon.

Amargo, 67.

ambiguella, Clysiana (Clysia, Conchylis).

ambiguus, Psallus.

Amblygonocarpus obtusangulus, resistance of timber of, to termites in N. Rhodesia, 247.

Amelanchier, Rhagoletis pomonella on, in Maine, 222.

Amelia (see Tortrix).

americana, Periplaneta; Schistocerca.

americanus, Syntomopus. amicus, Sympherobius.

Aminonitro-o-cresol, useless as an insecticide, 198.

Ammonia, and oleic acid, cottonseed oil emulsified with, 364.

Ammonium Magnesium Arsenate, in baits for cutworms, 140.

Ammonium Sulphamate, for destroying pear stumps and suckers against Psylla pyricola, 22, 258.

Amorbia emigratella, on Gliricidia sepium in Hawaii, 250.

Amphimallon (Amphimallus) solstitiale, in grass in Britain, 155.

amphis, Euderus.

Amphitetranychus viennensis (see Tetranychus crataegi).

n-Amyl Carbamate, action of, on insects,

Anabasine Sulphate, aircraft application of formulae containing, against Aphis gossypii, etc., 307.

Anabrus cerciata, potential importance of, in N. America, 114.

Anabrus longipes, potential importance of, in N. America, 114.

Anabrus simplex, in Canada, 114; U.S.A., 60, 113-115, 146, 259, nature of injury caused by, 113-115; natural enemies of, 146, 368; baits for, 60, 259.

Anacardic Acid, repellent to termites,

Anacardium occidentale, treatment of wood with oil from, against termites,

Anachaetopsis tortricis, parasite of Cydia pomonella in California, 240.

Anagrus yawi, sp. n. (parasite of Pycnoderes quadrimaculatus), in Mexico, 251; establishment of, in Hawaii, 251.

Anagyrus, attempted introduction against Pseudococcus comstocki in Virginia, 209.

Anagyrus aurantifrons, parasite of Pseudococcus perniciosus in Nyasaland, 86.

Anagyrus beneficians, parasite of Pseudococcus spp. in Nyasaland, 86.

anale, Sinoxylon.

analis, Philothalpus (Xenopygus).

Anaphaeis aurota, bionomics of, in Belgian Congo, 368.

Anaphes tingitiphagus, parasite of Corythaica passiflorae in Brazil, 264.

Anarhopus sydneyensis, utilisation of, against Pseudococcus adonidum in California, 59; experiment with P. comstocki and, 109.

Anarsia lineatella, on quince in France,

Anasa tristis, on cucurbits in U.S.A., 21, 179; utilisation of Trichopoda pennipes against, 21; tests of insecticides against, 179.

Anastatus disparis, introduced parasite of Lymantria dispar in U.S.A., 238.

Anastatus semiflavidus, effect of food in rearing of, against Hemileuca oliviae in U.S.A., 215.

Anastrepha, host-fruits, parasites and control ot, in Argentina, 134.

Anastrepha aphelocentema, on Lucuma in Mexico, 397.

Anastrepha chiclayae, on Passiflora in Mexico, 397; in Peru, 397.

Anastrepha distincta, host-fruits of, in Mexico, 397.

Anastrepha fraterculus, host-fruits of forms of, in S. America and Mexico, 134, 397; measures against, 134.

Anastrepha ludens, in Mexico, 52, 53, 190, 216, 340, 395-397; in Texas, 52; bionomics of, 52, 395-397; measures and experiments against, 190, 216, 340,

Anastrepha mombinpraeoptans, on mango in Porto Rico, 52, 397; bionomics of forms of, in Mexico, 52, 53, 397.

Anastrepha pallens (see Pseudodacus). Anastrepha pseudoparallela, on Passiflora

edulis in Argentina, 4.

Anastrepha serpentina, in Brazil, 52; in Mexico, 397; in Panama, 397; in Texas, 52, 397; bionomics of, 52, 397.

Anastrepha striata, investigations on, in Mexico, 395, 396, 397.

anchocelioides, Rhynchagrotis. ancilla, Othreis (Ophideres).

Ancylis comptana (fragariae), on strawberry in U.S.A., 47, 117, 262; parasites and biological control of, 47, 262; insecticides against, 262.

ancylivorus, Macrocentrus. Ancylonycha (see Lachnosterna).

ancylus, Diaspidiotus (Aspidiotus).

andereggiella, Argyresthia. andreae, Dysdercus.

Aneristus ceroplastae, parasite of Saissetia nigra in Australia, Hawaii and Porto Rico, 374.

Anethol, in baits for *Polychrosis viteana*, 27. Angitia apostata, parasite of Enarmonia diniana in Siberia, 401.

Angitia cerophaga, parasite of Plutella maculipennis in New Zealand, 269.

Angitia exareolata, parasite of Enarmonia diniana in Siberia, 401.

Angitia nana, establishment of, against Coleophora laricella in Ontario, 143.

Angitia obliterata, parasite of Polychrosis viteana in U.S.A., 28.

angulatus, Blepharidopterus (Aetorhinus). angulifera, Eublemma.

angusi, Datana.

angusticeps, Thrips.

angustiorana, Batodes (Ditula).

Anisandrus (see Xyleborus).

Anisolabis annulipes, bionomics and control of, on greenhouse radish in Ohio, 67, 68.

Anisota senatoria, tests of insecticides on, in U.S.A., **62**.

Annona cherimolia, Coccid on, in Argentina, 4: Anastrepha ludens on, Mexico, **395**.

Annona reticulata, Anastrepha ludens on. in Mexico, 395.

Annona squamosa, Coccid on, in S. Rhodesia, 126.

annulipes, Anisolabis (Euborellia).

Anobium punctatum (in timber), in Germany, 18; in New Zealand, 369; in Spain, 167; bionomics of, 18, 19, 167, 168, 369; larva of, 167.

Anodontonyx noxius (tetricus, auct.), on

cereals in Australia, 161.

Anogeissus latifolia, Sinoxylon in timber of, in India, 72.

bionomics of, in Anomala horticola, Britain, 155, 156; in Denmark, 381; resistance of, to derris, 381.

Anomis leona, on cacao in Gold Coast, 66. Anomis texana, on cotton in Peru, 132.

Anopheles quadrimaculatus, 160.

Anoplognathus, plants attacked by, in N. S. Wales, 162.

anselli, Tragocephala (see T. guerini). Antestia lineaticollis, parasite of, coffee in S. Rhodesia, 126.

Anthemis cotula, Diabrotica undecimpunctata on, in California, 30.

Anthicus troilus, n.n. (elegans, Lea), 330. Anthocoris nemorum, predacious on Paratetranychus pilosus in Britain, 234.

Anthonomus cinctus (see A. piri).

Anthonomus grandis (on cotton in U.S.A.), effect on yield of different rates of infestation by, 80; natural enemies and biological control of, 70, 122, 261; dusts against, 57, 80, 211, 260, 335, 336.

Anthonomus musculus, bionomics and control of, on blueberry in U.S.A.,

401, 402.

Anthonomus piri (on pear), in Germany, 117; sprays against, in Switzerland, 125. Anthonomus pomorum (on apple), in Britain, 171, 257, 359; in France, 331; in Germany, 35; measures and experi-

ments against, 35, 171, 257, 331, 359. Anthonomus rectirostris, on cherry in

Denmark, 382.

Anthonomus rubi, resistance of, to derris dusts in Denmark, 381.

Anthonomus signatus, on strawberry in Ohio, 263.

Anthonomus vestitus, on cotton in Peru, 132.

Anthrenocerus australis, bionomics and control of, in W. Australia, 163.

anthrenoides, Eucnocerus (see Trogoderma ornatum).

Anthrenus, differentiation of German species of, 230.

Anthrenus fasciatus, bionomics and control of, in India, 72, 73.

Anthrenus fuscus, in birds' nests in Germany, 230.

Anthrenus museorum, scarce in houses in Germany, 229, 230.

Anthrenus pimpinellae, scarce in houses in Germany, 229, 230; in horse-hair stuffing in Tanganyika, 85.

Anthrenus scrophulariae, bionomics of.

in Germany, 229, 230.

Anthrenus verbasci, in California, 102; bionomics of, in Germany, 229, 230; products infested by, 102, 230.

Antigua, plant pest legislation in, 328. antipoda, Apanteles (see A. ruficrus). antiqua, Hylemyia; Orgyia.

Antirrhinum, effect of aerosols against

pests of, in U.S.A., 44.

Ants, in tobacco seed-beds in Queensland, 369; associated with Cerosipha forbesi, 263; associated with Coccids, 65, 66, 147, 214, 366; mites spread by, 299; destroying other ants, 16: destroying other insects, 16, 28, 65, 122, 177, 243, 246; attempted use of, against Sahlbergella singularis, 195; termites working in wood occupied by, 69; measures and experiments against, 231, 257, 288, 318, 322, 339, 343, 370,

Ants, Funnel (see Aphaenogaster longi-

ceps).

Ants, Red Forest (see Formica rufa). Anua coronata, bionomics of, on Quis-

qualis in India, 386.

Anuraphis bakeri, on clover in Oregon,

Anuraphis eriophori, bionomics of, on Viburnum opulus in Britain, 382.

Anuraphis maidiradicis, on maize in Iowa, 101.

Anuraphis marutae, transmitting mosaic disease of cineraria in Washington, 70. Anuraphis piri, sprays against, on pear in Sweden, 36.

Anuraphis roseus, effect of DDT on, on apple in Indiana, 218.

Anuraphis schwartzi, natural enemies of, on peach in Argentina, 117.

anxia, Lachnosterna (Phyllophaga).

anxius, Agrilus.

Anysis saissetiae, parasite of Saissetia

nigra in Philippines, 374.

Aonidiella aurantii (on Citrus), in Australia, 32, 92, 322; in Brazil, 204; in California, 107, 113, 204, 253, 259, 271, 342, 346, 349, 350, 375; in China, 375; on rose, 107, 204, 205; other food-plants and races of, 204, 205; parasites and biological control of, 32, 92, 93, 107, 322, 375; predacious enemies and diseases of, 205; fumigation against, 204, 259, 271, 346; sprays against, 113, 204, 253, 342, 346, 349, 350; treatments favouring infestation by, 204, 205.

Aonidiella citrina, parasite of, in Asia,

Aonidiella eremocitri, Comperiella bifasciata reared from, 374.

Aonidiella orientalis, parasite of, in India, 374.

perniciosa (see Quadraspi-Aonidiella diotus).

Aonidiella taxus, parasite of, in Japan,

aonidum, auct., Chrysomphalus (see C.

Apanteles, parasite of Lepidoptera in Fiji, 202, 290.

Apanteles antipoda (see A. ruficrus). Apanteles carpatus (parasite of Tineola biselliella), in Australia, 320; in New York, 352.

Apanteles diatraeae (parasite of Diatraea distribution, biology spp.), attempted utilisation of, 176-177.

Apanteles dignus, parasite of Keiferia lycopersicella in Hawaii, 250.

Apanteles earterus, parasite of Diparopsis castanea in Nyasaland, 85.

Apanteles expulsus, in Fiji, 290, 291; parasite of Plusia chalcites, 290; recorded as Apanteles ruficrus, 291.

Apanteles flaviconchae, bionomics of, parasitising Colias in California, 81, 82.

Apanteles glomeratus, establishment of, against Pieris rapae in Australia, 322; parasite of P. rapae in New Zealand, 247.

Apanteles leptothecus, parasite of Eublemma costimacula in Nyasaland, 86.

Apanteles marginiventris, establishment of, against Laphygma exempta Hawaii, 250.

Apanteles parasae, parasite of Parasa lepida in Ceylon, 300.

Apanteles rubecula, introduction of, into Australia against Pieris rapae, 322.

Apanteles ruficrus, A. expulsus recorded as, in Fiji, 291.

Apanteles spurius, parasite of Ocnogyna in Morocco, 170.

Aphaenogaster longiceps, nesting habits and control of, in tobacco seed-beds in Queensland, 369.

Aphanotus destructor, in California, 102; bionomics and potential importance of, in stored products, 102, 156.

Aphanus sordidus, attacking stored ground-nuts in Nigeria, 370.

Aphelinus, parasite of Unaspis citri in N. S. Wales, 248.

Aphelinus mali (parasite of Eriosoma lanigerum), establishment and winter storage of, in Britain, 361; establishment of, in California, 282; imported into Mexico, 261.

aphelocentema, Anastrepha.

Aphidencyrtus aphidivorus, parasite of Anuraphis schwartzi in Argentina, 117.

Aphidiinae, of N. America, 96.

Aphidius (Lysiphlebus), parasite of Aphis gossypii in Oklahoma, 60; scope and subgenera of, 96.

Aphidius avenae, bionomics of, in Britain, 394.

Aphidius granarius, bionomics of, in

Britain, 77, 78, 394. Aphidius nigriteleus, sp. n., parasite of

Myzus persicae in U.S.A., 96. Aphidius platensis, parasite of Toxoptera

aurantii in Argentina, 4, 310; bionomics of, 311. Aphidius rosaphidis, sp. n., hosts of, in

U.S.A., 96. Aphidius testaceipes, parasite of Aphis

gossypii in Bermuda, 172. aphidivorus, Aphidencyrtus.

Aphids, and plant virus diseases, 1, 2, 33, 34, 38, 39, 40, 66, 70, 174, 247, 292, 323, 365, 369, 378, 392; comparison of injury to apple by Coccids and, 332; ants associated with, 263; natural enemies of, 4, 6, 60, 77, 86, 88, 96, 100, 106, 133, 172, 181, 194, 206, 225, 263, 290, 310, 359, 361, 365, 389, 394; tests of insecticides on, 15, 65, 179, 276, 277, 288, 328; index to food-plant catalogue of, 406.

Aphis abbreviata (see A. rhamni).

Aphis avenae, auct. (see Rhopalosiphum prunifoliae).

Aphis bakeri (see Anuraphis).

Aphis fabae,* bionomics of, on Viburnum in Britain, 382; on poppy in Hungary, 122; relation of, to virus diseases o lily and tulip in U.S.A., 1, 38.

Aphis ferruginea-striata (on carrots), in California, 338; in Hawaii, 250.

Aphis forbesi (see Cerosipha).

Aphis gossypii, in Bermuda, 172; Hawaii, 251; in Peru, 132; in Russia, 306, 307; in U.S.A., 1, 2, 38, 44, 60, 174, 261, 335, 336; on cotton, 132, 261, 306, 307, 335, 336; on other plants, 44, 60, 172, 251, 261; transmitting virus diseases of ornamental plants, 1, 2, 38, 174; natural enemies of, 60, 172; insecticides against, 44, 60, 172, 307, 336; arsenates favouring infestation of cotton by, 335, 336.

Aphis maidis, on vegetables in Hawaii, 251; on maize in Iowa, 101; on wheat in Tanganyika, 85.

* In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonomy.

Aphis marutae (see Anuraphis).

Aphis medicaginis,* food-plants of, in Hawaii, 251; on cotton in Louisiana.

Aphis middletoni, on papaya in Hawaii, 251.

Aphis papaveris,* on melons in Greece, 16. Aphis pomi (on apple), comparison of injury by Coccids and, in Br. Columbia, 332; measures against, in Holland, effect of DDT on, in Indiana, 218: use of eggs of, for testing ovicides, 15, 328.

Aphis rhamni (on potato), survey of populations of, in Britain, 358; measures against, transmitting leaf-roll in U.S.A., winter food-plants of, 358, 359;

natural enemies of, 359.

Aphis rumicis,* food-plants of, in Hawaii. tests of insecticides on, 54, 65, 276.

Aphis sacchari, food-plants of, in Hawaii,

Aphis viburni,* bionomics of, on Viburnum opulus in Britain, 382, 383.

Aphis, Bean Root (see Trifidaphis radi-

cicola). (see Toxoptera Aphis, aurantii).

Aphis, Cabbage (see Brevicoryne brassicae). Aphis, Cherry (see Myzus cerasi).

Aphis, Cotton (see Aphis gossypii).

Aphis, Mealy Plum (see Hyalopterus arundinis).

Aphis, Melon (see Aphis gossypii).

Aphis, Pea (see Macrosiphum onobrychis). Aphis, Woolly Apple (see Eriosoma lanigerum)

Yellow Sugar-cane (see Sipha Aphis,

flava).

Aphodius howitti, on grasses in N. S. Wales, 161.

Aphrophora parallela, A. saratogensis confused with, 107.

Aphrophora saratogensis, bionomics and control of, on pines, etc., in U.S.A., 107, 108.

Aphthona euphorbiae (virescens), dusts against, on flax in France, 331.

Aphycus flavus (see Metaphycus).

Aphytis diaspidis, parasite of Aulacaspis cinnamomi in Egypt, 228.

apicalis, Faustinus.

apicipennis, Exora.

Apion simplex, associated with apples in Argentina, 132.

Apion ulicis, utilisation of, against Ulex europaeus in New Zealand, 269.

abostata, Angitia.

Apple, Cydia pomonella on, in S. Africa,

127; pests of, in Argentina, 131, 132; Psylla mali on, in Austria, 117; pests of, in Britain, 155, 156, 171, 233, 234, 359, 360, 361; pests of, in Canada, 153, 154, 322, 334, 354, 361; pests of, in Denmark, 381, 382; Taeniocampa gracilis on, in Estonia, 119; pests gracus on, in Estonia, 119; pests of, in France, 125, 331, 354, 355, 356; pests of, in Germany, 34, 119; pests of, in Holland, 253; Quadraspidiotus perniciosus on, in India, 391; Cydia pomonella on, in Italy, 6; Anastrepha spp. on, in Mexico, 395, 397; pests of, in Nyasaland, 86; pests of, in Poland, 117; Curimosphena fasciculosa on, in S. Rhodesia, 126; Cydia pomonella on, in Spain, 125; pests of, in Switzerland, 124, 125, 394; pests of, in U.S.A., 3, 4, 22, 42, 57, 68, 78, 100, 108, 148, 149, 150, 179, 191, 208, 209, 218, 222, 240, 258, 273, 274, 275, 281, 282, 314, 315, 334; comparison of injury to, by Aphids and Coccids, 332; experiment with Thomasiniana theobaldi and, 308; effect of phenothiazine on, 258; repellents in sprays for, 212.

Apple (Fruit), Hepialus humuli in, in gas store in Britain, 359; restrictions on importation of, into Britain, 232; fumigation of, with methyl bromide, used for rearing Polychrosis viteana, 27; in baits for Otiorrhynchus

ligustici, 179.

Woolly Apple Aphis, (see Eriosoma lanigerum).

Apple Blossom Weevil (see Anthonomus pomorum).

Apple Capsid (see Plaesiocoris rugicollis). Apple Maggot (see Rhagoletis pomonella). Apple Sawfly (see Hoplocampa testudinea). Apple Sucker (see Psylla mali).

approximatus, Kalotermes.

Apricot, attraction of, for Capnodis in Palestine, 384.

aprobola, Argyroploce.

Aprostocetus roseveari, parasite of Phytolyma lata in Uganda, 329.

Arabia, Schistocerca gregaria in, 228. Arachis hypogaea (see Ground-nuts).

Avaecerus fasciculatus, inert dust ineffective against, in stored cacao in Gold Coast, 66.

Araucaria cunninghami, insects associated with injury to, in Queensland, 32.

arbustorum, Plagiognathus.

Archips fumiferana (see Harmologa).

Archytas, parasite of Heliothis virescens in Peru, **278.**

Arctium lappa, Myzus persicae on, in Hawaii, **251**.

^{*} In view of the divergent opinions on the identity of the Aphids of this group, no attempt is here made to show the synonomy.

Arescap, 240.

Argemone alba var. glauca, thrips on, in

Hawaii, 251.

Argentina, fruit pests in, 4, 94, 117, 131, 132, 133, 134, 242; miscellaneous pests in, 4, 263, 287; Aphids on sugar-cane and cereals in, 133, 310; Diabrotica of, 242; locusts in, 66, 67, 120, 252, 401; beneficial insects in, 4, 95, 117, 132, 133, 134, 242, 310; beneficial insects imported into U.S.A. from, 261; precautions against introduction of Leucinodes elegantalis into, from Brazil, 94. argentinus, Ephialtes (Calliephialtes). argillacea, Alabama.

Argyresthia andereggiella, on apple in

Switzerland, 394.

Argyresthia cornella, on fruit trees in Switzerland, 125, 394; bionomics of, 394.

argyrogaster, Aulacophora (Rhaphido-

palpa).

argyrolepidus, Proceras. Argyroploce aprobola, bionomics of, in Travancore, 387.

Argyroploce cespitana, on strawberry in Ohio, 263.

Argyroploce variegana, on apple in Switzerland, **394.**

Argyrotaenia citrana (see Tortrix). arida, Empoasca.

Aristotelia disconotella, 47.

Arizona, pests of beet in, 260, 279; Plusia brassicae on lettuce in, 112; Colias eurytheme on lucerne in, 339, 340; Diatraea grandiosella on maize in, 176, 177; grasshoppers in, 341; miscellaneous pests in, 101, 352, 353; beneficial insects in, 112, 176, 177.

Arkansas, Elasmopalpus lignosellus in,

191; Solubea pugnax in, 147.

armigera, Blondelia (Eucelatoria); Heliothis.

armoraciae, Phaedon.

Armyworms (see Cutworms).

Arrhinomyia tragica (see Elodia).

Arsenic, in baits, 140, 142, 154; treatment of timber with solutions containing, against Coleoptera and termites, 84, 247, 299, 371; danger to bees of insecticides containing, 6, 36, 212; problems of accumulation of, in soil, 63, 343; method for sampling deposits of, on apple leaves, 55.

Artichoke, Sphaeroderma rubidum on, in

Spain, 168.

Artocarpus spp., pests of, in Pacific Is., 161, 163.

arundinis, Hyalopterus.

Asamangulia cuspidata, parasites of, on sugar-cane in India, 391.

Ascia monuste, experiments with Mammea americana against, on cauliflower in Porto Rico, 339; susceptibility of, to DDT, 289.

Ascogaster, parasites of, in California and France, 240, 355.

Ascogaster? carpocapsae (quadridentata, auct.), parasite of Cydia pomonella in California, 240; introduction of, into Uruguay against C. molesta, 95; rearing of, on dead Ephestia larvae, 74; considered identical with A. quadridentata,

Ascogaster quadridentata, parasite of Cydia pomonella in France, 354, 355; A. carpocapsae (q.v.) considered identical with, **240**.

Asemum atrum, in fire-killed Pseudotsuga in U.S.A., 40.

Ash, as a repellent against beetles on plants, 17, 88, 304; experiments with, against pests of stored products, 300, 327; as a carrier for dusts, 37.

Asia, Central, cereal pests in, 159, 301, 302, 303; Melolontha spp. in, 159; insects pollinating Taraxacum koksaghyz in, 245; natural enemies of pests in, **301, 328.**

asparagi, Crioceris.

Asparagus, Agrotis fennica on, in Br. Columbia, 154; Aphids on, in Hawaii, 251; Crioceris asparagi on, in Washington, 21.

Aspen, American (see Populus tremuloides).

Aspen, European (see *Populus tremula*). Asphalt, resistance to insects of paper laminated with, 272.

Aspidiotiphagus, parasite of Unaspis citri in N.S. Wales, 248.

Aspidiotiphagus lounsburyi, parasite of Unaspis citri in Argentina, 133.

Aspidiotus, fungus infesting, in Ceylon, 232. Aspidiotus ancylus (see Diaspidiotus).

Aspidiotus combreti zizyphi, on papaya in S. Rhodesia, 126.

Aspidiotus destructor, Chilocorus spp. predacious on, on coconut in W. Indies, 110, 243,

Aspidiotus gracilis, on cacao in Belgian Congo, 365.

Aspidiotus jaboticabae (see Targionia). Aspidiotus juglans-regiae (see Quadraspidiotus).

Aspidiotus perniciosus (see Quadraspidiotus).

Aspidiotus silvaticus (see Selenaspidus). aspidistrae, Pinnaspis.

assectella, Acrolepia.

assimilis. Ceuthorrhynchus; Gryllulus (Gryllus).

Aster, Macrosteles divisus transmitting virus of, to potato in W. Virginia, 173. Aster Leafhopper (see Macrosteles divisus). Asterolecanium pustulans, bionomics and control of, in Egypt, 226, 227; on fig in Nyasaland, 86; synonymy of, 226.

Asterolecanium pustulans sambuci (see A. pustulans).

asynamorus, Tachycines.

Asystasia gangetica, thrips on, in Hawaii,

atalanta, Vanessa (Cynthia).

aterrimus, Carpophilus (see C. hemip-

Athalia himantopus, on crucifers in Nyasaland, 86.

Athalia rosae (spinarum), dusts against, on radish in Denmark, 381.

Athous, populations of, in Britain, 405.

atomaria, Epicauta.

Atractotomus mali, predacious on Paratetranychus pilosus in Britain, 234.

Atrichopogon brevipalpis, sp. n., on cacao in Trinidad, 78.

atriplicis, Trachea.

atropos, Acherontia. atrum, Asemum.

Atta laevigata, experiments with DDT against, in Brazil, 288.

Attagenus pellio, in Germany, 229, 230;

infesting drugs, 230.

Attagenus piceus, in California, 102; in Germany, 229, 230; products infested by, 102, 230; killed by infra-red rays, 53.

attenuata, Psylliodes.

Aulacaspis cinnamomi, parasites and control of, on mango in Egypt, 228; on rose in Fiji. 290.

Aulacaspis cinnamomi mangiferae (see A.

cinnamomi).

Aulacaspis tegalensis (see Chionaspis). Aulacophora abdominalis, auct. (see A. foveicollis).

Aulacophora argyrogaster, plants attacked

by, in New Caledonia, 161. Aulacophora foveicollis, bionomics and control of, in Greece, 16, 17.

aurantifrons, Anagyrus. aurantii, Aonidiella; Scirtothrips; Toxop-

aurata, Cetonia.

auricularia, Forficula.

aurifrons, Eumerus.

aurota, Anaphaeis.

australasiae, Periplaneta.

Australia, Coccids on Citrus, etc., in, 32, 92, 322, 374; Lyctus in, 181; miscellaneous pests in, 269, 322; pests and virus diseases of potato and crucifers in, 33, 93, 322, 323, 378; pests of stored products in, 318, 319, 320, 372; natural enemies and biological control of insects in, 32, 92, 181, 320, 322, 374; biological control of noxious plants in, 321; insects imported into other countries against noxious plants from, 264,

269; termites introduced into New Zealand from, 234. (See also under the various States.)

Australia, South, Austroicetes cruciata in, 76: miscellaneous pests in, 129, 319; establishment of parasite of Aonidiella aurantii on Citrus in, 322.

Australia, Western, Smynthurus viridis in, 129; pests of stored products in, 163,

318.

australicum, Trichogramma.

australis, Anthrenocerus; Teleonemia,

Austria, pests of vines and fruit trees in, 117.

Austrodacus (see Dacus).

Austroicetes cruciata, in S. Australia, 76; physiology of eggs of, 76, 136.

Autoba gayneri, predacious on Asterolecanium pustulans in Egypt, 227.

Autographa brassicae (see Plusia).

Autographa egena, on beans in California,

Automalus alboguttatus, parasite of Dasychira pudibunda in Sweden, 20. avenae, Aphidius; Macrosiphum.

avenae, auct., Aphis (see Rhopalosiphum

prunifoliae)

Avocado, 373; Protopulvinaria longi-valvata on, in Brazil, 205; Pseudococcus adonidum on, in California, 59; Lachnosterna mindanaona on, in Guam, 163; Helopeltis sanguineus on, in S. Rhodesia, 126.

Axonopus compressus, Crambus haytiellus on, in Florida, 115; Coccids on, in Br. Guiana, 147.

azureus, Rhopalicus (Pteromalus).

Azya luteipes, predacious on Aonidiella auvantii in Brazil, 205.

В.

B-1956, **54**, **148**, **149**, **378**.

Babiana, Aphid transmitting virus to Tritonia from, in U.S.A., 2.

Bacillus popilliae, tests and use of, against Popillia japonica in U.S.A., 79, 280.

Bacteria, Anastrepha attracted by culture of, from mango fruit, 396; (infecting insects), 20, 81, 205, 280, 303, 307; utilisation of, 79, 128.

baetica (see boetica).

"Baggo", **283.**

Bagrada hilaris, measures against, on cabbage in S. Africa, 293.

Bahamas, Cryptocephalus marginicollis in, 116.

Baikiaea plurijuga, resistance of timber of, to termites in N. Rhodesia, 247.

Bait-sprays, for Dysdercus, 88; for fruitflies, 132, 162, 172, 190, 256, 363; for Teleonemia australis, 126; for thrips,

52, 97.

Baits, for Anabrus simplex, 59, 259; for ants, 370; for Coleoptera, 55, 73, 83, 179, 259, 345, 347, 348; for crickets, 257, 270; for Ctenolepisma urbana, 344; for earwigs, 68, 141; for Hylemyia antiqua, 154; for locusts and grasshoppers, 60, 144, 201, 257, 259, 321; for moths, 27, 292, 340; for Lepidopterous larvae, 140, 183, 252, 280, 402; for Trypetids, 134, 162, 216, 340, 396; formulae for, 55, 59, 60, 68, 134, 140, 142, 183, 201, 216, 257, 259, 270, 280, 344, 348.

bajulus, Hylotrupes.

bakeri, Anuraphis (Aphis); Pseudococcus (see P. maritimus); Tarsonemus.

Balsam Fir (see Abies balsamea).

balteatus, Syrphus.

Bamboos, insects on, in Porto Rico and Asia, 243, 388; (culms), Coleoptera in, 71, 312; resistance of species of, to Dinoderus minutus, 312.

Bambusa (see Bamboos).

Banana, Cosmopolites sordidus on, Belgian Congo, 365; Pseudococcus maritimus on, in Egypt, 227; C. sordidus on, in Honduras, 261; Pentalonia nigronervosa on, in India, 392; pests of, in Pacific Is., 161, 163, 201, 202; Dacus musae on, in Queensland, 162; Coccids on, in Venezuela, 147; stems of, as traps for C. sordidus, 365.

bancrofti, Tarsonemus.

Bands, for Cydia pomonella, 57, 69, 240, 274, 275, 316, 355; used in tests of repellents for C. pomonella, 274; for other Lepidopterous larvae, 14, 169, **347.** (See Adhesives.)

banian, Hieroglyphus.

Barathra brassicae (see Mamestra).

barbatus, Encyrtus.

Baricide, 280.

Baris laticollis, on swedes in Denmark,

Barium Carbonate, ineffective in baits for Heliothis virescens, 280; toxicity of derris dust to Musca domestica, 78.

Barium Fluosilicate, 280; dusting with, 192, 246, 333; spraying with, 17, 176, 185; mixed with stored cowpeas against Bruchus chinensis, 327; causing injury to potato, 185.

Bark-beetles, relation of, to Ceratostomella ulmi, 31, 43, 110, 173; fungi associated with, in *Pinus echinata*, 286,

Barley, pests of, in Britain, 297, 307; Cephus pygmaeus on, in Denmark, 382; Zabrus tenebrioides on, in Hungary, 11; Laphygma exempta on, in S. Rhodesia, 126; pests of, in U.S.A., 70, 101; susceptibility of varieties of, to Toxoptera graminum, 70; relation of wire-worms to, 260, 265, 294; dinitro compound not injurious to, 54.

Barley (Stored), DDT against Calandra oryzae in, 288; (meal), unsatisfactory carrier for cubé, 283.

Barriers, against Blissus leucopterus, 57. Barringtonia asiatica, Aspidiotus des-tructor on, in Mona I., 110.

basalis, Ceresa; Microphanurus.

Basella alba, mites on, in Ceylon, 299. basilare, Xylobiops.

Bassus [Braconid] (see Agathis).

Bathyplectes curculionis, parasite of Hypera variabilis in California, 41.

Batodes angustiorana, on Taxus in Britain, 171.

Batrachedra, parasites and control of, on pineapple in Porto Rico, 97.

Batrachedra mathesoni, on coconut Florida, 97.

Bats, destroying noxious insects, 112, 333.

Bean Beetle, Mexican (see Epilachna varivestis).

Bean Root Aphis (see Trifidaphis radicicola).

Bean Weevil (see Sitona lineatus).

Beans, Epilachna varivestis on, in Canada, 139, 144; pests of, in Fiji, 290; Tipula czizeki on, in Germany, 182; Lachnosterna mindanaona on, in Guam, 163; pests of, in Hawaii, 217, 250, 251; pests of, in Holland, 12; larvae in nodules of, in Italy, 120; pests of, in U.S.A., 3, 22, 30, 48, 49, 79, 175, 191, 337, 338; relations of wireworms to, 266, 294, 337; insecticides causing injury to, 176, 217.

Beans (Stored), pests in, and their control, 3, 290, 371.

Beans, Broad (see Broad Beans).

Beans, Lima (see Lima Beans). beckii, Lepidosaphes (Mytilococcus).

Bed-bugs (see Cimex lectularius).

Beech, Cryptococcus fagi on, in Canada, 155, 353; Dasychira pudibunda on, in Sweden, 19; experiment with Malacosoma neustria and, 119.

Beech Scale (see Cryptococcus fagi). Beer, in bait for fruit-flies, 134.

Bees, Bumble (see Bombus).

Bees, Honey, pollinating Taraxacum koksaghyz in Russian Union, 245; honey obtained from Aphid honeydew by, 6: natural enemies and diseases of, 5, 6, 328; Leptocoris trivittatus associated with, 106; risk of poisoning, by insecticides, 6, 36, 212, 213; repellents for, in lead-arsenate sprays, 212; measures to eradicate colony of, 345; book on, 328.

INDEX. 427

Beet, Pemphigus betae on, in Alberta, 144; favouring damage by wireworms in Britain, 294; thrips on, in Denmark, 381; Nezara on, in Fiji, 291; Myzus persicae on, in Hawaii, 251; Cleonus punctiventris on, in Russia, 166, 302; pests of, in Spain, 120, 166; pests of, in U.S.A., 23, 63, 101, 206, 260, 279, 338; Eutettix tenellus transmitting curly-top of, 101; (dried), insects infesting, in California, 102.

Belgium, Agromyzids on chicory in, 120; Leptinotarsa decemlineata in, 157, 158.

Bemisia, relation of, to virus diseases of plants in Africa, 126, 330.

Bemisia gossypiperda (see B. tabaci),

Bemisia rhodesiaensis, measures against, transmitting virus disease of tobacco in S. Rhodesia, 126.

Bemisia tabaci, bionomics of, on cotton in Punjab, 391, 392.

benedettoi, Empoasca.

beneficians, Anagyrus.

Bengal, miscellaneous pests in, 128, 363, 393.

Bentonite, nicotine fixed on, 23, 28, 43, 149, 150, 239, 272, 273, 274; other uses and effects of, in sprays and dusts, 152, 180, 211, 276, 282.

Benzene, as a solvent, 239, 244.

Benzene Hexachloride, isomers and value of, as an insecticide, 256-258, 331; toxicity of, to vertebrates, 257, 258; derivatives of, 331.

berberidis, Eulecanium (see E. persicae).

bergrothi, Helopeltis. berlesei, Prospaltella.

Bermuda, miscellaneous pests in, 38, 172, 289, 290; beneficial insects in, 172, 289, 290.

Bermuda Cedar (see Juniperus bermudiana).

Bermuda Grass (see Cynodon dactylon). berwicki, Acropyga (Rhizomyrma).

betae, Pemphigus (Prociphilus).

Betula (see Birch).

betulaecola, Aenoplex (see Mastrus pilifrons).

bicolor, Tenthecoris.

Bidens pilosa, experiment with Ecpantheria hambletoni and, in Brazil, 204.

bidentatus, Ips (Pityogenes).

bifasciata, Comperiella.

bifasciculatus, Chrysomphalus.

Bilberry (see Vaccinium myrtillus).

bilineatus, Agrilus.

bimaculatus, Tetranychus.

binotalis, Crocidolomia.

bioculatus, Paratetranychus (Tetranychus). Biological Control, importance of diapause in parasitic Hymenoptera in relation to, 105; diseases of insects and, 128; book on, in Stain, 357; of noxious plants, 32, 269, 291, 321.

Biosteres (see Opius). biplaga, Earias. biplagiata, Coelophora, bipunctata, Adalia, bipunctatus, Collops.

bipustulatus, Corymbites (Calambus).

Birch (Betula), Geometrid on, in Britain, 360; pests of, in Canada, 154, 155; factors affecting infestation of, by Agrilus anxius in U.S.A., 43, 271.

Birch Borer, Bronzed (see Agrilus anxius). Birds, destroying noxious insects, 9, 14, 20, 72, 112, 121, 146, 155, 182, 246, 263, 267, 333, 356, 371, 401, 404; encouragement of, in forests, 404; Coccids possibly spread by, 171, 248.

biselliella, Tineola.

bisoctonotata, Thea.

bispinosus, Dendrothrips.

bituberculatum, Eulecanium,

bivittatum, Trypodendron (see T. lineatum). bivittatus, Melanoplus.

bivulnerus, Chilocorus (see C. stigma).

Black Leaf 10, 50.

Black Leaf 155, 23, 28, 43, 49, 149, 179, 274.

Blackberry, experiments with Thomasiniana theobaldi and, in Britain, 308. blackburni, Chelonus.

Blastophagus (see Myelophilus).

parasite Blastothrix erythrostethus, Coccus hesperidum in Palestine, 326.

germanica, bromide methyl against, in cheese factories in Wisconsin, 343; value of laminated paper wrapping against, 272.

Blennocampa geniculata, dusts against, on strawberry in Denmark, 381.

Blepharidopterus angulatus, predacious on Paratetranychus pilosus in Britain, 233. Blepharoneura, bait attractive to, in Mexico, 216.

Blissus leucopterus, barriers against, in Iowa, **57.**

Blondelia armigera, rearing and utilisation of, against Laphygma exempta in Hawaii, 250.

Blood, Dried, as carrier for cubé, 283.

Blood Albumin, uses of, in sprays, 58, 97, 148, 149; in formula for treating bands against Cydia pomonella, 274.

Blow-lamp, use of, to destroy egg-masses of Lymantria dispar, 309.

Blueberry, Tortrix rosana on, in Holland, 12; Rhagoletis pomonella on, in Nova Scotia, 144; pests of, in U.S.A., 4, 223, 401, 402.

Blueberry Fruitfly (see Rhagoletis pomon-

Boarmia roboraria, on apple and oak in Britain, 359.

boetica, Cosmolyce (Lampides, Polyommatus); Ocnogyna.

boeticus, Carabus rugosus.

Bolivia, Chalcodermus niger in, 312.

bolliana, Gretchena.

Bollworms (see Cotton Bollworms).

Bombax stolzii, Dysdercus spp. on, in Nyasaland, 91.

Bombay, Aphid and virus disease of cardamom in, 392; Tribolium destroying eggs of laboratory host of Trichogramma in, 393.

Bombus, pollinating Taraxacum kok-saghyz in Kazakstan, 245.

Bombyx mori, strains and continuous rearing of, as host for Compsilura concinnata, 324, 325; use of, in insecticide tests, 131, 288, 289.

bondari, Chalcodermus.

Books, insects damaging, in Fiji, 136, 290. Borax, treatment of timber with, against

beetles, 84.

Bordeaux Mixture, against Empoasca fabae, 180, 206, 317; against Epitrix cucumeris, 180, 206; against Gnorimoschema operculella, 185; effect of, on entomogenous fungi, 204; formulae for, 94, 148, 180, 185, 206, 207, 268, 317; formulae containing, 28, 94, 148, 317; and arsenicals, 22, 28, 62, 94; reducing effectiveness of dinitro-o-cresol, 35; and nicotine sulphate, 180, 317; and oil emulsions, 268, 364; as an emulsifier for oils, 58, 148; and sulphur, 317; and zinc sulphate, 364; wetting agent for, 169; and injury to plants, 62, 105, 317.

Boric Acid, treatment of timber with, against beetles, 84.

bosqi, Celatoria.

Bostrychoplites cornutus, on Eucalyptus in Mauritius, 371.

bostrychorum, Coeloides.

Boswellia servata, protection of timber of, against insects in India, 372.

Bothynoderes (see Cleonus).

botrana, Polychrosis.

Boxelder (see Acer negundo).

Boxelder Bug (see Leptocoris trivittatus).

brachialis, Bruchus.

Brachiaria plantaginea, Aonidiella aurantii on, in Brazil, 204.

Brachycolus heraclei, on celery in Hawaii, 251.

Brachymeria, parasite of Anaphaeis aurota in Belgian Congo, 368; parasite of Batrachedra in Porto Rico, 97.

Brachyrrhinus (see Otiorrhynchus).

Brachystegia utilis, resistance of timber of, to termites in N. Rhodesia, 247.

brachytarsus, Macrosiphum.

Bracon, parasite of Asamangulia cuspidata in India, 391.

Bracon gelechiae, parasite of Polychrosis viteana in U.S.A., 28.

Bracon hellulae, parasite of Hellula phidilealis in Brazil, 95.

Bracon kirkpatricki, attempted spread of, against Platyedra gossypiella in Belgian Congo, **367**.

Bracon mellitor, parasite of weevils in U.S.A., 70, 216; effect of population density on productivity of, 70.

Bracon punctatus, parasite of Cylas formicarius in Louisiana, 216.

Bracon pygmaeus, parasite of Coleophora laricella in Ontario, 143.

Bracon vestiticida, imported into U.S.A. against Anthonomus grandis, 261.

Bracon vulgaris, imported into against Platyedra gossypiella, 261.

Bran (including middlings and pollard), in baits, 59, 60, 68, 134, 140, 142, 179, 183, 201, 257, 259, 270, 402; ineffective as dust against Calandra granaria, 231; feeding experiments with insects and, 235, 320.

Eutinobothrus brasiliensis, (Gastero-

cercodes).

Brassica campestris, Myzus persicae on, in U.S.A., **2.**

Brassica pekinensis, Aphids on, in Hawaii,

Brassica perveridis, Myzus persicae on, in Porto Rico, 339.

brassicae, Brevicoryne; Hylemyia; Mamestra (Barathra); Microplitis; Pieris; Plusia (Autographa).

Brazil, pests of cereals and grasses in, 4, 93, 147, 205; coffee pests in, 147, 203, 311; cotton pests in, 4, 93, 202, 204, 278, 311; fruit pests in, 52, 94, 204, 205; miscellaneous pests in, 4, 13, 95, 205, 401; pests of ornamental plants in, 134, 204, 205; pests of solanaceous plants in, 94, 263, 264, 287, 288, 289; pests of stored products in, 375: Bruchophagus gibbus intercepted in lucerne seed in, 375; beneficial insects in, 95, 133, 204, 205, 264, 278, 311; beneficial insects imported into other countries from, 261, 312.

brazzai, Sinoxylon. brevicauda, Oecanthus.

Brevicoryne brassicae (Cabbage Aphis), in Australia, 322, 323; in U.S.A., 100, 338; transmitting virus disease of cauliflower and swede, 323; reducing seed production of cabbage, measures against, 322, 338; toxicity of DDT to, 288.

brevipalpis, Atrichopogon.

brevipes, Daihinia; Pseudococcus.

brevis, Kalotermes (Cryptotermes).

Bridelia, Lasiocampids on, in Nyasaland, 86.

British Columbia, forest pests in, 354; miscellaneous pests in, 154.

INDEX. 429

orchard pests in, 142, 153, 332, 334; Depressaria heracleana on parsnip in, 333.

British Isles, cereal pests in, 77, 92, 199, 293, 297, 307, 393; forest pests in, 171, 359, 360, 404; insects damaging grasses in, 92, 155, 156, 297, 359; pests of heather in, 232, 233; hop pests in, 359, 361; orchard pests in, 96, 155, 156, 171, 359, 360, 361; pests of small fruits in, 117, 181, 308, 358, 359, 360, 361; pests of ornamental plants in, 118, 171, 382; pests of stored products in, 135, 181, 232, 233, 234, 359; Lyctus in timber in, pests of vegetable and root crops in, 121, 156, 232, 256, 257, 358, 359, 361, 404, 405; wireworms in, 92, 123, 198, 232, 265, 294, 296, 368, 405; beneficial insects in, 77, 122, 181, 233, 308, 360, 361, 394; biological control of Eriosoma lanigerum in, 361; birds destroying noxious insects in, 404; entomogenous fungi in, 232; legislation against introduction of Leptinotarsa decemlineata into, 232; beneficial insects introduced into other countries from, 143, 269, 322.

Beans (Vicia faba), Anomala horticola on, in Britain, 156; Bruchid in, in France, 298; not colonised by Aphis viburni, 383.

bromeliae, Diaspis.

4-Bromoacetophenone, action of, as an insecticide, 279.

brouni, Kalotermes (Calotermes).

Brown-tail Moth (see Nygmia phaeo-

Bruchophagus gibbus (funebris), intercepted in lucerne seed in Brazil, 375; in Russia, 305; in U.S.A., 101; clover, 101; measures against, on lucerne, 101, 305; review of data on,

Bruchus, fumigation against, in stored

beans in Mauritius, 371.

Bruchus brachialis, on vetch in U.S.A.,

101, 259; dust against, 101.

Bruchus chinensis, dusts against, in stored cowpeas in S. Africa, 327; tests of mercury vapour against eggs of, in India, 164.

Bruchus obtectus, in imported beans in Fiji, 290; on beans in U.S.A., 3; experiments with infra-red rays and, 53.

Bruchus pisorum (pisi), measures against, on peas in U.S.A., 100, 260.

Bruchus rufimanus, in broad beans in France, 298.

brumata, Operophtera (Cheimatobia).

brunnea, Colaspis; Serica.

brunneus, Heterobostrychus; Ptinus (see P. hirtellus).

Brushware, insects damaging, 72.

Bryobia praetiosa, bionomics of, on almond and pear in Morocco, 356.

Bryocoropsis laticollis, bionomics of, on cacao in W. Africa, 65, 66, 196; experiments against, 66.

bubalus, Ceresa.

Buckwheat, Lygus oblineatus on, in apple orchard in Br. Columbia, 153.

Buffalo Treehopper (see Ceresa bubalus). Bufo americanus, destroying insects in U.S.A., 96.

Bufo cognatus, destroying insects in N. Dakota, 96.

Bufo marinus, imported into Florida from Porto Rico, 313.

Building Timbers, Coleoptera in, 18, 176, 228, 229, 255, 287, 403, 406; termites in, 73, 298, 318; *Xylocopa* in, 371–372; survey of data on pests of, 232.

Bulgaria, experiments with Empusa against Calliptamus italicus, 122; Lepidoptera on plum and oak in, 119.

Bumelia spiniflora, Pseudodacus pallens on, in Mexico and Texas, 397.

Bupalus piniarius (Pine Geometrid), factors affecting abundance of, in Germany,

buprestoides, Spondylis. buqueti, Tragocephala guerini.

Burma, Pyrausta coclesalis on Cephalostachyum in, 388.

burrelli, Allotropa.

Burseria simaruba, used in tests of preservatives against termites, 243.

Busseola fusca, on maize and sorghum in E. Africa, 85, 86, 88.

Butane, for producing aerosols, 261. Butyl Carbitol Thiocyanate, 30, 44, 198.

(See Thiocyanates.) n-Butyl Phthalate, as solvent for derris extractives in oil, 349.

2-n-Butyl-pyridine, as a fumigant, 275. Byturus urbanus (tomentosus, auct.), dusts against, on raspberry in Denmark, 381.

C.

Cabbage, pests of, in S. Africa, 293; Myzus persicae on, in Australia, 323; Hellula phidilealis on, in Brazil, 95; pests of, in Britain, 121, 358; pests of, in Canada, 139, 154; Pieris brassicae on, in Denmark, 381; pests of, in Fiji, 290, 291; Tipula czizeki on, in Germany, 182; Ceuthorrhynchus rapae on, in Holland, 12; Lepidoptera on, in New Zealand, 247, 376; pests of, in Russia, 304; Hellula undalis on, in S. Rhodesia, 126; pests of, in U.S.A., 22, 24, 48, 49, 100, 177, 178, 220, 276, 338, 348; creolin emulsion causing

scorching of, 304; (dried), insects in, 102, 290.

Cabbage, Chinese (see Brassica pekinen-

Cabbage Aphis (see Brevicoryne brassicae). Cabbage Looper (see *Plusia brassicae*).

Cabbage Maggot (see Hylemyia brassicae). Cabbage Moths (see Mamestra brassicae and Plutella maculipennis).

Cabbage Seed-pod Weevil (see Ceutho-

rrhynchus assimilis). Cacao, pests of, in W. Africa, 65, 66, 195, 196, 364, 365, 366, 383; Chalcodermus marshalli on, in Brazil, 312; pests of, in Trinidad, 147; Ceratopogonids and pollination of, 78; insects and swollenshoot disease of, 66, 196.

Cacao (Stored), pests in, 66.

Cacao Moth (see Ephestia elutella).

Cacoecia (see Tortrix).

cacti, Chilocorus.

Cadelle (see Tenebroides mauritanicus).

cahirensis, Haplothrips.

Cajanus cajan (indicus), pests of, in E. Africa, 86, 88, 89; Saissetia coffeae on, in Porto Rico, 243.

Calambus bipustulatus (see Corymbites).

Calandra, measures against, in stored

grain in Algeria, 403.

Calandra granaria (Grain Weevil), in Australia, 129, 318, 319, 320, 372; in Germany, 228; experimentally infesting dehydrated vegetables, 102; fumigants against, 26, 229, 230, 295; gammexane against, 257; inert dusts against, 156, 157, 231, 319, 372; other

measures against, 228.

Calandra oryzae, associated with fallen apples in Argentina, 132; on cassava in Belgian Congo, 367; experimentally infesting dehydrated vegetables, 102; (in stored cereals), in S. Africa, 327; in Australia 120, 218, 210, 220, 276; in Australia, 129, 318, 319, 320, 372; in Brazil, 375; in Fiji, 290; in India, 165; in Peru, 131; in Porto Rico, 312; in S. Rhodesia, 127; in U.S.A., 25, 69, 236; bionomics of strains of, in stored wheat and maize, 361; DDT against, 288; inert dusts against, 131, 157, 228, 312, 319, 327, 372; fumigants against, 25, 69, 70, 127, 236, 312, 319, 335; effect of mercury vapour on eggs of, 165.

calceata, Lachnosterna (Phyllophaga). Calcid, 255, 256.

Calcite (see Calcium Carbonate).

Calcium Arsenate, against cabbage caterpillars, 293, 337; against cotton pests, 80, 202, 210, 211, 261, 306, 335, 336; effect of, on infestation of cotton by Aphis gossypii, 335, 336; against pests of leguminous plants, 3, 101, 176, 192: against Bruchid in stored cowpeas, 327; against pests of solanaceous plants, 22, 51, 59, 154, 180, 185, 260; against pests of other crops, 4, 28, 63, 64, 112, 168, 191; tests of toxicity of, to Drosophila, 144; formulae for, in dusts, 3, 22, 63, 64, 154, 185, 293, 336; carriers for, 3, 59, 144, 185, 293; spraying with, 22, 28, 168, 176, 180, 185, 191, 260, 306; adhesives for spray or wash containing, 202; atomised sprays of, 51, 254; applied by aircraft, 306; and Bordeaux Mixture, 22, 28, 62; and nicotine, 336, 337; and Paris green, 59, 63, 279; and pyre-thrum, 64; and rotenone, 336; and sulphur, 63, 64, 101, 261, 336; and injury to plants, 4, 22, 62, 105, 176; "safeners" for, 62, 176, 191, 211; soluble arsenic and copper in dust mixtures containing fungicides and, 221; forms of, 335, 336; other insecticides compared with, 176, 279, 335, 336.

Calcium Arsenite, aeroplane dusting with, against pests, 303, 304, 401; in baits for cutworms, 140.

Calcium Carbonate (Calcite, Limestone), tests with, against beetles in stored wheat, 372; benzene hexachloride not decomposed by, 257; in DDT spray, 213: reducing toxicity of derris dust, 78; reaction of lead arsenate with, in spray water, 39.

Calcium Chloride, not increasing efficiency of cutworm baits, 140; effect of, in oil sprays against Tetranychus telarius, 404. Calcium Cyanamide, tests and uses of, for

insect control, 12, 158, 260.

Calcium Cyanide, 44, 345; dusting with, against Anasa tristis, 21; tent fumigation with, against Chrysomphalus ficus, 255, 256; ineffective against Rhizoglyphus echinopus in bulbs, 35; soil treatment with, against pests, 109, 337; use of, for estimating populations of Empoasca on cotton, 296; prietary forms of, 35, 255, 256.

Calcium Hydroxide, reducing toxicity of derris dust, 78; reaction of lead arsenate with, in spray water, 39.

Calcium Sulphate, copper arsenicals precipitated on, 278.

Calendra zea-mais (see Calandra oryzae).

Calgreen, 63.

Calidea, on cotton in Tanganyika, 88, 89. Calidea dregii (on cotton), in Belgian Congo, 367; bait-spray for, in Tanganyika, 88.

California, Citrus pests in, 59, 97, 98, 107, 113, 204, 253, 259, 271, 342, 346, 349, 373, 397; pests of forest and shade trees in, 282, 352, 373, 398; lucerne INDEX. 43I

pests in, 40, 81, 241, 280, 338, 339, 340; maize pests in, 30, 337; miscellaneous pests in, 107, 114, 258, 368; orchard pests in, 111, 219, 240, 258, 282, 314; Pseudococcus spp. on avocado and vines in, 59, 214; pests of ornamental plants in, 107, 111, 373; pests of Parthenium argentatum in, 104, 259; pests in stored products and houses in, 25, 30, 102, 106, 343; pests of vegetable and root crops in, 30, 39, 220, 337, 338; insects and virus diseases of lettuce and flax in, 39, 378; wireworms in, 55, 104, 260, 337; beneficial insects and biological control in, 30, 41, 59, 61, 81, 104, 107, 109, 214, 219, 240, **241, 259, 282, 374;** predactious mite in, 397; proposed introduction of Pyralid against Opuntia in, 242; beneficial insects imported into other countries from, 32, 61, 96, 322, 385; Hercothrips fasciatus probably introduced into Hawaii from, 252.

California Red Scale (see Aonidiella

aurantii).

californica, Chrysopa; Coccinella; Lema trilineata; Pristiphora (Gymnonychus); Tomocera.

californicus, Ligyrus (see L. gibbosus); Limonius (Pheletes).

Calivoa limacina (on fruit trees), in Argentina, 131; in Britain, 360; in Denmark, 381; bionomics of, in Washington, 285; on Cydonia japonica, 360; insecticides against, 285, 360, 381; inert dust against, 131.

Calliephialtes (see Ephialtes).

Callimome druparum, timing of seasonal cycles of, on apple in Maine, 42.

Calliptamus italicus, Empusa grylli infesting, in Bulgaria, 122.

Callosobruchus (see Bruchus).

Calluna vulgaris (see Heather).

Calocampa exoleta (see Xylena).

Calocarpum mammosum, Anastrepha serpentina on, 52.

Calochortus, Aphids and virus diseases in, 2, 38,

Calocoris fulvomaculatus, bionomics and control of, on hops and bush fruits in Britain, 361.

Calomel (see Mercurous Chloride).

Calophyllum inophyllum, Toxoptera aurantii on, in Hawaii, 251.

Caloptenus (see Calliptamus).

Calosoma peregrinator, predacious on Plusia brassicae in Arizona, 112.

Calosoma sycophanta (predacious on Lymantria spp.), in Portugal, 310; in Saxony, 8; established in U.S.A., 238; polyphagous habits and rearings of, 8. Calotermes (see Kalotermes).

Calpe emarginata, on Citrus and guava in India, 386.

Calycotome spinosa, Dacus oleae on, in Italy, 118.

Camellia, Pseudococcus maritimus on, in Egypt, 227; new mite on, in U.S.A., 111.

Cameroons (British and French), pests of cacao in, 195, 196; insects on other plants in, 196.

Camphor Oil, proprietary insecticide containing, 51.

camphorae, Xyleborus.

Campoplex tortricidis (see Omorgus).

Campsomeris mansueta, encouragement of, against Cochlictis melolonthoides in

Tanganyika, 85.

Canada, forest pests in, **8**, **180**, **313**, **314**, **353**, **354**, **398**; Lachnosterna spp. in, **140**, **141**, **180**; miscellaneous pests in, **107**, **114**, **220**, **339**; Cydia nigricana on peas in, **141**, **145**; Plodia interpunctella in stored wheat in, **402**; beneficial insects in, **9**, **180**, **353**, **398**; proposed biological control of Cydia pomonella in, **240**, **241**. (See also under the various Provinces.)

canadaria, Melanolophia.

Canary Is., Bruchophagus gibbus intercepted in Brazil from, 375.

Canavalia ensiformis, Rhopalocampta forestan on, in Nyasaland, 86.

candida, Saperda.

Canna, Pseudococcus maritimus on, in Egypt, 227.

Cantaloupe (see Melon).

Cantharis, possibly predacious on Cydia pomonella in France, 356.

canus, Limonius (Pheletes).

Cape Gooseberry (see Physalis peruviana). capensis, Nudaurelia cytherea; Pachypasa. capitata, Ceratitis.

Capitophorus, new parasite of, on strawberry in U.S.A., 96.

Capitophorus fragaefolii, on strawberry in Argentina, 4.

Capitophorus fragariae, Staphylinids attacking, on strawberry in Britain, 181.

Capitophorus potentillae, new parasite of, on rose in U.S.A., 96.

Capitophorus ribis, on currant and gooseberry in Ontario, 64.

Capnodis spp., bionomics of, in Palestine, 75, 384.

capreae, Cavariella.

Caproic Acid, attractiveness of, for Cotinis nitida, 347, 348.

Capsicum, Scaptocoris terginus on, in Argentina, 4; Faustinus cubae on, in Brazil, 288; condition due to mites on, in Ceylon and India, 299, 300; Nezara on, in Fiji, 291; Tmolus echion on, in

Hawaii, 250; Gnorimoschema gudmannella on, in Mexico and Porto Rico, 97; experiments with Mexican species of Anastrepha and, 395, 397.

capsulatus, Stomacoccus.

Capsus meriopterus, predacious on Paratetranychus pilosus in Britain, 233, 234. Capsus ruber (see Deraeocoris).

Carabus rugosus, predacious on Ocnogyna in Morocco, 170.

Carabus rugosus var. boeticus, predacious

on Ocnogyna in Spain, 170.

Carbolic Acid (Phenol), in spray against Bagrada hilaris, 293; unsatisfactory as a bee repellent in sprays, 212; treatment of timber with, against termites, 244.

Carbolineums (see Tar Distillates).

Carbon, Activated, tests with dust of, against pests of stored products, 231; pyrethrum extracts purified with, 189, 190.

Carbon Bisulphide, 345; fumigating ants' nests with, 322; soil fumigation with, against Coleoptera, 17, 56; against Lamiid borers, 365; against pests of stored products, 25, 69, 94, 312, 318, 319, 326, 375; sorption of, by wheat, 237; effects of, on plants and products, 17, 94, 259, 320, 326, 375; ineffective against mites in Sitotroga cultures, 125; vacuum fumigation with, 375.

Carbon Dioxide, in fumigant mixtures,

104, 231, 296.

Carbon Tetrachloride, against pests of stored products, 83, 104, 235, 236, 237, 371; mixed with other fumigants, 83, 104, 235, 236, 237; decreasing viability of wheat, 237; effect of sorption of, by wheat, 236; as a solvent for benzene hexachloride, 257.

carbonaria, Capnodis.

carbonarium, Monomorium.

Carcelia gnava, parasite of Dasychira pudibunda in Sweden, 20.

Cardamom (see Elettaria cardamomum).

cardamomi, Taeniothrips. cardinalis, Dysdercus; Rodolia (Novius). cardini, Aleurodicus (Metaleurodicus).

Cardiochiles nigriceps, parasite of Heliothis vivescens in Virginia, 281.

Cardol, tests with, for treating wood against termites, 244.

Cardolite 627, 349, 350.

Carduus pycnocephalus, Eurygaster integriceps on, in Uzbekistan, 301.

Carex, Dociostaurus maroccanus on, in Iraq, 200.

Carica papaya (see Papaya).

caricis, Vesiculaphis.

caridei, Sarcophaga (Acridiophaga). carinatus, Cheiloneurus; Eriophyes. cariosa, Capnodis.

Carnation, Pseudococcus maritimus on, in Egypt, 227; Epichorista ionephela on, in S. Rhodesia, 126; pests and diseases of, in U.S.A., 174, 379; spray causing injury to, 379.

Carneocephala fulgida, in U.S.A., 104, 172; transmitting virus disease of vines, 172; measures against, on

Parthenium argentatum, **104.**

Carneocephala triguttata, transmitting virus disease of vines in U.S.A., 172.

Carolina, North, Ctenolepisma urbana in, 106; forest pests in, 286; Rhagoletis pomonella on blueberry in, 401.

Carolina, South, Capsid predacious on Pseudococcus comstocki on apple in, 209; miscellaneous pests in, 41, 348.

carpatus, Apanteles.

Carpet Grass (see Axonopus compressus). Carpocapsa (see Cydia).

carpocapsae, Ascogaster; Mastrus (Aenoplex).

Carpophilus aterrimus (see C. hemipterus). Carpophilus decipiens (see C. ligneus).

Carpophilus hemipterus, in Queensland, 31, 291; infesting Citrus fruits injured by moths, 291; measures against, in stored ground-nuts, 31; synonymy of,

Carpophilus ligneus, synonymy of, 330. Carrot, Cavariella aegopodii and virus disease of, in Australia, 292, 369; Aphis ferruginea-striata on, in California, 338; Tipula czizeki on, in Germany, 182; Aphids on, in Hawaii, 250, 251; experiments with Psila rosae and varieties of, in Sweden, promoting rapid growth of Agriotes, 294; (dehydrated), insects infesting, 102.

Carrot Fly (see Psila rosae).

Cartox, 231.

carus, Paratillus.

caryae, Acrobasis.

casei, Piophila; Tyrolichus.

Casein, Anastrepha striata attracted by, partial development of Tineola biselliella on, 351.

Casein-lime, as a spreader for sprays, 248. Cashew (see Anacardium occidentale).

Casimiroa edulis, Anastrepha ludens on, in Mexico, 395.

Cassava, Bemisia transmitting disease of, in E. Africa, 330; Erinnyis ello on, in Brazil, 13; pests of, in Belgian Congo, 367, 368; Lachnosterna mindanaona on, in Guam, 163; Eumerus spp. on, in Hawaii, 250; Ecpantheria

castanea, Diparopsis.

hambletoni fed on, 204. castaneiceps, Spatulifimbria.

castaneum, Nicobium; Tetropium; Tribolium.

Castor, Helopeltis orophila on, in Belgian Congo, 37; pests of, in India, 386; Omocena convergens on, in Nyasaland, 87; Ecpantheria hambletoni fed on, 204; experiments with products of, as insecticides, 106, 149.

Castor Oil, Sulphonated, as an emulsifier,

275.

Casuarina, Icerya purchasi on, in Mona Island, 110.

Cathartus quadricollis, measures against, in stored maize in Porto Rico, 312.

Catorama herbarium, attacking book covers and leather in Fiji, 290.

Cattleya (see Orchids).

caudatus, Ephialtes (Ichneumon).

Cauliflower, Brevicoryne brassicae transmitting virus disease of, in Australia, 323; Prodenia litura on, in Bengal, 363; Hellula phidilealis on, in Brazil, 95; Hylemyia floralis on, in Britain, 121; Scaptomyzella incana on, in Denmark, 381; Rhopalosiphum pseudobrassicae on, in Hawaii, 251; Plutella maculipennis on, in Porto Rico, 339; pests of, in U.S.A., 177, 178, 338.

cautella, Ephestia.

Cavariella aegopodii, transmitting virus disease of carrot in Australia, 292, 369; on willow, 292.

Cavariella capreae, on carrot in Hawaii,

251.

cavus, Dibrachys.

Cedar Oil, solution of, protecting wood from termites, **244.**

Ceiba pentandra, insects on, in W. Africa, 196; Lachnosterna mindanaona on, in Guam, 163; Dysdercus spp. on, in Nyasaland, 91.

Celatoria bosqi, parasite of Diabrotica

speciosa in Argentina, 242.

Celatoria diabroticae, parasite of Diabrotica undecimpunctata in California, 30. Celerio lineata var. livornica, on vines in

Spain, **298.**

Celery, Hylemyia cilicrura on, in California, 338; pests of, in Hawaii, 250, 251.

Cellosolve (see Ethylene Glycol Monoethyl Ether).

cephalonica, Corcyra.

Cephalostachyum pergracile, Pyrausta coclesalis on, in Burma, 388.

Cephus cinctus, relation of, to flax in Canada, 98.

Cephus pygmaeus, on cereals in Denmark, 382.

Ceramica picta, parasite of, 281.

Ceraphron, parasite of Fornicia ceylonica in Ceylon, 300.

cerasi, Myzus.

Ceratitis capitata, 134; in Bermuda, 172; in Egypt, 256; on Momordica bal-

samina in Hawaii, 250; measures against, on Citrus, 172, 256.

Ceratitis rosa, on mango in S. Rhodesia, 126.

ceratoniae, Myelois.

Ceratostomella pini, relation of Dendroctonus frontalis to, on Pinus echinata in U.S.A., 286, 287.

Ceratostomella pseudotsugae, relation of Dendroctonus pseudotsugae to, in Douglas fir in U.S.A., 40.

Ceratostomella ulmi, relation of Scolytids to, in elm in U.S.A., 31, 43, 110, 173; defoliation by insects influencing development of, in elm, 173.

cerciata, Anabrus.

cerealella, Sitotroga.

Cereals, Laphygma exempta ovipositing on, in E. Africa, 245; pests of, in Britain, 77, 199, 393; Agrotis fennica on, in Br. Columbia, 154; Eurygaster integriceps on, in Persia and Iraq, 302; pests and disease of, in Russian Union, 183, 301, 302; pests of, in U.S.A., 57, 101, 114, 147. (See Maize, Wheat, etc.)

Cereals (Stored), pests in, and their control, 25, 69, 102, 127, 129, 131, 164, 224, 228, 229, 230, 232, 236, 257, 259, 312, 318, 319, 320, 327, 330, 361, 362, 372, 393, 402, 403.

Ceresa basalis, bionomics and control of,

on pear in Washington, 285.

Ceresa bubalus (on fruit trees), in Br. Columbia, 153; bionomics of, in Washington, 285; measures against, 153, 285.

Ceromasia senilis, auct. (see Lydella stabulans grisescens).

cerophaga, Angitia.

ceroplastae, Aneristus.

Ceroplastes destructor, spray against, on Citrus in N. S. Wales, 268.

Ceroplastes ficus, parasite of, on Aleurites fordi in Nyasaland, 86.

Ceroplastes floridensis, parasites of, in Palestine, 325, 326.

Ceroplastes rubens, on fruit trees in New Caledonia, 161.

Ceroplastes rusci, parasite of, in Palestine, 326.

Ceroplastes toddaliae, on Annona squamosa in S. Rhodesia, 126.

Cerosipha forbesi, bionomics of, on strawberry in Ohio, 263.

Cerotoma ruficornis, tests of insecticides on, in Porto Rico, 339.

Ceruraphis (see Anuraphis).

cervinatus, Oxycanus.

cespitana, Argyroploce (Olethreutes).

Cetonia aurata, on rye in Denmark, 382. Ceuthorrhynchus albovittatus, on Papaver rhoeas in Hungary, 122. Ceuthorrhynchus assimilis, on cabbage in U.S.A., 22, 24, 100; dusts against, 24.

Ceuthorrhynchus denticulatus, dust against, on Papaver somniferum in Hungary, 122.

Ceuthorrhynchus macula-alba, on Papaver spp. in Hungary, 122.

Ceuthorrhynchus rapae, on cabbage in Holland, 12.

Ceylon, symptoms caused by mites on tomato, etc., in, 299; tea pests in, 294, 300; termites in, 73; beneficial insects in, 300, 385; fungi infesting insects in, 232; Icerya purchasi probably introduced into India from, 385. ceylonica, Fornicia.

Chaetodacus (see Dacus).

Chaff, use of, to protect stored potatoes from *Gnorimoschema operculella*, 300; unsatisfactory in baits for *Oxycanus cervinatus*, 183.

chalcites, Plusia.

Chalcodermus bondari, on cotton in Brazil, 311; possibly a synonym of C. niger, 311.

Chalcodermus marshalli, on cacao and cotton in Brazil, 311, 312.

Chalcodermus niger, C. bondari possibly a synonym of, 311.

chalcographus, Ips (Pityogenes).

chalybea, Haltica (Altica).

Chamaecyparis pisifera, reaction of, to methyl bromide, 50.

Charcoal, action of, on insects, 189. Cheese, pests in, and their control, 77,

Cheese Mite (see Tyrolichus casei).

Cheiloneurus carinatus, parasite of Pseudococcus adonidum in Nyasaland, 86.

Cheiloneurus cyanonotus, parasite of Pseudococcus spp. in Nyasaland, 86.

Cheimatobia brumata (see Operophtera). Cheletogenes ornatus, predacious on other mites in California, 397.

Chelonella (see Chelonus).

Chelonus (parasite of Lepidoptera), in Eritrea, 379; in Fiji, 290; in India, 389.

Chelonus blackburni, parasite of Keiferia lycopersicella in Hawaii, 250.

Chelonus pectinophorae, utilisation of, against Platyedra gossypiella in Texas, 261.

Chelonus phthorimaeae, establishment of, against Gnorimoschema operculella in N. S. Wales, 322.

Chelonus texanus, parasite of Loxostege in Montana, 74; reared and imported into S. Africa in Ephestia, 75.

Cherimoya (see Annona cherimolia).

Chermes nordmannianae (on Abies), in Hungary, 11; bionomics and control of, in Switzerland, 194.

Chermes nüsslini (see C. nordmannianae). Chermes piceae, bionomics and control of, on Abies in Switzerland, 194.

Chermes prelli, on Abies spp. in Switzer-

and, 194.

Cherry, Myzus cerasi on, in Br. Columbia, 153; pests of, in Denmark, 381, 382; Rhagoletis spp. on, in U.S.A., 23, 24, 222; Thomasiniana theobaldi not ovipositing on, 308.

Chierry Aphis (see Myzus cerasi). Chick-pea (see Cicer arietinum).

Chickadee (see Penthestes atricapillus).

chiclayae, Anastrepha.

Chicory, Agromyzids on, in Belgium, 120; thrips on, in Denmark, 381.

Chicory (Ground), development of *Plodia* interpunctella on, 231.

chilensis, Hyalomyia. Chilli (see Capsicum).

Chilo trypetes, on sugar-cane in India, 406. Chilocorus bivulnerus (see C. stigma).

Chilocorus cacti, distribution of, attacking Coccids in W. Indies, 110, 243, 313; imported into Panama, 313.

Chilocorus distigma, predacious on cassava Coccid in Belgian Congo, 368.

Chilocorus nigritus, predacious on Chionaspis tegalensis in Mauritius, 371.

Chilocorus politus, predacious on Chionaspis tegalensis in Mauritius, 371.

Chilocorus stigma, in California, 282, 374; in Canada, 354; predacious on Coccids, 354, 374; predacious on Eriosoma lanigerum, 282.

Chilomenes sexmaculata, not parasitised by

Perilitus coccinellae, 369.

China, hosts of Comperiella bifasciata in, 374; parasite of Coccinellids in, 369. chinensis, Bruchus (Callosobruchus); Estigmena; Parlatoria.

Chionaspis citri (see Unaspis).

Chionaspis furfura, spread of, on apple in New York, 148.

Chionaspis tegalensis, Coccinellids predacious on, on sugar-cane in Mauritius, 370. Chives, Idiopterus formosanus on, in

Hawaii, 251.

Chloratone, and methyl chloride, aerosol of, against Tarsonemus pallidus, 44.

chlorea, Sphingomorpha.

Chloridea obsoleta (see Heliothis armigera). Chlorion laeviventris, bionomics of, destroying Anabrus simplex in Nevada, 146, 368.

chlorizans, Malacocoris.

Chlorochroa sayi, in U.S.A., 63, 101; effect of, on seed production of crops, 101; dusts against, 63.

Chloroclystis, bionomics of, on mango in

Travancore, 387.

Chloroclystis rectangulata, on fruit trees in Switzerland, 125, 394.

INDEX. 435

Chlorophora excelsa, Psyllid on, in Uganda, 329; termites making soil favourable for, 329.

Chlor-2-phenylphenol, in barriers against

Blissus leucopterus, 57.

Chlorpicrin, fumigation with, reducing viability of wheat, 237, 259; (as a soil fumigant), against Lamellicorns, 10, 58; increasing susceptibility of pineapple to mealybug wilt, 249.

Chortoicetes terminifera, outbreaks and biology of, in Australia, 321, 322;

experiments against, 321.

Chrome Alum insects repelled by

Chrome Alum, insects repelled by solution

containing, 272.

Chrysocharis laricinellae, establishment of, against Coleophora laricella in Canada, 143, 155.

Chrysolina (see Chrysomela).

Chrysomela decemlineata (see Leptinotarsa). Chrysomela gemellata, Rossi (see C. quad-

rigemina).

Chrysomela hyperici, establishment of, against Hypericum perforatum in Australia, 321; introduction of, into New Zealand, 269, 321.

Chrysomela quadrigemina, establishment of, against Hypericum perforatum in

Victoria, 321.

chrysomelina, Epilachna.

Chrysomphalus agavis, on Juniperus bernudiana in Bermuda, 290.

Chrysomphalus aonidum, auct. (see C. ficus).

Chrysomphalus bifasciculatus, parasite of, in China, 374.

Chrysomphalus dictyospermi, parasite of,

in China, 374.

Chrysomphalus ficus (on Citrus), in Egypt, 255; in Florida, 211, 341; cultural practices favouring infestation by, 341; measures against, 211, 255.

Chrysopa, predacious on Ormenius infuscata in Bermuda, 289; predacious on Icerya purchasi in Mona I., 110.

Chrysopa californica, predacious on Sais-

setia nigra in California, 374. Chrysopa vulgaris, predacious on Leptinotarsa decemlineata in France, 298.

chrysophthalmus, Meteorus.

Chrysophyllum cainito, Anastrepha ser-

pentina on, 52.

Chrysoplatycerus splendens, not reproducing on Pseudococcus comstocki, 109. chrysorrhoea, auct., Euproctis (Liparis) (see Nygmia phaeorrhoea).

chrysotheme, Colias.

Cicer arietinum, ineffective as trap-crop for Heliothis virescens in Peru, 278.

Cidaria citrata (see Dysstroma).

ciliata, Corythucha.

cilicrura, Hylemyia (Chortophila).

Cimex lectularius, use of heat against, in houses in Germany, 255; experiments with trichloracetonitrile against, 296.

Cinara, natural enemies and importance of, on Juniperus bermudiana in Bermuda, 289, 290; value of honeydew produced by, on conifers in Germany, 6. Cinara thujafolia, on Widdringtonia whytei

in Nyasaland, 86.

Cinchona spp., pests of, in Belgian Congo, 36, 38, 363.

cincta, Illeis.

cinctipennis, Closterocerus.

cinctus, Anthonomus (see A. piri); Cephus. Cineraria, insects and virus diseases of, in Washington, 70.

cinereus, Hesperophanes. cingulata, Rhagoletis.

cinnamomi, Aulacaspis.

circellata, Neokolla.

circumflexus, Myzus.

Cirrospilus pictus, establishment of, against Coleophora laricella in Ontario, 143.

Cistus, Quadraspidiotus perniciosus on, in France, 5.

Cistus monspeliensis, Dacus oleae on, in Italy, 118.

Citharexylum spinosum, Aleurodids on, in Bermuda, 172, 290.

citrana, Tortrix (Argyrotaenia). citrata, Dysstroma (Cidaria).

citri, Contarinia; Paratetranychus; Pseudococcus; Scirtothrips; Unaspis (Chionaspis, Prontaspis).

Citricola Scale (see Coccus pseudomag-

Citrullus vulgaris (see Watermelon).

noliarum).

citricola, Coccidophilus. citrifolii, Dialeurodes. citrina, Aonidiella.

Citrus, pests of, in Argentina, 4, 132, 134; pests of, in Australia, 32, 162, 248, 267, 268, 291, 322; pests of, in Bermuda, 172, 290; Aonidiella aurantii on, in Brazil, 204; pests of, in Belgian Congo, 364, 365; Icerya montserratensis on, in Ecuador, 95; pests of, in Egypt, 228, 255, 256; Coccids on, in France, 5, 357; pests of, in India, 246, 385, 386; new Cecidomyiid on, in Mauritius, 74; Anastrepha ludens on, in Mexico, 52, 340, 395, 396; not infested by Mexican forms of A. fraterculus, 397; Distantiella theobroma on, in Nigeria, 196; pests of, in Pacific Is., 161, 163; pests of, in Palestine, 225, 325, 326; Coccids on, in Porto Rico, 110, 243; pests of, in S. Rhodesia, 126; Lasiocampid on, in

Spain, 170; pests of, in U.S.A.,

59, 97, 98, 107, 113, 204, 211, 253, 259, 271, 341, 342, 346, 349, 350, 373;

fumigation causing injury to, 255, 256;

juice of fruit of, in bait for fruit-flies, 134.

Citrus trifoliata, Quadraspidiotus perniciosus on, in France, 5.

Citrus Aphis, Black (see Toxoptera aurantii).

Citrus Blackfly (see Aleurocanthus wog-lumi).

Citrus Scale, White (see *Unaspis citvi*). Citrus Thrips (see *Scivtothrips citri*).

Cladis nitidula, predacious on Coccids in Porto Rico, 313; imported into Panama,

Porto Rico, 313; imported into Panama, 313.

Cladosporium, infesting Aphids in Belgian Congo, 365; infesting *Icerya purchasi* in

India, 385; associated with thrips injury on pyrethrum in Tanganyika, 85. Cladosporium aphidis, growing on honeydew of Phorodon humuli in New York,

317.

claripalpis, Theresia.

Clausenia purpurea, parasite of Pseudococcus comstocki in U.S.A., 109, 208; parasites of, 109.

clavatus, Prochiloneurus.

Clays, as carriers for dusts, 48, 239, 283; as an emulsifier, 371.

Clemora smithi, factors reducing infestation by, on sugar-cane in Mauritius, 370.

Cleome monophylla, not an important foodplant of Myzus persicae in Nyasaland, 85. Cleonus mendicus, on beet in Spain, 166.

Cleonus punctiventris (on beet), in Russia, 166, 302; in Spain, 166; use of fowls against, 302.

Clinodiplosis, of no importance on raspberry in Britain, 308; occurrence of, among stored seeds, 170.

Closterocerus, parasite of Asamangulia cuspidata in India, 391.

Closterocerus cinctipennis, parasite of Neodiprion spp. in Canada, 9.

Clothes Moths, toxicity of isomers of benzene hexachloride to, **257**, **331**. (See *Tinea pellionella* and *Tineola biselliella*.)

Clover (Trifolium), Smynthurus viridis on, in W. Australia, 129; observations on wireworms and, in Britain, 266, 294; Coleophora spissicornis on, in New Zealand, 163, 184; pests of, in U.S.A., 22, 101, 241; relation of races of Colias chrysotheme to, in N. America, 339, 340.

Clover Butterfly (see Colias chrysotheme). Clover Leaf Weevil (see Hypera punctata). Clysiana (Clysia) ambiguella (on vines), bionomics and control of, in Germany,

bionomics and control of, in Germany, 13,14; in Spain, 298; on Symphoricarpus racemosus, 13.

Cnephasia longana, bionomics of, in Oregon, 342, 343.

Cnephasia wahlbomiana, on flax and peas in Holland, 12.

coarctata, Hylemyia (Leptohylemyia).

coccidarum, Turanodinia.

coccidivorus, Eupelmus.

Coccidophilus citricola, predacious on Aonidiella aurantii in Brazil, 205.

Coccidoxenus, parasite of Ceroplastes ficus in Nyasaland, 86.

Coccinella californica, predacious on Eriosoma lanigerum in California, 282.

Coccinella hieroglyphica, predacious on Lochmaea suturalis in Britain, 233.

Coccinella septempunctata, in Britain, 259, 394; in Czechoslovakia, 252; parasite of, in Yunnan, 369; biology of, in different climatic regions, 225; predacious on Aphids, 225, 359, 394; doubtful value of, 225; device for separating, from trapped insect pests, 252.

coccinellae, Perilitus (Dinocampus).

coccisugus, Hemisarcoptes (see H. malus). Coccophagus (parasite of Coccids), in N.S. Wales, 248; in Palestine, 326.

Coccophagus cowperi, parasite of Saissetia nigra in California, 374.

Coccophagus gurneyi, parasite of Pseudococcus spp. in U.S.A., **59**, **109**; experiments with, **109**.

Coccophagus hawaiiensis, parasite of Saissetia nigra in Hawaii, 374.

Coccophagus isipingoensis, parasite of Saissetia nigra in Transvaal, 374.

Coccophagus lunulatus (see C. scutellaris). Coccophagus modestus, parasite of Saissetia nigra in Dahomey, 374.

Coccophagus ochraceus, parasite of Saissetia nigra in California, 374.

Coccophagus pulvinariae, parasite of Saissetia nigra in California, 374.

Coccophagus rusti, introduced parasite of Saissetia oleae in California, 374; attempted use of, against S. nigra, 374. Coccophagus saissetiae, parasite of Sais-

setia nigra in Panama, 374.

Coccophagus scutellaris (parasite of Coccids), in Palestine, 326; in Porto Rico, 374.

coccorum, Pachyneuron.

Coccus acuminatus, sprays against, on mango in Egypt, 227, 228.

Coccus deltae, on Citrus in Argentina, 4. Coccus elongatus, spray against, on Platanus in U.S.A., 353.

Coccus hesperidum, on Cinchona in Belgian Congo, 36; natural enemies of, on Citrus, etc., in Palestine, 325, 326; spray against, on Platanus in U.S.A., 353.

Coccus pseudomagnoliarum, spray against, on orange in California, 98.

Coccus viridis, on celery in Hawaii, 250.

Coccygomimus instigator (see Pimpla). cochleariae, Phaedon.

Cochliotis melolonthoides, parasite of, on sugar-cane in Tanganyika, 85.

cockerelli, Lecaniobius; Paratrioza.

Cockroaches, measures against, 255, 343; value of laminated paper wrapping against, 272; insecticide tests on 64. 339, 347, 368.

Cocksfoot Grass (see Dactylis glomerata).

coclesalis, Pyrausta.

Coconut, Batrachedra on, in Florida, 97; pests of, in Pacific Is., 161, 163; Aspidiotus destructor on, in W. Indies, 243.

Coconut Oil, Sulphonated, proprietary spray preparation containing, 51.

Coconut Oil Soap (see Soap).

Codiosoma spadix, humidity requirements of, in timber in Germany, 18.

Codling Moth (see Cydia pomonella).

Coeloides abdominalis, hosts of, in Finland, 11, 12.

Coeloides bostrychorum, parasite of Pissodes pini in Finland, 12.

Coeloides melanotus, parasite of Pissodes pini in Finland, 12.

Coelonotus (see Protaphidius).

Coelophora biplagiata, not parasitised by Perilitus coccinellae, 369.

Coelostethus pertinax, humidity requirements of, in timber in Germany, 18.

coerulea, Zicrona.

Coffea (see Coffee). coffeae, Saissetia.

coffearia, Homona.

Coffee, pests of, in W. Africa, 166, 196; pests of, in Brazil, 147, 203, 311; Coccids on, in Colombia, 147; Coccids transmitting phloem necrosis of, in Dutch Guiana, 147; pests of, in Pacific Is., 161, 163; Antestia lineaticollis on, in S. Rhodesia, 126; pests of, in W. Indies, 97, 147.

coffeella, Leucoptera.

coiti, Paratetranychus.

Cola, insects on, in W. Africa, 66, 195, 196. Colaphellus (Colaphus) sophiae, bionomics of, on crucifers in Denmark, 381.

Colaspis brunnea, dusts against, on strawberry in Ohio, 262.

Colasposoma acaciae, bionomics and control of, on Acacia in S. Africa, 83, 84. Colchicum, Aphid and virus diseases in, in

U.S.A., **2.**

Cold Storage, use of, against pests, 3, 25, 191; survival of Aphelinus mali in, 361; effect of, on eggs of Bombyx mori, 324, 325.

Coleophora laricella, parasites and biological control of, on larch in Canada, 142, 143, 155; bionomics of, 142.

Coleophora malivorella, on apple in Delaware, 78.

Coleophora spissicornis, bionomics of, on clover in New Zealand, 163, 184.

Colias chrysotheme (Clover Butterfly), forms and races of, in N. America, 81, 339, 340. (See C. eurytheme.)

Colias eurytheme (Alfalfa Butterfly), natural factors in control of, in N. America, 81, 82, 206, 339, 340; considered a race of C. chrysotheme, 81, 339; effect of interbreeding between C. chrysotheme and, 340.

Collabismodes (see Faustinus).

collaris, Degeeria; Thyraeella (Diadromus). Collops spp., feeding habits of, in Utah, 106.

Colocasia esculenta, Aphids on, in Hawaii, 251.

Colombia, ants and Coccids on coffee in, 147.

Colorado, Anabrus simplex in, 114; pests of forest and shade trees in, 314, 353; pests of vegetables in, 30, 175, 219, 220; Melitara dentata destroying Opuntia macrorrhiza in, 241.

Colorado Potato Beetle (see Leptinotarsa

decemlineata).

comariana, Peronea. combreti, Aspidiotus.

Combretum, Pachypeltis on, in Br. Cameroons, 196.

comes, Erythroneura.

Comperiella bifasciata, hosts and races of, in Asia, 374; utilisation of Chinese race of, against Aonidiella aurantii in Australia, 32, 92, 322; bionomics of, 92, 93, 375.

Compound A, 173.

Compsilura concinnata, rearing of Bombyx mori as host for, in S. Africa, 324; established parasite of Lymantria dispar in U.S.A., 238.

comptana, Ancylis.

Comptonia peregrina, Aphrophora on, in U.S.A., 107.

Comstockaspis (see Quadraspidiotus). comstocki, Euplectrus; Pseudococcus. Conchylis ambiguella (see Clysiana).

concinnata, Compsilura.

concinnus, Phaedon armoraciae. concolor, Nadiasa (Taragama).

condonensis, Kalotermes (Calotermes).

confluens, Neokolla. confraterna, Corythucha.

confusum, Tribolium.

Congo, Belgian, Agromyza phaseoli on beans in, 363; cacao pests in, 364, 365, **366**; pests of cassava in, **367**, **368**; pests of Cinchona in, 36-38, 363, 364; cotton pests in, 127, 128, 366, 367; fruit pests in, 364, 365; Helopeltis orophila on tea in, 37; insects and virus diseases of potato and sweet potato in, 330, 365; locusts in, 366; beneficial insects in, 37, 363, 366, 368; suggested introduction of *Rodolia cardinalis* into, 365.

conigerum, Sinoxylon.

Connecticut, Pseudococcus comstocki and its biological control on apple in, 208, 209, 258; pests of forest and shade trees in, 173, 313, 354; Pyrausta nubilalis on maize in, 151; pests of vegetables in, 48.

Conopia tipuliformis (see Aegeria).

Conorrhynchus (see Cleonus).

Conotrachelus nenuphar (in U.S.A.), on apple, 4, 42, 151; on blueberry, 402; on peach and plum, 314, 315, 317; bionomics of, 315; timing of seasonal cycles of, 42; measures against, 151, 314, 315.

consobrina, Ernestia.

consobrinus, Encyrtus (Microterys).

conspersa, Narosa.

Contarinia, on Juniperus bermudiana in Bermuda, 289; measures against, on lucerne in Russia, 305, 306.

Contarinia citri, sp. n., bionomics of, on Citrus in Mauritius, 74.

Contarinia juniperina, 289.

Contarinia merceri, 170.

Contarinia pyrivora, measures against, on pear in Denmark, 12.

Contarinia tritici, infestation by, as basis for forecasting wheat yield in Britain, 203

convergens, Omocena.

convexifrons, Allotropa.

convolvulii, Myzus (see Macrosiphum solani).

Conwentzia hageni, predacious on Saissetia nigra in California, 374.

cooki, Cremastus.

cophta, Gryllotalpa gryllotalpa.

Copidosoma tortricis, parasite of Peronea

comariana in Britain, 360.

Copidosoma truncatellum (see Litomastix). Copper Arsenate (Basic or Hydroarsenate), against cotton pests, 210, 211, 261, 335, 336; against pests of tobacco and potato, 45, 185, 260, 280, 322, 323; against insects on other plants, 28, 63; tests of toxicity of, to insects, 278, 279; in bait, 280; dusting with, 45, 63, 210, 211, 261, 335, 336; carriers for, 45, 335; spraying with, 28, 45, 185, 322, 323; formulae containing, 45, 63, 185, 211, 278, 280, 323, 335; precipitated on calcium sulphate, 278; and copper hydroarsenite, 278; and soap, 45; and sulphur, 63, 211, 261.

Copper Arsenites, tests of toxicity of dusts of, to insects, **278**, **279**; and basic copper arsenate, precipitated on calcium sul-

phate, 278.

Copper Carbonate, experiments with, against pests in stored grain, 312, 327, 328; and lime, 312.

Copper Naphthenate, solution of, protecting wood from termites, 244.

Copper Oxide, lead arsenate compatible with spray containing, **54**.

Copper Oxychloride, compatibility of insecticides with, 173, 221.

Copper Pentachlorphenate, solutions of, protecting wood from termites, **245**.

Copper Sulphate, 271; colloidal spray of, against *Gnorimoschema operculella*, 185; for treating timber against termites, 244, 247; in calcium arsenate dusts, 3, 144. (See Bordeaux Mixture.)

Copper Sulphate, Tribasic, effect of, in calcium-arsenate dusts, 221, 222.

Copper Sulphide, increasing toxicity of derris dust to Musca domestica, 78.

Coptotermes, characters of introduced species of, in New Zealand, 234.

Corchorus olitorius, parasitism of Pempherulus affinis on, in India, 21.

Corcyra cephalonica, in India, 164, 362, 390, 393; in Mauritius, 370; in Porto Rico, 312; experiments against, in stored grain, 164, 312, 362; Tribolium destroying eggs of, 393; Trichogramma reared on, 370, 393; not parasitised by Rogas testaceus, 390.

Cordyceps militaris, infesting Dasychira pudibunda in Sweden, 20.

coridon, Perichares.

Cork Oak (see Quercus ilex var. suber).

Corn Borer, European (see Pyrausta nubilalis).

Corn Borer, Southwestern (see Diatraea grandiosella).

Corn Earworm (see Heliothis armigera).

cornella, Argyresthia.

corni, Eulecanium (Lecanium).

Cornstalk Borer, Lesser (see Elasmo-

palpus lignosellus).

Cornus spp., Quadraspidiotus perniciosus on, in France, 5; susceptibility of, to Macremphytus varianus in Pennsylvania, 4.

cornutus, Bostrychoplites; Gnathocerus.

corollae, Syrphus. coronata, Anua.

Corrosive Sublimate (see Mercuric Chloride).

ide).
Corylus avellana, Curculio nucum on, in

Denmark, 382; pests of, in U.S.A., 341. Corymbites spp., effect of altitude on populations of, in Wales, 405.

Corythaica, food-plants, synonymy and distribution of species of, 263.

Corythaica monacha (cyathicollis), confusion of C. passiflorae with, 263.

Corythaica passiflorae (planaris), foodplants and distribution of, 263, 264; parasite and control of, 264.

Corythucha spp., bionomics and control of.

on Platanus in U.S.A., 353. coscinia, Panoquina nyctelius.

Cosmolyce boetica, in harvested pea pods in Bengal, 128; on Gliricidia sepium in Hawaii, 250.

Cosmopolites sordidus (on banana), measures against, in Belgian Congo, 365; in Honduras, 261; in Pacific Is., 161, 202; utilisation of Plaesius javanus against, 202, 261.

Costa Rica, Corythaica passiflorae on Solanum torvum in, 263.

costalis, Psylla.

costimacula, Eublemma.

Cotinis nitida, in U.S.A., 344, 348; control of, in tobacco plant beds, 344, 345; attracted by caproic acid, 348.

Cotoneaster, Quadraspidiotus perniciosus

on, in France, 5.

Cotton, pests of, in S. Africa, 87, 127; pests of, in Brazil, 4, 93, 202, 204, 278, 311, 375; pests of, in Belgian Congo, 127, 366; pests of, in India, 20, 164, 296, 362, 386, 388, 389, 390, 391; Platyedra gossypiella on, in Mexico, 286: Helopeltis on, in Nigeria, 91, 383; pests of, in Nyasaland, 85, 86, 89-91, 358; pests of, in Peru, 132, 278; Heliothis armigera on, in Queensland, 32, 87; pests of, in S. Rhodesia, 87; pests of, pests of, in S. Rindessia, 87; pests of, in Russian Union, 306; pests of, in Sudan, 87; pests of, in Tanganyika, 88, 89; pests of, in Uganda, 88, 127; pests of, in U.S.A., 57, 70, 80, 102, 210, 211, 237, 260, 261, 270, 278, 286, 335, 336; pests of, in W. Indies, 97, 278; spread of Nematospora gossypii by Dysdercus on, 357; Saissetia nigra on, 373; varieties of, in relation to pests, 87, 88, 132, 278, 296, 362, 386, 387, 392; favouring Campsomeris mansueta, 85: effect of removal of squares on yield of, 80; dusting with calcium cyanamide to cause fall of leaves and squares of, 260; (harvested), regulations regarding importation of, into U.S.A., 193, Platyedra gossypiella intercepted in, in U.S.A., 100. (See Cottonseed.)

Cotton Aphis (see Aphis gossypii).

Weevil (see Anthonomus Cotton Boll grandis).

Cotton Bollworm, American (see Heliothis armigera).

Cotton Bollworm, Pink (see Platyedra gossypiella).

Cotton Bollworm, Red or Sudan (see Diparopsis castanea).

Cotton Bollworms, Spiny (see Earias). Cotton Fleahopper (see Psallus seriatus). Cotton Leafworm (see Alabama argillacea). Cotton Stainers (see Dysdercus and Oxycarenus).

Cotton Stem Weevil (see Pempherulus

affinis).

Cottonseed, regulations regarding importation and movement of, in U.S.A., 286; treatment of, against Platyedra gossypiella, 286, 375; manuring with, favouring infestation of cotton by pests, 88; (hulls), in baits for grass-

hoppers, **259.** Cottonseed Oil, emulsions of, against Coccids, 364; uses of, in experiments against Cydia pomonella, 108, 109, 274.

Couch Grass (see Agropyrum).

Cowpeas, mites on, in Ceylon, 299; pests of, in U.S.A., 191, 270.

cowperi, Coccophagus.

Crab Grass (see Digitaria sanguinalis). Crambus haytiellus, bionomics of, Florida, 115, 116. Crambus malacellus, measures against,

on maize and rice in Mauritius, 371. Cranberry, 343; pests of, in U.S.A., 23,

402.

Cranberry Fireworm (see Rhopobota naevana).

Cranberry Fruitworm (see Mineola vaccinii).

crassicornis, Meloboris. crassiseta, Ephialtes.

crataegi, Tetranychus; Thomasiniana.

Crataegus, Paratetranychus pilosus on, in Britain, 233; Quadraspidiotus perniciosus on, 5; Rhagoletis pomonella on, in Maine, 222; experiment with Thomasiniana theobaldi and, 308.

crawfordi, Opius.

Cremastus cooki, utilisation of, against Ancylis comptana in Ohio, 262.

Cremastus flaviventris, parasite of Cydia molesta in Argentina, 95.

Cremastus rubeo, parasite of Cydia molesta in Argentina, 95.

Creolin, tests and uses of, against pests, 304.

Creonaph, 268.

Creontiades femoralis, on cotton in Arizona, 101.

Creosote, types of, as barriers against Blissus leucopterus, 57; and oil, treatments of timber with, against Coleoptera, 72, 84, 371; injected into termite galleries in timber, 73; experiments with derivatives of, against termites, 243, 244; experiments with, against Tineola biselliella, 320; ineffective as a bee repellent in sprays, 212.

Cresol (Cresylic Acid), soil treatment with, against Aulacophora foveicollis, 17; experiments with, against Tineola bisel-

liella, 320.

Criocephalus productus, in fire-killed Douglas fir in U.S.A., 40.

Crioceris asparagi, on asparagus in Washington, 21.

critolaus, Spathius.

Crocidolomia binotalis (on crucifers), dip against, in S. Africa, 293; parasites of, in Fiji, 290.

Crotalaria, Agromyza phaseoli on, in

Belgian Congo, 363.

Crotalaria juncea, Exora spp. on, in S. Rhodesia, 126.

Crotalaria mucronata, Aphids on, in Hawaii, 251; Nezara on, in Fiji and Montserrat, 291.

Croton, Pseudococcus maritimus on, in Egypt, 227.

cruciata, Austroicetes.

crudum, Eulecanium (Lecanium) corni.

Cryolite, 239; against cabbage pests, 48, 293, 337; against cotton pests, 210, 211, 335, 336; against pests of leguminous plants, 3, 22, 48, 79, 176, 192, 277, 338; not protecting Narcissus bulbs from Merodon equestris, 24; against orchard pests, 22, 148, 149, 150, 179, 258, 284, 380; unsatisfactory against Pyroderces rileyi on orange, 113; against Acrobasis caryae on pecan, 210; against pests of small fruits, 4, 23, 262; against pests of solanaceous plants, 22, 45, 48, 51, 54, 59, 260, 280, 323, 324; against pests of other plants, 104, 112, 212, 260, 334; against Bruchid in stored cowpeas, 327; testing of, on Drosophila melanogaster, 144; in baits, 140, 252, 280; dusting with, 3, 4, 22, 24, 45, 48, 54, 59, 79, 104, 105, 112, 113, 144, 192, 210, 211, 212, 252, 260, 262, 293, 334, 335, 337; carriers for, 3, 22, 24, 45, 48, 54, 79, 112, 144, 212, 262, 293, 334; spraying with, 22, 23, 51, 53, 105, 113, 148, 150, 176, 179, 210, 252, 258, 260, 262, 277, 284, 323, 380; adhesives for, 51, 148, 149; formulae containing, 3, 22, 23, 24, 45, 48, 51, 54, 59, 79, 104, 112, 140, 148, 150, 176, 210, 212, 262, 280, 284, 293, 323, 334, 337; and nicotine, 337, 338, 380; and oil emulsion, 23, 149, 210, 284; and sulphur, 59, 335; effect of, on plants, 22, 24, 105, 149, 258; problems of residues of, 53, 149, 284; not toxic to residues of, 53, 149, 284; not toxic to residues of the residues 56; Alorco form of, 280; other insecticides compared with, 48, 176, 323, 324, 335, 336.

Cryptocephalus marginicollis, in Bahamas, 116; on orange in Cuba, 116.

Cryptochetum iceryae (parasite of Icerya spp.), bionomics of, in Ecuador, 95; introduced into Mona Island from California, 110.

Cryptococcus fagi (on beech), spread of, in

Canada, 155, 353, 354; in U.S.A., 354; bionomics of, 354.

Cryptognatha, predacious on Aonidiella aurantii in Brazil, 205.

Cryptognatha signata, predacious on Unaspis citri in Argentina, 133.

Cryptohypnus riparius (see Hypnoidus). Cryptolaemus montrouzieri, ineffective against Pseudococcus adonidum and P. comstocki in U.S.A., **59**, **209**.

Cryptorrhynchus mangiferae, on mango in New Caledonia, 161.

Cryptotermes (see Kalotermes).

Cryptus sexannulatus (parasite of Cydia pomonella), introduced into Canada, 356; in France, 355, 356; attempted rearing of, in New Zealand, 269; imported into U.S.A., 261; parasites of, 355; method of rearing on Ephestia, 74; not developing on artificial media, 75.

Crystox, 198.

Ctenolepisma quadriseriata, distribution and injuriousness of, in N. America, 107; value of laminated paper wrapping against, 272.

Ctenolepisma urbana, in buildings in U.S.A., 106, 107, 344; baits for, 344.

Cuba, Cryptocephalus marginicollis on orange in, 116; pests of solanaceous plants in, 263, 278; parasite of Diatraea saccharalis in, 176; beneficial insects imported into other countries from, 110, 177, 261.

cubae, Faustinus (Collabismodes).

Cubé, dusting with, 48, 50, 178, 239, 277, 282, 283; carriers for, 48, 239, 282, 283; spraying with, 113, 176, 239, 258, 268, 350; wetting agents for, 239, 240; formulae containing, 113, 239, 349; and DDT, 113; and oil emulsions, 113, 258, 350; and nicotine, 50, 239, 240; and soap, 268; and sulphur, 239, 240; and thiocyanates, 49; rotenone contents of, 49, 50, 176, 178, 239, 283; other insecticides compared with, 49, 176.

cucujiforme, Smodicum.

Cucumber, Tipula czizeki on, in Germany, 182; Trogophloeus elongatulus on, in Holland, 12; pests of, in Virginia, 212; squash preferred to, by Anasa tristis, 179.

cucumeris, Epitrix.

cucumis, Dacus (Austrodacus).

Cuerna occidentalis, transmitting virus disease of vines in U.S.A., 172.

Culex fatigans (quinquefasciatus, auct.), tests of plant extracts on larvae of, 65. culinaris, Rhyncolus.

Cultural Practices, influence of, on insect pests, 185, 186, 215, 216, 224, 258, 259, 297, 341, 383, 384, 392.

Cumquat, Unaspis citri on, in Argentina, 132.

cunea, Hyphantria.

cupreus, Corymbites.

Cupric Compounds (see under Copper). Cuprous Cyanide, causing spots on apples,

Cuprous Oxide, Red, effect of, in calcium arsenate dusts, 221.

Curculio nucum, bionomics of, in Denmark, 382.

Curculio uniformis, bionomics and control of, in U.S.A., 341.

Curculio, Plum (see Conotrachelus nenu-

curculionis, Bathyplectes.

curculionum, Eurytoma.

Curimosphena fasciculosa, on apple and peach in S. Rhodesia, 126.

Currant, pests of, in Britain, 360, 361; pests of, in Canada, 64; Lygus pabulinus on, in Holland, 253; Thomasiniana theobaldi not ovipositing on, 308.

Currant, Black, Otiorrhynchus singularis on, in Britain, 360.

curvicauda, Toxotrypana.

Cuspicona simplex, on Solanum spp. and tomato in New Zealand, 93.

cuspidata, Asamangulia.

Custard Apple (see Annona squamosa). Cutworms, Bufo spp. destroying in Oklahoma, 96; baits for, 140, 252, 263, 402. cyanea, Scutellista.

Cyanogas, 35, 255, 256.

cyanonotus, Cheiloneurus.

cyathicollis, Corythaica (see C. monacha). Cybocephalus, predacious on Chionaspis tegalensis in Mauritius, 371.

Cyclamen Mite (see Tarsonemus pallidus). Cycloneda munda, predacious on Saissetia nigra in California, 374.

Cycloneda sanguinea, predacious on Sipha

flava in Argentina, 133.

Cydia funebrana (on stone fruits), bionomics of, in Britain, 360; in Denmark, 381.

Cydia illutana, bionomics of, on conifers in Finland, 12.

Cydia latiferreana, bionomics and control of, on Corylus avellana in U.S.A., 341. Cydia microgrammana, cultural measures

against, on lucerne in Russia, 305. Cydia molesta (Oriental Peach Moth), bionomics and spread of, in S. America, 94; in U.S.A., 46, 105, 213, 219, 258; in Victoria, 322; on quince, 94; parasites and biological control of, 46, 95, 105, 213, 219, 322, 348; measures against, 94, 213; method of obtaining pupae of, in winter, 348.

Cydia nigricana (Pea Moth), bionomics of,

in Canada, 141, 145; in U.S.A., 22, 24, 100, 101, 141; on vetch, 22, 101; liberation of Macrocentrus ancylivorus against, 24.

44I

Cydia pomonella (Codling Moth), in S. Africa, 127; in Argentina, 94, 131; in Brazil, 94; in Britain, 359; in Canada, 74, 240, 241, 354, 356; in France, 354-356; in Germany, 119; in Italy, 6; in New Zealand, 269; in Poland, 117; in Spain, 125; in U.S.A., 22, 42, 56, 68, 108, 148, 149, 150, 151, 179, 191, 218, 240, 241, 258, 261, 273, 274, 281, 284, 314, 315; in Uruguay, 94; on pear, 240, 284, 355; on quince, 355; on stone fruits and nuts, 240, 355; bionomics of, 68, 284; graphic method of indicating population of, 56; parasites and biological control of, 74, 125, 240, 261, 269, 354-356; other natural enemies of, 240, 356; technique for obtaining pupae of, in winter, 348; fatty materials in diapausing larvae of, 264; chemical composition of cocoons of, 281; dusts against, 42, 149, 150, 179, 274, 275; experiments with inert dust against, 131; sprays against, 6, 22, 42, 119, 148, 149, 150, 151, 179, 218, 258, 273, 284, 315, 359; basic lead arsenate ineffective against, 314; timing of sprays against, 68; banding and orchard sanitation against, 69, 275, 316; repellents against full-fed larvae of, 274; soil treatment against cocoon stages of, 108, 281; fumigation of stored apples against, 127; tests of insecticides on, 106, 107, 214, 344.

Cydia strobilella, on conifers in Finland, 12.

cydiae, Ephialtes.

Cydonia japonica, Caliroa limacina on, in Britain, 360.

Cylas, favoured by virus disease on sweet potato in Uganda, 330.

Cylas formicarius (on sweet potato), in Fiji, 290; parasites of, on Ipomoea spp. in Louisiana, 216; possibly in S. Rhodesia, 126.

Cylas formicarius elegantulus, 216.

cylindricollis, Sitona.

Cylindrodendrum suffultum, sp. n., infesting Tipulid pupae in Britain, 232.

Cymatodera spp., predacious on Cydia pomonella in California, 240.

Cynara scolymus (see Artichoke).

Cynodon, Laphygma exempta E. Africa, 245.

Cynodon dactylon, Crambus haytiellus on, in Florida, 115.

Cynthia atalanta (see Vanessa). Cypripedium (see Orchids).

cytherea, Nudaurelia.

Cytisus, Quadraspidiotus perniciosus on, in France. 5.

Czechoslovakia, miscellaneous pests in, 8,

czizeki, Tipula.

D.

dacicida, Dinarmus.

Dacnusa leptogaster, parasite of Agromyzid in Belgium, 120.

Dactylis glomerata, Glyphipteryx achyloessa on, in New Zealand, 269.

dactylopii, Leptomastix.

Dactylosphaera (see Phylloxera).

Dacus, parasites and attempted biological control of, in Fiji, 161, 290, 291; peach, etc., in New Caledonia, 161.

Dacus cucumis, on cucurbits and papaya in Queensland, 162.

Dacus ferrugineus tryoni, bionomics and control of, in Queensland, 162.

Dacus musae, on banana in Queensland,

Dacus oleae, parasites of, on olive in Italy, 118.

Dacus passiflorae, host fruits of, in Fiji, 161.

Dacus xanthodes, host fruits and parasite of, in Fiji, 161, 290.

Dadap (see Erythrina).

Dahlia (host of tobacco virus), prohibition of introduction of, into Nyasaland, 85. Daihinia brevipes, bionomics and control

of, in U.S.A., 270.

Daikon, Aphids on, in Hawaii, 251.

Dakota, North, Daihinia brevipes in, 270; Epitrix cucumeris in, 220; natural enemies of noxious insects in, 96.

Dakota, South, Anabrus simplex in, 114; Epitrix cucumeris in, 220; mite attacking

grasshoppers in, 193.

Dalbergia sissoo, insects on, in India, 389, 390; treatment of timber of, against Sinoxylon, 371.

Dambala (see Psophocarpus tetragonolobus).

Daphne gnidium, Polychrosis botrana on, in Italy, 118.

Dasychira pudibunda, bionomics of, in Sweden, 19, 20.

Dasyhelea, on cacao in Trinidad, 78.

Dasyneura alopecuri, 170.

Dasyneura ignorata, cultural measures against, on lucerne in Russia, 305, 306. Dasyneura leguminicola, reducing seed production of clover in Montana, 101.

Dasyneura papaveris, bionomics of, on Papaver spp. in Hungary, 122.

Datana angusi, measures against, blueberry in U.S.A., 402.

Date Moth (see *Ephestia cautella*).

Datura stramonium, Gnorimoschema operculella on, in N. S. Wales, 185; dust of, ineffective against Calandra oryzae, 327.

Datura tatula, Gnorimoschema operculella on, in N. S. Wales, 185.

(parachlorphenyl)-1, 1, DDT [2, 2-bis 1-trichlorethane, against Aonidiella aurantii, 113; against Aphids, 218, against other Hemiptera, 173, 207, 322; against Atta laevigata, 288; against Coleoptera, 36, 125, 171, 172, 207, 217, 218, 289; mixed with grain against Calandra oryzae, 288; soil treatment with, against Diptera on vegetables, 288; in paints against Musca domestica, 137; against Lepidopterous larvae, 197, 213, 218, 288, 289, 344; effect of treating timber with, against termites, 244; against Thrips tabaci, 338; testing of, on earwigs and woodlice, 197; action of, on honey bees, 36, 213; use of, to eradicate honey-bee colony, 345; toxicity of, to Macrocentrus ancylivorus, 213; modes of action of, on insects, 137, 197, 257, 288, 289; question of fungicidal action of, 173, 207; studies on toxicity of, to mammals, 160, 245; poisoning of man by, 406; and injury to plants, 218; dusting with, 36, 113, 207, 217, 218, 282, 288; spraying with, 113, 171, 172, 173, 213, 217, 218, 288, 289, 338; doublenozzled spray apparatus for applying water and, 379; slow rate of evaporation of dusts and spray deposits of, formulae containing, 113, 173, 207, 213, 217, 218, 244, 338; adhesive for, 213; bentonite reducing toxicity of, 282; and calcite, 213; and cubé, 113; used with lead arsenate and limesulphur, 171; and oil emulsions, 113; pyrophyllite as carrier for, 207; compatible with fungicides, 173; solvents for, 244, 379; colorimetric tests for, 264, 368; research leading to discovery of, 137; proprietary preparations of, 36, 125, 171, 288, 345; DDT analogues (q.v.) compared with, 137, 197, 245, 344, 347; gammexane compared with, 257.

DDT Analogues, tests of, as insecticides, 137, 197, 245, 344; determination of distribution of, in insect tissue, 347; toxicity of, to rats, 245; colorimetric tests for, 368.

debellator, Alomya.

Decahydronaphthalene, as solvent for benzene hexachloride, 257.

decemlineata, Leptinotarsa (Chrysomela, Doryphora). decipiens, Carpophilus (see C. ligneus).

De-di-tox, **171**.

Degeeria collaris, parasite of Lochmaea suturalis in Britain, 233.

Delacroixia coronata (see Entomobhthora). Delaware, miscellaneous pests and biological control in, 78, 79, 80.

delegorguei, Encosternum (Haplosterna).

delicatus, Macrocentrus.

Delomerista diprionis, parasite of Gilpinia

hercyniae in Quebec, 224.

Delphacodes (Delphax) striatellus, experiments with virus disease of cereals and, in Russia, 183.

delta, Helochara.

deltae, Coccus (Lecanium).

Dendrobium pertinax (see Coelostethus).

Dendrocalamus (see Bamboos).

Dendroctonus frontalis, fungi associated with, on Pinus echinata in U.S.A., 286, 287.

Dendroctonus micans, study of damage to

spruce by, in Finland, 7, 8.

Dendroctonus pseudotsugae, fungus associated with, on Pseudotsuga taxifolia in U.S.A., 40.

Dendroctonus terebrans, factors affecting infestation of Pinus rigida by, in Massachusetts, 43.

Dendroctonus valens, factors affecting infestation of Pinus rigida by, in Massachusetts, 43.

Dendrothrips bispinosus, bionomics of, on

tea in Madras, 246.

Denmark, orchard pests in, 12, 380, 381, 382; miscellaneous pests in, 381. 382; parasite of Melolontha in, 380.

dentata, Melitara.

denticulatus, Ceuthorrhynchus.

dentipes, Monodontomerus.

Deob-dryer, 255.

Depressaria heracleana, bionomics and control of, on parsnip in Br. Columbia, 333.

Devaeocoris ruber, predacious on Paratetranychus pilosus in Britain, 234.

Dermestes, rare in houses in Germany, 229: feeding habits of, on cork oak in survey of data on Portugal, 310; injurious species of, 230.

Dermestes maculatus (vulpinus), products infested by, in U.S.A., 82, 102; bionomics and control of, in sheepskins, 82, 83; toxicity of gammexane to, 257.

derogata, Sylepta.

Derris, against Aphids, 2, 50, 276, 282, 283, 381; against Capsids, 37, 66, 153; against Coccids, 211, 212, 342, 346, 349, 350; against other Hemiptera, 154, 262; against Coleoptera, 3, 24, 48, 154, 176, 231, 233, 255, 256, 381; against Hylemyia brassicae, 154, 178; against Lepidoptera, 31, 48, 112, 139, 185, 197, 212, 231, 239, 332, 381, 404; treatments of stored potatoes with, against

Gnorimoschema operculella, 187; against Paratetranychus pilosus, 234; against sawflies, 360, 380, 381; against Thrips tabaci, 161; penetration of, through spiracles and cuticle of Melophagus ovinus, 383; dusting with, 2, 3, 24, 30, 37, 48, 50, 66, 78, 112, 153, 154, 161, 178, 185, 187, 212, 231, 233, 239, 256, 260, 262, 276, 282, 283, 322, 333, 360, 381, 404; carriers for, 24, 48, 112, 185, 187, 239, 262, 283; effect of carriers on insecticidal action of dusts of, 78, 152, 282, 283; conditioning agents for dusts of, 24, 50, 152, 240, 276, 277, 283; spraying with, 2, 3, 50, 139, 176, 185, 211, 212, 233, 239, 342, 345, 349, 350, 360, 380; spreaders for, 2, 139, 240; atomised sprays containing, 255; formulae containing, 50, 78, 112, 139, 152, 185, 187, 197, 211, 212, 239, 255, 349, 350; and nicotine, 2, 50, 66, 239; and oil emulsions, 51, 211, 212, 234, 255, 342, 345, 346, 349, 350; mutual solvents for extractives of when used with a discovery containing the second for extractives of, when used with oils, 211, 349, 350; and pyrethrum, 231, 233; and sulphur, 112, 239; rotenone contents of, 3, 24, 37, 48, 78, 112, 139, 176, 178, 185, 197, 211, 239, 255, 260, 262, 276, 277, 283, 381; insecticidal action of constituents of, 350; loss of rotenone and deguelin from mixtures containing, 152; other insecticides compared with, 48, 139, 176, 256.

Derris elliptica, D. malaccensis compared with, 49.

Derris malaccensis, rotenone contents and insecticidal value of dust of, 48, 49.

Desert Locust (see Schistocerca gregaria). desertana, Enarmonia (Steganoptycha) diniana.

desiliens, Gelechia.

destructor, Aphanotus (Tribolium); Aspidiotus; Ceroplastes; Glycyphagus; Mayetiola (Phytophaga); Phyllocoptes. detectus, Iridomyrmex.

detorquens, Technomyrmex.

devastans, Empoasca.

Dexia rustica, bionomics of, in Denmark,

diabola, Solenopsis geminata.

Diabrotica, of Argentina, 242.

Diabrotica duodecimpunctata, tests insecticides on, 278, 279.

Diabrotica longicornis, on maize in Iowa, 101.

melanocephala, Diabrotica insecticides against, on cucurbits in Virginia, 212. Diabrotica speciosa, bionomics of, in

Argentina, 242.

Diabrotica undecimpunctata, bionomics and control of, in California, 30, 104. diabroticae, Celatoria.

diadema, Sinea.

Diadromus collaris (see Thyracella).

Dialdehyde, insects repelled by solution containing, 272.

Dialeurodes citrifolii, measures against, on

Citrus in Bermuda, 172.

Diamond-back Moth (see Plutella maculipennis).

Diaphania hyalinata, tests of insecticides on, in Porto Rico, 339.

Diaphania nitidalis, insecticides against,

on cucurbits in Virginia, 212. Diaprepes abbreviatus, seasonal abundance of, on sugar-cane in Porto Rico, 243.

Diaspidiotus ancylus, spray against, on blueberry in U.S.A., 401.

diaspidis, Aphytis.

Diaspis bromeliae, bionomics and control of, on pineapple in Queensland, 248.

Diaspis echinocacti, on Opuntia in Nyasaland, 86.

Diatomaceous Earths, as carrier for cubé and derris dusts, 283; calcined forms of, against pests in stored wheat, 372.

Diatraea grandiosella, on sorghum and maize in Arizona, 102, 176, natural enemies of, 176, 177.

Diatraea lineolata, parasite of, in Trinidad,

Diatraea saccharalis (Sugar-cane Moth Borer), in Mexico, 176; in U.S.A., 26, 116, 177, 260, 261; in W. Indies, 26, 97, 176, 243, 132; on maize, 26; bionomics of, 116; ant destroying eggs of, 243; parasites and biological control of, 26, 97, 176, 177, 243, 260, 261, 312; other measures against, 25, 26, **260:** methods for rearing and observing, 205.

Apanteles; diatraeae, Lixophaga; Paratheresia (see Theresia claripalpis).

Dibrachys affinis, parasite of Cydia pomo-

nella in France, 355.

Dibrachys cavus (parasite and hyperparasite of Cydia spp.), in Brazil and Uruguay, 95; in California, 240; in France, 355; Trichomalus trujilloi treated as distinct from, 95.

Dibromodiphenyltrichlorethane (see DDT

Analogues).

Dibutyl Phthalate, mixture containing, as intermediary solvent for derris resins with oils, 211.

4,4'-Dichlorbenzophenone (see DDT Ana-

logues).

Dichlordifluoromethane, use of, for producing aerosols, 43, 99, 142, 261; calculation of change of insecticide concentration in, during aerosol discharge, 332.

 α - α -Dichlor-diphenyl- $\beta\beta$ -trichlorethane

(see DDT).

Dichlorethane (see Ethylene Dichloride). Dichlorethyl Ether, experiments with, against cheese mites, 130, 131, 269; in oil for treating maize ears against Heliothis armigera, 337; in sprays against Hylemyia spp., 276, 338; soil treatment with emulsions of, against pests, 104, 337, 338; in oil emulsions against cocoon stages of Cydia pomonella in soil, 108; effect of addition of, to nicotine dusts, 338; formulae containing, 104, 108, 276, 337.

1,1-Dichlor-1-nitroethane, 237.

Dichlorpropane, and dichlorpropylene, experiments with, against wireworms, 55, 56.

Dichlorpropylene, and dichlorpropane, experiments with, against wireworms,

Dichomeris ventrellus, on pecan in Texas, 347.

dicincta, Mylabris.

Dicladocerus westwoodi, establishment of, against Coleophora laricella in Ontario, 143.

dictyospermi, Chrysomphalus.

Dicyclohexylamine Dinitro-o-cyclohexylphenate (see Dinitro-ortho-cyclohexylphenol).

Dielis (see Campsomeris).

Dieseline, ineffective against biselliella, 320.

japanica Diestrammena (japonica, marmorata, De Haan), confusion of Tachycines asynamorus with, 146. differentialis, Melanoplus.

Digitalis purpurea, Macrosiphum solani on, in Britain, 358.

Digitaria sanguinalis, Elasmopalpus lignosellus on, in Arkansas, 192.

dignus, Apanteles,

9,10-Dihydroanthracene, toxicity of, to Cydia pomonella, 215.

Dihydro - methoxy - methyl - oxonicotinonitrile, toxicity of, to Cydia pomonella, 106.

Dihydro - trimethyl - oxonicotinonitrile, toxicity of, to Cydia pomonella, 106.

Dikraneura, on cotton in Peru, 132. Dilachnus (see Cinara).

Dilophonota ello (see Erinnyis).

Dimethyl Ether, use of, for producing aerosols, 99.

Dimethyl-pentanone Semicarbazone, 99.

dimidiatus, Phymatodes. Dinarmus dacicida, parasite of Dacus

oleae in Italy, 118. Dinarmus sauteri, parasite of Pempherulus

affinis in India, 21. diniana, Enarmonia (Steganoptycha).

Dinitro-\alpha-naphthol, Anthrenus not festing materials dyed with, 73.

Dinitrocresol, preparation containing, against Psylla mali, 117. (See Dinitroortho-cresol.)

INDEX. 445

Dinitro-o-cresol, in dormant sprays against orchard pests, 34, 35, 58, 153, 179, 233, 253; other uses of, against insects, 108, 109, 153, 319; reactions of mature larvae of Cydia pomonella to, 274; not repellent to honey-bees, 6: tests of toxicity of, to insect eggs, 15. Parthenium argentatum injured by dust containing, 105; effects of, on cattle and fowls, 35; factors affecting solubility of, in water, 58; formulae containing, 58, 109, 153, 274, 319; Bordeaux mixture reducing effectiveness of, 35; lye not increasing effectiveness of, 179; and oil emulsions, 108, 109, 153, 179, 233, 253, 319; effect of oil emulsions on, 15, 179; derivatives compared with, as insecticides, 198; question of toxicity of salts of, 15. (See Sodium Dinitro-o-cresylate.)

Dinitro-o-cresol Propionate, toxicity of, to insects and plants, 279.

Dinitro-o-cyclohexylphenol, and sulphur, spraying with, against Paratetranychus citri, 172; increasing efficiency of oil emulsions against Parlatoria chinensis, Parthenium argentatum injured by, 105; dicyclohexylamine salt of, in dust and sprays, 45, 53, 54, 208.

Dinitro-o-tolyl Acetate, insecticidal pro-

perties of, 198.

Dinitrophenol Propionate, toxicity of, to insects and plants, 279.

Dinocampus (see Perilitus).

Dinoderus, bionomics and control of, in stored bamboos in India, 71, 72.

Dinoderus minutus, in stored maize and bamboos in Porto Rico, 312; imported bamboos resistant to, 312.

Dioryctria abietella, on conifers in Finland,

Diospyros (see Persimmon).

Diospyros ebenaster, Anastrepha ludens on, in Mexico, 340.

Diospyros virginiana, Schistocerca obscura feeding on, in Oklahoma, 237.

2,2 - Di - para - anisyl - 1,1,1 - trichlorethane (see DDT Analogues).

2.2 - Di - paraphenetyl - 1,1,1 - trichlorethane (see DDT Analogues).

2,2 - Di - paratolyl - 1,1,1 - trichlorethane (see DDT Analogues).

Diparopsis castanea (on cotton), in S. in Nyasaland, 89; Rhodesia, 87; bionomics of, in Nyasaland, 85, 89-91.

Diphenyl Sulphide and Sulphoxide, effect of aerosols of, on Tarsonemus pallidus and Antirrhinum, 44.

 $\beta\beta\beta$ - trichlorethane αα - Diphenyl (see DDT Analogues).

Diplazon agilis, parasite of Coleophora laricella in Ontario, 143.

Diplazon laetatorius (parasite of Syrphids)? parasite of, in Argentina, 117: Fiji, **290.**

Diplogaster, 61.

Diprion pini, on pine in Germany, 14, 16: ant predactions on, 16: disease of Gilpinia hercyniae attacking, 398.

Diprion similis, on Pinus mugo in Pennsylvania, 4.

diprionis, Delomerista.

Dirhinus, failure to establish, against Dacus in Fiji, 291.

discoidalis. Exora.

disconotella, Aristotelia.

Disodium Ethylene Bisdithiocarbamate, in sprays against mites, 378, 379.

dispar, Lymantria (Liparis, Porthetria); Xyleborus (Anisandrus).

disparis, Anastatus.

disstria, Malacosoma,

Distantiella theobroma, bionomics and distribution of, on cacao, etc., in W. Africa, 65, 66, 195, 196; on Citrus, 196; insecticides against, 66.

distigma, Chilocorus.
Distillers' Grain, ineffective as baits for cutworms and earwigs, 140, 142.

distincta, Anastrepha. Dithane, 378, 379.

Ditula angustiorana (see Batodes).

divergens, Phyrdenus.

diversa, Agathis (Bassus).

Diversinervus elegans, parasite of Saissetia nigra in Australia, 374; hosts of, in Palestine, 326.

Diversinervus smithi, introduced parasite of Saissetia oleae in California, 374; attempted use of, against S. nigra, 374. divisus, Macrosteles.

Dixippus, tests with charcoal dust on, 189. DN-111, **53, 208.**

Dociostaurus maroccanus, in Iraq, 199–201; in Spain, 166, 167; bionomics and ecology of, 167, 199-201; forms of, 166, 167; measures against, 201.

Dodecyl Nitrile, action of, on insects, 198. Dodecyl Thiocyanates, 198. (See Thiocyanates.)

Dogwood (see Cornus).

Dolichosoma lineare, possibly predacious on Cydia pomonella in France, 356.

Dolomite, tests with, against pests in stored wheat, 318, 319, 372; ineffective against Gnorimoschema operculella, 31.

domestica, Musca; Thermobia.

domesticus, Gryllulus; Opilo.

dominica, Rhizopertha.

Doralis (see Aphis).

Doryphora decemlineata (see Leptinotarsa). Dosage-mortality Curves, 64.

Douglas Fir (see Pseudotsuga taxifolia). Dowicide G, 272.

Downy Mildew (see Pseudoperonospora humuli).

Dracaena, Pseudococcus spp. on, in California and Egypt, 59, 227.

Draeculacephala minerva, in U.S.A., 104, 172; measures against, on Parthenium argentatum, 104; transmitting virus diseases of vines, 172.

Dragonflies, economic importance of, 5.

dregii, Calidea.

Drepanosiphum platanoides, spray against, on Platanus and maples in U.S.A., 353. Dreyfusia (see Chermes).

Dried Fruit Beetle (see Carpophilus hemipterus).

Dried Fruit Moth (see Plodia interpunctella).

Drosophila melanogaster, experiments with, as host for parasites of Oscinella frit, 75; use of, for testing insecticides, 144, effect of temperature on 188, 189; development of, 187.

Drosophila subobscura, action of charcoal

dust on, 189.

drosophilae, Spalangia. drummondi, Melanophila. druparum, Callimome.

Dryobates villosus, destroying Depressaria heracleana in Br. Columbia, 333. dubiosus, Neodiprion.

Duboisia hopwoodii, insecticidal constituent of, 54.

duodecimpunctata, Diabrotica.

Dusts, use of oil-impregnated, 24, 150, 158, 159, 238, 276, 277, 283, 316, 317; electrostatic charges of, 277, 283; particle size of, 42, 283; apparatus for applying, 57, 214, 334; application of, from aircraft, 238, 260, 401; method of using explosives for applying, in forests, 46; (inert), tests and use of, against pests of stored products, 31, 66, 131, 156, 157, 231, 318, 319, 327, 372; against field pests, 131, 231; action of, on insects, 131, 156, 188, 231, 372.

Dutch Elm Disease (see Ceratostomella

ulmi).

Dutox, 280.

Dysdercus (Cotton Stainers), in Nyasaland, 91, 358; in Peru and Porto Rico, 312; in Tanganyika, 88; method of spread of Nematospora by, 357; natural enemies and biological control of, 88, bait-spray for, 88; tests of charcoal dust on, 189.

Dysdercus andreae, parasites introduced against, in Porto Rico, 312.

Dysdercus cardinalis, on cotton in Tangan-

yika, 88, 89. Dysdercus fasciatus (on cotton), in Belgian Congo, 367; in Nyasaland, 91;

S. Rhodesia, 87; in Tanganyika, 88; bionomics of, 91, 367.

Dysdercus howardi, effects of temperature and humidity on, 135.

Dysdercus intermedius (on cotton), bionomics of, in Nyasaland, 91; in S. Rhodesia, 87; in Tanganyika, 88.

Dysdercus melanoderes, bionomics of, on cotton in Belgian Congo, 127, 128, 367; status and interbreeding of,

D. nigrofasciatus, 127.

Dysdercus neglectus (see D. sanguinarius). Dysdercus nigrescens, sp. n., a mutant of D. nigrofasciatus in Belgian Congo, 127.

Dysdercus nigrofasciatus (on cotton), in E. Africa, 88, 89, 127; in S. Africa, 127; in Belgian Congo, 127, 367; bionomics of, 127, 128, 367; status, forms and interbreeding of, with D. melanoderes, 127.

Dysdercus ruficollis, on cotton in Peru,

Dysdercus sanguinarius, parasites introduced against, in Porto Rico, 312.

Dysdercus superstitiosus (on cotton), in E. Africa, 88, 89; in Belgian Congo, 127, 137; in S. Rhodesia, 87; bionomics of, 367; D. nigrofasciatus and D. melanoderes distinct from, 127.

Dysstroma citrata immanata, on strawberry, etc., in Britain, 360.

Ε.

Earias, on cotton in Tanganyika, 88, 89. Earias biplaga (on cotton), in Belgian Congo, 367; in Tanganyika, 88.

Earias fabia, bionomics of, on cotton in

India, 389.

Earias insulana (on cotton), in Belgian Congo, 367; bionomics of, in India, 389; in Tanganyika, 88.

earterus, Apanteles.

Earwig, European (see Forficula auricularia).

Earwig, Ring-legged (see Anisolabis annulipes).

ebeninum, Monomorium carbonarium.

echinocacti, Diaspis.

echinopus, Rhizoglyphus. echion, Tmolus (Thecla).

Economic Entomology, bibliography of, in Finland, 406; organisation and problems of, in various countries, 5: book on, 328.

Ecpantheria hambletoni, bionomics of, in Brazil, 204.

Ecuador, miscellaneous pests in, 5, 95; natural enemies and biological control of Icerya montservatensis in, 95, 96. egena, Autographa; Pentilia.

Egg, Trogoderma versicolor infesting dried. 102; nutritional value of constituents

of, for Tineola biselliella, 351.

Egg-plant, Corythaica passiflorae on, in Brazil and Porto Rico, 263, 264; Epilachna vigintioctopunctata on, in Fiji, 290; pests of, in Hawaii, 250, 251; pests of, in U.S.A., 51.

Egius platycephalus, predacious on Coccids in Porto Rico, 313; imported into

Panama and Trinidad, 313.

Egypt, pests of *Citrus* in, 228, 255, 256; miscellaneous pests in, 5, 226, 227; beneficial insects in, 225, 227; Rodolia cardinalis introduced into India from, 385.

Eichhornia, Aphid on, in Fiji, 290.

Elachista, on sweet potato in Mauritius, 371.

Elaeis guineēnsis, Bruchid infesting field and stored nuts of, in Nigeria, 165, 370. Elasmopalpus lignosellus, food-plants of,

bionomics

in U.S.A., 102, 191, 192; and control of, 191, 192.

Elasmus, parasite of Asamangulia cuspidata in India, 391.

Elaterids (see Wireworms).

Eldana saccharina, bionomics of, on cassava in Belgian Congo, 367.

Electro-magnetic Radiation (see Infra-red

Rays and Light). Electrostatic Charges, of dusts, 277, 283.

elegans, Diversinervus. elegans, Lea, Anthicus (see A. troilus).

elegantalis, Leucinodes.

elegantulus, Cylas formicarius; Lixus.

Elettaria cardamomum, Aphid transmitting virus disease of, in India, 392.

Eleusine coracana, Lygaeid on, in S. Rhodesia, 126.

Elgetol (see Sodium Dinitro-o-cresylate). elisus, Lygus.

elita, Eumacronychia.

ello, Erinnyis (Dilophonota).

Elm, Paratetranychus pilosus Britain, 233; Agelastica alni on, in Germany, 231; pests of, in U.S.A., 43, 110, 173, 180, 237, 270, 313; factors affecting Scolytids and Ceratostomella ulmi in, 43, 110, 173, 174, 313.

Elm Leaf Beetle (see Agelastica alni). Elodia tragica, parasite of Cydia pomonella in France, 355; suggested introduction of, into Canada, 356.

elongatulus, Trogophloeus.

elongatus, Coccus; Eremotes; Thysanus.

Elsinoe fawcetti, 205. elutella, Ephestia.

emarginata, Calpe. emigratella, Amorbia.

Empoasca, on cotton in Peru, 132.

Empoasca arida, measures against, on Parthenium argentatum in California, 104. Empoasca benedettoi, variety of cotton resistant to, in Sudan, 88.

Empoasca devastans, studies on resistance of varieties of cotton to, in India, 296, 362, 386; methods of estimating populations of, 296.

Empoasca fabae, measures and experiments against, on potato and hops in

U.S.A., 173, 180, 206, 207, 317. Empoasca solana, on beet in Arizona, 279. Empria ignota, insecticides against, on

strawberry in Ohio, 263.

Empria maculata, insecticides against, on strawberry in Ohio, 263.

Empusa forficulae var. major, infesting Forficula auricularia in Britain, 232.

Empusa grylli, infesting Calliptamus italicus in Bulgaria, 122.

Enargopelte, parasite of Asterolecanium

pustulans in Egypt, 227.

Enarmonia diniana, 400; parasites of, 401. Enarmonia diniana desertana, bionomics of, on Larix sibirica in Siberia, 400, 401; dust against, 401.

Encelia farinosa, Plusia brassicae on, in

Arizona, 112.

Encosternum delegorguei, use of, as food in Africa, 327.

Encyrtus (parasite of Coccids), in Australia, 374; in Palestine, 325, 326.

Encyrtus barbatus, parasite of Saissetia nigra in Hawaii, 374.

Encyrtus consobrinus, 326.

Encyrtus frontatus, hosts of, in Palestine,

Encyrtus infelix (see Eucomys).

enecator, Trichomma.

Entandrophragma caudatum, resistance of timber of, to termites in N. Rhodesia,

Entedon pempheridis, parasite of Pempherulus affinis in India, 21.

Entomology, book on, 379. (See Economic

Entomology.) entomophagus, Thyreophagus (Monieziella). Entomophthora anglica, sp. n., infesting Coleoptera in Britain, 232.

Entomophthora coleopterorum, infesting Sitona flavescens in Britain, 232.

Entomophthora coronata, experimentally infesting Kalotermes approximatus in Bermuda, 290.

Entomophthora virescens, infesting cabbage caterpillar in Britain, 232.

Eosin, unsatisfactory for marking Pyrilla,

Ephestia cautella, inported into Germany, 231; in Gold Coast, 66; measures against, in stored ground-nuts in Queensland, 31; in stored cacao, 66, 231: inert dust ineffective against, 66.

Ephestia elutella, products infested by in Germany, 230, 231; eradication of, in tobacco factory in S. Rhodesia, 127; other pests confused with, 231.

Ephestia kuehniella, in dehydrated vegetables in California, 102; E. elutella confused with, in Germany, 232; fumigation controlling, in flour mills in Victoria, 130; tests of fumigants on, 295; experiments with inert dusts and, 157, 231; use of eggs of, for testing ovicides, 15; effect of temperature on development of, 187; as host for parasites of other Lepidoptera, 74, 75.

Ephialtes aequalis (see Pimpla).

Ephialtes argentinus, parasite of Cydia

molesta in Argentina, 95.

Ephialtes caudatus (parasite of Cydia pomonella), introduction and liberation of, in Canada, 356; in France, 355, 356; utilisation of, in New Zealand, 269; imported into U.S.A., 261; method of rearing, on Ephestia, 74; not developing on artificial media, 75; recorded as E. extensor, 356.

Ephialtes crassiseta, parasite of Cydia pomonella in France, 355, 356; recorded

as E. extensor, 356.

Ephialtes cydiae, parasite of Cydia pomon-

ella in France, 355, 356.

Ephialtes extensor, É. caudatus and E. crassiseta recorded as, 356; synonym of E. punctulatus, 356.

Ephialtes inquisitor, bionomics of, parasitising Dasychira pudibunda in Sweden,

20.

Ephialtes punctulatus, E. extensor (q.v.) a synonym of, 356.

Ephialtes ruficollis, parasite of Cydia

pomonella in France, 355, 356. Ephialtes sanguinipes (see Pimpla).

Ephialtes terebrans, parasite of Pissodes validirostris in Finland, 11.

Epicauta atomaria, tests with DDT against, on potato in Brazil, 289.

Epichorista ionephela, on carnation in S. Rhodesia, 126.

Epilachna, not parasitised by Perilitus coccinellae, 369.

Epilachna chrysomelina, on melons in Greece, 16.

Epilachna varivestis (on beans), in Canada, 139, 144; tests and use of insecticides against, in U.S.A., 3, 48, 49, 65, 79, 175, 179, 260, 277; Lydinolydella metallica imported into U.S.A. against, 261; on lima beans, 79; bionomics of, 3; oil-burner for destroying beans infested by, 139.

Epilachna vigintioctopunctata, on vegetables in Fiji, 290.

epinotiae, Phaeogenes.

Epitetranychus (see Tetranychus).

epitricis, Microctonus.

Épitrix, distribution and identity of black

species of, on potato, etc., in N.America, 220.

Epitrix cucumeris, in Canada, 220; measures and experiments against, on potato in U.S.A., 48, 180, 206, 207, 260; distribution and confusion of, with other species, 220.

Epitrix hirtipennis, bionomics and control of, on tobacco in U.S.A., 41, 45;

recorded as E. parvula, 41.

Epitrix parvula, E. hirtipennis confused with, in U.S.A., 41.

Epitrix similaris, sp. n., on potato in California, **220.**

Epitrix subcrinita, on potato, etc., in U.S.A., 22, 220; adult habits and control of, 22.

Epitrix tuberis, sp. n., bionomics of, on potato, etc., in U.S.A., 22, 220; measures against, 22.

Epiurus inquisitor (see Ephialtes).

epunctifera, Sesamia.

equestris, Merodon.

Erechtites valerianaefolia, Aphids on, in Hawaii, 251.

eremocitri, Aonidiella.

Eremotes elongatus, humidity requirements of, in timber in Germany, 18.

Eretmocerus serius, imported into Mexico against Aleurocanthus woglumi, 261.
Ergates faber, bionomics and physiology

of, in timber in Germany, 18, 19.

Ergates spiculatus, in fire-killed Douglas

fir in U.S.A., 40. ericae, Nysius.

Erigorgus melanobatus, parasite of Ocnogyna in Spain, 170.

Erinnyis ello, bionomics of, on cassava in Brazil, 13.

Eriobotrya japonica, Ecpantheria hambletoni on, in Brazil, 204.

Eriocampoides (see Caliroa).

eriophori, Anuraphis (Ceruraphis).

Eriophyes carinatus, on tea in Madras, 246. Eriophyes litchii, sp. n., on Litchi chinensis in Hawaii, 111, 250.

Eriophyes oleivorus (see Phyllocoptruta).

Eriophyes pyri, bionomics and control of, on pear in Washington, 285.

Eriophyes sheldoni, mite predacious on, on lemon in California, 397.

Eriophyes vaccinii, on blueberry in U.S.A., 401.

Eriosoma lanigerum (on apple), in Britain, 361; in California, 282; in Germany, 34; in Mexico, 261; natural enemies and biological control of, 261, 282, 361; factors affecting sprays against, 34, 253.

Eritrea, introduction of Gnorimoschema operculella and its parasites into, 379. Ernestia consobrina, bionomics of, in

Russia, 304, 305.

Erodium, Dociostaurus maroccanus on, in Iraq, 200.

Erythrina, Terastia meticulosalis on, in Madras, 246.

erythrocephala, Acantholyda.

Erythroneura comes, on vines in U.S.A., 28, 44; method of estimating populations of, 44; sprays against, 28.

Erythrophloeum africanum, resistance of timber of, to termites in N. Rhodesia. 247.

erythrostethus, Blastothrix.

Estigmena chinensis, on bamboos in India,

Estigmene acraea, measures against, on Parthenium argentatum in California,

Estonia, Taeniocampa gracilis on apple in, 119.

Ether, as a solvent, 281.

Ethyl Alcohol (see Alcohol).

2-Ethyl-pyridine Sulphate, effect of dust containing, against Hylemyia brassicae,

Ethylene Dichloride, in fumigant mixtures against pests of stored cereals, 235, 236, 237, 318, 319; sorption by wheat and wheat products of, 236, 237, 264; determination of, in wheat products, 264; and carbon tetrachloride, not highly effective against Dermestes in sheep skins, 83; and oil emulsion, soil treatment with, against cocoons of Cydia pomonella, 108; in oil for treating maize ears against Heliothis armigera, 337.

Ethylene Glycol Monoethyl Ether, as a solvent, 347.

Ethylene Oxide, tests and use of, as a fumigant against pests of stored products, 103, 231, 295, 296, 403; and carbon dioxide, 103, 231; sorption and determination of, in wheat, 128; recirculation increasing effectiveness of, in vacuum chamber, 103.

Etiella zinckenella, on peas in Bengal, 128. Euaraba tergata, relation of, to Anabrus simplex in Nevada, 146.

Eublemma angulifera, bionomics of, on mango in Travancore, 387.

Eublemma costimacula, predacious on Pseudococcus spp. in Nyasaland, 86; parasite of, 86.

Eublemma scitula, predacious on Coccus hesperidum in Palestine, 326.

Euborellia annulipes (see Anisolabis).

Eucalyptus, pests of, in S. Africa, 83; Helopeltis orophila on, in Belgian Congo, 37; pests of, in Mauritius, 371; Anoplognathus on, in N.S. Wales, 162.

Eucalyptus gigantea, infestation of firekilled timber of, by borers in Victoria, 377, 378.

Eucalyptus goniocalyx, Platypodids in timber of, in Victoria, 378.

Eucalyptus regnans, infestation of firekilled timber of, by borers in Victoria, 377, 378.

Eucelatoria armigera (see Blondelia).

Eucnocerus anthrenoides (see Trogoderma ornatum).

Eucoila, parasite of Agromyza strigata in Fiji, 290.

Eucomis undulata, possible transmission of Ornithogalum mosaic to, 1.

Eucomys infelix, parasite of Saissetia nigra in Hawaii, 374.

Eucomys lecaniorum, parasite of Coccus hesperidum in Palestine, 325, 326.

Euderus amphis, parasite of Coleophora laricella in Ontario, 143.

Euderus pempheriphila, parasite of Pempherulus affinis in India, 21.

Eugenia, Anastrepha fraterculus on, in

S. America, 397. Eugenia cumini (jambolana), Argyroploce

aprobola on, in Travancore, 387. Eugenia jambos, Anastrepha spp. on, in

Mexico, 52, 395, 397.

Eugenol, in baits for Polychrosis and Popillia, 27, 55.

Eulecanium berberidis (see E. persicae). Eulecanium bituberculatum, on apple in Denmark, 381.

Eulecanium corni, spray against, Platanus in U.S.A., 353.

Eulecanium corni crudum, bionomics and control of, on Taxus in Britain, 171. Eulecanium excrescens, spray against, on

Platanus in U.S.A., 353.

Eulecanium nigrofasciatum, on blueberry in U.S.A., 401.

Eulecanium persicae, measures against, on mulberry and vines in Egypt, 228.

Eulecanium pruinosum, spray against, on Platanus in U.S.A., 353.

Eulimneria, parasite of Gnorimoschema operculella in Eritrea, 379.

Eulimneria pattoni, parasite of Crambus haytiellus in Florida, 115.

Eulophonotus myrmeleon, bionomics of, on cacao in Belgian Congo, 366.

Eulophus longulus, parasite of Dacus oleae in Italy, 118.

Eumacronychia elita, relation of, to Anabrus simplex in Nevada, 146.

Eumaenas salaminia, measures against, damaging fruits in Queensland, 291.

Eumerus aurifrons, on cassava in Hawaii, 250.

Eumerus marginatus, on cassava Hawaii, 250.

Euonymus, Icerya purchasi on, in France, 357; experiment with Aphis viburni and, 383.

Eupelmus coccidivorus, hyperparasite of

Saissetia nigra in Porto Rico and Panama, 374.

Eupelmus urozonus, hosts and ecology of, in Italy, 118.

Euphoria inda, bionomics and control of, on strawberry in Ohio, 262.

Euphorus miridicidus, parasite of Helopeltis orophila in Belgian Congo, 37.

Euphorus sahlbergellae, parasite of Capsids in W. Africa, 65, 66, 195, 196; failure to rear, on Distantiella theobroma, 66; parasite of, 195.

Euplectrus, parasite of Prodenia litura in Fiji, 202.

Euplectrus comstocki, parasite of Plusia brassicae in Arizona, 112.

Euplectrus mellipes, parasite of Coleophora laricella in Ontario, 143.

Euproctis chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Euproctis terminalis, on pines in S. Africa, 83, 324; proposed biological control of, 324.

Eurychilopterella luridula, predacious on Pseudococcus comstocki in U.S.A., 209.

Eurygaster integriceps (on cereals), in Persia and Iraq, 302; bionomics of, in Russian Union, 159, 301-303; parasites and biological control of, 301, 302, 303; use of fowls to destroy, 159, 303; aeroplane dusting and spraying against, 303; other measures against, 159.

eurytheme, Colias (Eurymus).

Eurytoma, parasite of Ecpantheria hambletoni in Brazil, 204.

Eurytoma curculionum, parasite of Pissodes validirostris in Finland, 12.

Eurytoma ischioxanthos, parasite of Ips spp. in Finland, 12.

Eurytoma nobbei, parasite of Magdalis violacea in Finland, 12.

Eurytoma rosae, hosts and ecology of, in Italy, 118.

Eurytoma rufipes, hosts of, in Finland, 12. euschisti, Trissolcus.

Euschistus servus, on peach in Illinois, 316. Euschistus tristigmus, on peach in Illinois, 316.

Euschistus variolarius, on peach in Illinois, 316.

Eutermes walkeri, in imported timber in New Zealand, 234, 235.

Eutettix tenellus (in U.S.A.), transmitting curly-top in beet and flax, 101, 378; on weeds, 281, 314; new weed host of curly-top transmitted by, 281; effects of varying temperatures on development of, in U.S.A., 220; factors affecting catches of, by different methods, 314. Eutinobothrus brasiliensis, bionomics and

control of, on cotton in Brazil, 202,

203, 311.

Eutrombidium locustarum (see E. trigonum).

Eutrombidium trigonum, bionomics and synonymy of, attacking grasshoppers in U.S.A., 193.

Euxenus variegatus (see Faustinus cubae). Euxoa, on peas in Quebec, 223.

Euxoa nigricans, on vines in Holland, 12.

Euxoa segetum (see Agrotis).

Evergreen, 51.

exarator, Spathius. exareolata, Angitia.

excrescens, Eulecanium (Lecanium).

exempta, Laphygma.

Exenterus, utilisation of, against Gilpinia hercyniae in Br. Columbia, 155.

exigua, Laphygma. exitiabilis, Pseudococcus.

exitiosa, Aegeria.

Exochomus melanocephalus (nigromaculatus), predacious on Pseudococcus spp. in Nyasaland, 86.

Exochomus quadripustulatus, establishment of, attacking Coccids and Eriosoma lanigerum in California, 282.

exoleta, Xylena (Calocampa).

Exora apicipennis, measures against, on Crotalaria juncea in S. Rhodesia, 126. Exora discoidalis, measures against, on Crotalaria juncea in S. Rhodesia, 126.

Exorista larvarum (see Tachina).

exotica, Allograpta.

Explosives, method of using, to apply insecticide dusts in forests, 46; for destroying termite nests, 73.

expulsus, Apanteles. extensor, Ephialtes (see E. punctulatus).

Extrax, 51.

Eye-spotted Bud-moth (see Spilonota oçellana).

F.

fabae, Aphis (Doralis); Empoasca.

faber, Ergates. fabia, Earias.

fagi, Cryptococcus.

farinae, Tyroglyphus.

fascialis, Hymenia (see H. recurvalis).

fasciatus, Anthrenus; Dysdercus; Hercothrips; Oncopeltus.

fasciculatus, Araecerus.

fasciculosa, Curimosphena.

fatigans, Culex. fausta, Rhagoletis.

Faustinus apicalis, on solanaceous plants in Brazil, 288.

Faustinus cubae (tabaci, variegatus), bionomics of, in Brazil, 288; synonymy of,

Feathers, insects infesting, 163, 229, 351. felinelle, Lithocolletis.

femoralis, Creontiades. femur-rubrum, Melanoplus. fenestralis, Scenopinus.

fennica, Agrotis.

Fenugreek (see Trigonella foenum-grae-cum).

Fermate (see Ferric Dimethyl Dithiocarbamate).

Ferric Chloride, for treating wood against termites, 244.

Ferric Dimethyl Dithiocarbamate, 150; increasing infestation of apple by Paratetranychus pilosus, 208; not reducing effect of oil emulsions on P. pilosus, 148, 208; and bentonite, as repellentspray against Popillia japonica, 180; and lead arsenate in sprays and dusts, 54, 151; solutions of, protecting wood against termites, 244.

Ferrisia (Ferrisiana) virgata, food-plants and distribution of, in W. Africa, 196, 366; transmitting virus disease of cacao in Gold Coast, 66, 196; on Lantana

camara, etc., in Egypt, 227.

Ferrous Sulphate, costs of using mixture containing, against Aglaope infausta, 169; as a safener for arsenicals, 62.

ferruginea-striata, Aphis.

ferrugineus, Dacus (Strumeta).

Ferula, 302.

festucae, Macrosiphum (Myzus).

Ficus, Parlatoria chinensis on, in California, 258; Pseudaonidia trilobitiformis on, in Belgian Congo, 365; species of, infested by Pulvinaria psidii, in Egypt, 228. (See Fig.)

ficus, Ceroplastes; Chrysomphalus.

Fiddlewood (see Citharexylum spinosum). Fidia viticida, sprays against, on vines in U.S.A., 28.

Fig, Asterolecanium pustulans on, in Egypt, 227; experiments with Anastrepha spp. and, in Mexico, 395, 397; Ceroplastes rubens on, in New Caledonia, 161; Asterolecanium pustulans on, in Nyasaland, 86.

Fig Moth (see Ephestia cautella).

Fiji, miscellaneous pests in, 161, 201, 202, 290, 291; notice of lists of insect pests in, 136; natural enemies and biological control of insect pests in, 161, 202, 269, 290, 291; insects attacking noxious weeds in, 136, 201, 291; beneficial insects imported into other countries from, 202, 261, 291.

Filbert (see Corylus avellana).

Filbert Weevil (see Curculio uniformis).

Filbert Worm (see *Cydia latiferreana*). Finland, forest pests in, **7**, **11**, **12**; parasites of bark-beetles in, **11**; bibliography of economic entomology in, **406**.

Firebrat (see *Thermobia domestica*). Fish, Dried, insects infesting, **82**, **102**.

Fish Meal, as bait for Anthrenus fasciatus, 73; development of Tineola biselliella on, 351.

Fish Oil, in nicotine-sulphate spray, 142; causing loss of rotenone in stored derris dusts, 152.

flammea, Panolis.

flava, Sipha,

flavedana, Platynota.

flavescens, Sitona.

flaviconchae, Apanteles.

flavipes, Karnyothrips; Reticulitermes.

flaviventris, Cremastus.

flavus, Metaphycus (Aphycus); Microterys. Flax, 278; relations between wireworms and, in Britain, 265, 294; insects on, in Canada, 98, 144; Aphthona euphorbiae on, in France, 331; Cnephasia wahlbomiana on, in Holland, 12; Prodenia litura on, in Nyasaland, 86; pests of, in U.S.A., 342, 343, 378; Jassids and virus diseases of, 378; cultivation of, as measure against Cephus cinctus, 98; Tyroglyphus farinae in stored seed of, 269.

Flea-beetle, Eastern Potato (see *Epitrix* tuberis).

Flea-beetle, Hop (see Psylliodes attenuata). Flea-beetle, Potato (see Epitrix cucumeris).

Flea-beetle, Tobacco (see Epitrix hirtipennis).

Flea-beetle, Turnip (see Phyllotreta nemorum).

Flea-beetle, Western Potato (see Epitrix subcrinita).

Fleahopper, Cotton (see Psallus seriatus). floralis, Hylemyia (Phorbia); Plagionotus.

Florida, pests of Citrus in, 211, 341; Batrachedra on coconut in, 97; Crambus haytiellus damaging grass in, 115; pests of solanaceous plants in, 41, 54, 215, 220; sugar-cane pests in, 116, 177, 260, 261; termites in, 69; Aphis gossypii transmitting lily rosette virus in, 174; natural enemies and biological control of insects in, 115, 177, 260, 261, 313; Trichopoda pennipes introduced into New South Wales from, 322.

Florida Red Scale (see Chrysomphalus ficus).

floridensis, Ceroplastes.

Flour, pests in, and their control, 104, 129, 130, 232, 235, 236, 237; sorption of fumigants by, 236, 237; effect of fumigation and insect infestation of wheat on qualities of, 237, 320, 403; insects reared on, 156, 181, 235, 351; in baits, 144, 145, 259, 343; as carrier for dusts, 3, 262, 283; as an adhesive for sprays, 271, 272.

Flour Beetle, Confused (see *Tribolium confusum*).

Flour Beetle, Rust-red (see Tribolium castaneum)

Flour Moth (see Ephestia kuehniella). Fluoranthene, protecting wood from ter-

Fluorene, protecting wood from termites,

Fluorine, problem of residues of, 79, 149,

forbesi, Cerosipha (Aphis).

Forcipomyia quasi-ingrami, and pollination of cacao in Trinidad, 78.

forestan, Rhopalocampta.

Forests, pests of, in S. Africa, 83; pests of, in Australia, 32, 162, 377, 378; pests of, in Britain, 359, 360, 404; pests of, in Bulgaria, 119; pests of, in Canada, 8, 71, 142, 143, 154, 155, 180, 223, 313, 314, 353, 354, 398; pests of, in Czechoslovakia, 8; pests of, in Finland, 7, 11, 12; pests of, in Germany, 6, 7, 8, 9, 10, 14, 15, 122, 231; pests of, in Holland, 7; pests of, in Hungary, 18; pests of, in India, 7, 385; pests of, in Jugoslavia, 5; Melolontha in, in Latvia, 7; pests of, in Mauritius, 371; Lymantria dispar in, in Morocco, 310; Sirex noctilio in, in New Zealand, 269; Sphinx pinastri in, in Poland, 7; pests of, in Portugal, 309, 310; pests of, in Queensland, Enarmonia diniana in, in Siberia, 400; pests of, in Spain, 357; pests of, in Sweden, 19, 32; pests of, in Switzerland, 194; pests of, in Turkey, 7; pests of, in U.S.A., 4, 31, 40, 43, 71, 107, 110, 173, 180, 238, 270, 271, 282, 286, 287, 310, 313, 314, 352, 354, 379, 398; symposium on pests of, 6-9.

Forficula, tests of insecticides on, 197, 198. Forficula auricularia, fungus infesting, in Britain, 232; baits for, in Ontario, 141.

Formaldehyde, tests of solutions containing, to protect fur, etc., against Tineola biselliella, 266, 267.

Formica, pollinating Taraxacum saghyz in Kazakstan, 245.

Formica rufa, forms, rearing and value of, against forest pests in Germany, 15, 16, 122.

Formica rufa rufa, 16.

Formica rufa rufo-pratensis, 16.

formicarius, Cylas; Stictococcus. formosanus, Idiopterus (Micromyzus).

fornicata, Phytodecta.

fornicatior, Xyleborus fornicatus.

Fornicia ceylonica, parasite limiting value of, against Limacodids in Ceylon, 300. foveicollis, Aulacophora.

Fowls, use of, to destroy insect pests, 134, 159, 302, 303; favouring infestation of *Citrus* by Coccids, 268.

Foxglove (see Digitalis purpurea).

fragaefolii, Capitophorus.

fragariae, Ancylis (see A. comptana); Capitophorus; Tarsonemus (see T. pallidus); Tyloderma.

decemlineata on Leptinotarsa France. potato in, 157, 158, 298, 331; miscellaneous pests in, 297, 331, 357; orchard pests in, 5, 125, 331, 354-356; natural enemies and biological control of insects in, 298, 354-356, 357; parasites of Cydia pomonella imported into Canada from, 356; Bruchophagus gibbus intercepted in Brazil from, 375.

Frankliniella, sprays against, on Parthenium argentatum in California, 104. Frankliniella helianthi, dust against, on

beans and peas in California, 338. Frankliniella minuta, on Parthenium argentatum in California, 104.

Frankliniella moultoni, on Parthenium argentatum in California, 104.

Frankliniella occidentalis, on lucerne and Parthenium in U.S.A., 101, 104.

fraterculus, Anastrepha. frenchi, Coptotermes.

Freon-12, 99.

Frianite, as diluent for cubé and derris dusts, 283.

frit, Oscinella.

Fritillaria pudica, transmission of virus diseases from lilies to, 2, 38.

Frogs, destroying beetles in wood stacks, 72.

frontalis, Dendroctonus.

frontatus, Encyrtus (Microterys).

frugiperda, Laphygma.

Fruit, Dried, pests in, and their control, 25, 102, 231, 296.

Fruit-flies, pits for disposal of fruit infested with, 134, 162.

Fruit-fly, Blueberry (see Rhagoletis pomonella).

fugax, Solenopsis. fulgida, Carneocephala.

fuliginosus, Stenocarus.

fullonia (fullonica), Othreis (Ophideres).

fulvicornis, Hoplocampa (see H. minuta). fulvoguttata, Melanophila.

fulvomaculatus, Calocoris.

fulvum, Stromatium.

fumida, Rodolia.

fumiferana, Harmologa (Archips).

Fumigants, sorption prolonging effect of, in type of container for grain and flour, 236, 237; recirculation of, 103, 104; methods of testing, 252.

funebrana, Cydia (Laspeyresia).

funebris, Bruchophagus (see B. gibbus). Fungi, Entomogenous, 20, 41, 100, 110, 119, 122, 204, 209, 232, 241, 242, 248, 258, 290, 303, 356, 364, 374, 381, 385, 389; utilisation of, 110, 128, 204, 364.

Fungi, Injurious, 8, 16, 35, 43, 54, 59, 85, 103, 150, 171, 173, 194, 204, 206, 207, 239, 240, 255, 272, 285, 317, 322, 326, 372; relation of insects to, 31, 40, 79, 110, 228, 286, 317, 354, 357, 389.

Funnel Ant (see Aphaenogaster longiceps).

fur. Ptinus. furcifera, Sogata.

furfura, Chionaspis.

Furniture, Coleoptera in, 167, 176, 403; Tineola biselliella damaging upholstered, 351.

Furs, insects damaging, 72, 163, 229, 351; proofing of, against clothes moths in Britain, 266.

fusca, Busseola.

fuscifrontia, Sesamia.

fuscipenne, Microplectron.

fuscum, Tetropium.

fuscus, Anthrenus. Fusicladium, 35.

G.

gahani, Pseudococcus.

Galeruca tanaceti, on potato in Denmark,

Galerucella luteola, favouring infestation of elm by Scolytus multistriatus in U.S.A., 43.

Galerucella tenella, bionomics of, on straw-

berry in Denmark, 381.

Galerucella xanthomelaena (see G. luteola). Galleria mellonella, experiments with Ernestia consobrina and, 305; distribution of DDT homologue in tissue of, 347.

Galtonia candicans, Myzus persicae transmitting virus to Ornithogalum from, in

U.S.A., 1.

Gammexane, properties and value of, as an insecticide, 257, 258; y isomer of benzene hexachloride (q.v.), 257.

Garden Centipede (see Scutigerella imma-

culata).

Garlic, pests and virus diseases of, in U.S.A., 70, 338.

Gasterocercodes brasiliensis (see Eutinobothrus).

Gastroides viridula, used for tests, 253.

gayneri, Autoba.

gei, auct., Macrosiphum (see M. solanifolii).

Gelatine, Tineola biselliella not develop-

ing on, 351.

Gelechia desiliens, bionomics of, on Platanus in U.S.A., 353.

gelechiae, Bracon (Microbracon).

Gelignite, for destroying termite nests, 73. Gelis, hyperparasite of Polychrosis viteana in U.S.A., 28.

gemellata, Rossi, Chrysomela quadrigemina).

geminata, Solenopsis.

geniculata, Blennocampa; Pristiphora.

geniculati, Stenodiplosis.

geniculatus, Uloborus.

Geocoris punctipes, predacious on Epitrix hirtipennis in U.S.A., 41.

Georgia, biological control of Pseudococcus comstocki on apple in, 208, 209; tobacco pests in, 54, 215; vetch Bruchid in, 259.

Geraniol, in baits for Popillia japonica, 55.

germanica, Blattella.

Germany, Acarapis on bees in, 6: importance of Aphid honeydew for bees in, 6; forest pests in, 6, 7, 8, 9, 10, 14, 15, 16, 122; Leptinotarsa decemlineata in, 157, 158; miscellaneous pests in, 36, 182, 231; orchard pests in, 34, 35, 117, 119; insects on *Ribes* spp. in, 14, 308; vine pests in, 13, 14; pests of stored products in, 228, 229, 230, 231, 232, 255; Coleoptera in building timber in, 18, 228, 229, 255; predacious ants and their value in, 15, 16, 122; other natural enemies of insect pests in, 8, 14, 19, 119, 120, 225; characters of Elaterid larvae of, 10, 11.

Gesapon, 288.

Gesarol Preparations, 36, 125, 288, 345.

gibbosus, Ligyrus.

Gilpinia hercyniae (on spruce), in Canada, 155, 223, 398; in Maine, 47; natural enemies, biological control and disease of, 47, 155, 223, 224, 398.

Gilpinia pallida, disease of, in Europe, 398. Gilpinia polytoma, disease of, in Europe,

Gipsy Moth (see Lymantria dispar). gladioli, Taeniothrips (see T. simplex).

Gladiolus, Taeniothrips simplex on, in Quebec, 224; Aphids and virus disease of, in U.S.A., 2.

Glenea, on cacao in Belgian Congo, 366. Gliricidia sepium, insects on, in Hawaii,

Glischrochilus hortensis, on raspberry and strawberry in Denmark, 382.

glomeratus, Apanteles.

Gloriosa, susceptibility of, to Aphid-borne virus diseases of Lilium, 2.

Glossolepis pilgeriana, Mesohomotoma tessmanni on, in Gold Coast, 66.

Glue, in DDT spray, 213; insects attracted by adhesive containing, 272.

Glyceryl Mono-oleate, effect of, in retarding penetration of oil into plant tissue, 253.

Glyceryl Oleate, effect of, in retarding penetration of oil into plant tissue, 253. Glycol Dichloride, proprietary spray pre-

paration containing, 51.

Glycol Mono-oleate, effect of, in retarding penetration of oil into plant tissue, 253. Glycyphagus destructor, factors affecting production of hypopi by, in stored grain, 330.

Glyoxal (see Dialdehyde).

Glyphipteryx achyloessa, bionomics and control of, on Dactylis glomerata in New Zealand, 269.

Glypta mutica, parasite of Polychrosis viteana in U.S.A., 28.

Glypta rufiscutellaris, introduction of, into Uruguay against Cydia molesta, 95. Gnathocerus cornutus, experimentally in-

festing dehydrated vegetables, 102. Gnathotrichus spp., in fire-killed Douglas

fir in U.S.A., 40.

gnava, Carcelia.

Gnorimoschema gudmannella, on Capsicum in Mexico and Porto Rico, 97.

Gnorimoschema ocellatella, bionomics and control of, on beet in Spain, 120, 121.

Gnorimoschema operculella (Potato Moth), in S. Africa, 326; in Argentina, 4; in Australia, 31, 93, 184–187, 322, 323, 404; in Eritrea, 379; in India, 300; in U.S.A., 51, 146, 214; bionomics of, 184, 185; parasites and biological control of, 322, 379; measures against, on field potatoes, 51, 185, 186, 322, 323; method of estimating damage to potato foliage by, 93; measures and experiments against, in stored potatoes, 31, 186, 187, 214, 300, 326, 404; use of, for rearing Macrocentrus ancylivorus, 218, 219.

Gnorimoschema plaesiosema, experiments against, on tomato in New Zealand, 269.

Gold Coast, pests and disease of cacao in, 65, 66, 195, 196, 383; pests of stored cacao in, 66; beneficial insects, 65, 196. Golden-rod (see Solidago).

Golf Courses (see Turf).

Gonimbrasia tyrrhea, on pines and Acacia in S. Africa, 83.

Gonipterus scutellatus, on Eucalyptus in S. Africa, 83.

Gooseberry, pests of, in Canada, 64; Nematus ribesii on, in various countries, 14, 64, 360, 381; Thomasiniana theobaldi not ovipositing on, 308.

Gorse (see *Ulex europaeus*) gossypiella, Platyedra (Pectinophora). gossypii, Aphis; Phenacoccus. gossypiperda, Bemisia (see B. tabaci).

Gossypium (see Cotton). gothica, Neokolla.

Goulac (see Sulphite Lye).

gracilis, Aspidiotus; Taeniocampa.

Grain Beetle, Saw-toothed (see Oryzaephilus surinamensis).

graminum, Toxoptera. Granadilla (see Passiflora). granaria, Calandra (Sitophilus). granarium, Macrosiphum (see M. avenae); Trogoderma.

granarius, Aphidius.

Granary Weevil (see Calandra granaria).

grandiosella, Diatraea. grandis, Anthonomus.

granella, Tinea.

granicola, Liposcelis.

Berry Moth Grape (see Polychrosis viteana).

Grape Leafhopper (see Erythroneura comes).

Grapefruit, pests of, in Bermuda, 172; Coccids on, in Belgian Congo, 364; Anastrepha ludens on, in Mexico, 395; pests of, in U.S.A., 211, 397.

Graphognathus (see Pantomorus). Grapholitha molesta (see Cydia).

Graptolitha, on pear in Washington, 285.

Grass, Bermuda (see Cynodon dactylon).

Grass, Carpet (see Axonopus compressus). Grass, Cocksfoot (see *Dactylis glomerata*).

Grass, Couch (see Agropyrum).

Grass, Crab (see Digitaria sanguinalis). Grass, Japanese (see Zoysia japonica).

Grass, St. Augustine (see Stenotaphrum secundatum).

Grass, Timothy (see Phleum pratense).

Grasselli Spreader-sticker, 339.

Grasses, pests of, in Australia, 161, 269; pests of, in Brazil, 93; pests of, in Britain, 92, 155, 156; Melanoplus mexicanus damaging, in Br. Columbia, 154; new Cecidomyiid on, in Mauritius, 74; pests of, in New Zealand, 183, 269, 375; Laphygma exempta on, in S. Rhodesia, 126; pests of, in U.S.A., 114, 115, 147, 191, 192, 215, 219; cereal pests on, 11, 93, 98, 147, 191, 199, 301. Grasshopper, Two-striped (see *Melano-*

plus bivittatus).

Greece, pests of melon in, 16.

gregaria, Schistocerca.

Grenada, parasite of Diatraea saccharalis in, 176.

Gretchena bolliana, on pecan in Texas, 347. Grevillea robusta, reducing infestation of tea by Paratetranychus bioculatus in Nyasaland, 86.

grisescens, Lydella stabulans.

Ground-nuts, Diabrotica speciosa on, in Argentina, 242; Elasmopalpus lignosellus on, in Arkansas, 191.

Ground-nuts (Stored), insects damaging, 31, 370.

Gryllotalpa, feeding habits of, in Egypt, 226.

Gryllotalpa gryllotalpa var. cophta, 226. Gryllulus (Gryllus) assimilis, Bufo cognatus destroying in N. Dakota, 96.

Gryllulus domesticus, gammexane against, 257.

Guadeloupe, parasite of Leucoptera coffeella introduced into Porto Rico from, 97.

Guaiacum spp., tests with gum from, against termites, 244.

Guam, Lachnosterna mindanaona in, 163.

Guatemala, lists of pests in, 289.

Guava, pests of, in India, 386, 393; Anastrepha spp. on, in Mexico and S. America, 395, 397; Helopeltis sanguineus on, in S. Rhodesia, 126; experiments with fruit-flies and, 52, 161.

Guayule (see Parthenium argentatum).

gudmannella, Gnorimoschema.

guerini, Tragocephala:

Guerinia spinosula (see Tachina rustica). Guiana, British, ant associated with Coccids on Axonopus in, 147; Pentatomid on rice in, 147; parasite of Saissetia nigra in, 374.

Guiana, Dutch, ants associated with Coccids transmitting phloem necrosis of coffee in, 147; Pentatomid on rice

in, **147**,

Guinea, Spanish, Bryocoropsis laticollis in, 196; new Coccid on coffee in, 166. guineensis, Stictococcus.

Gum Arabic, and mercurous chloride, against Hylemyia brassicae, 178.

gurneyi, Coccophagus.

guttifera, Teragra.
Gymnococcus (Coccid) (see Ovaticoccus).
Gymnonychus californicus (see Pristiphora).

Gynandropsis pentaphylla, Myzus persicae

on, in Nyasaland, 85.

Gypsum, as a carrier for dusts, 144, 197, 256; accelerating action of insecticides on *Drosophila*, 144.

H.

Habrocytus phycidis, parasite of Coleophora laricella in Ontario, 143.

Habrolepis rouxi (parasite of Coccids), in S. Africa, 107; establishment of, in California, 107.

haemorrhoidalis, Heliothrips.

hageni, Conwentzia.

Hairy Woodpecker (see *Dryobates villosus*). Haiti, new ant associated with Coccids on coffee in, **147**.

Halictus, pollinating Taraxacum saghyz in Russian Union, 245.

Halogeton glomeratus, relation of Eutettix tenellus and curly-top to, in U.S.A., 281. Haltica chalybea, sprays against, on vines

in U.S.A., 28.

Haltichella, parasite of Batrachedra in Porto Rico, 97.

Halticoptera aenea, parasite of Agromyza virens in California, 104.

Halticus saltator, on beans in Holland, 12. Halyzia, associated with mildew on pear in India, 389.

Halyzia sanscrita, not parasitised by Perilitus coccinellae, 369; possibly disseminating fungi on Dalbergia sissoo in India, 389.

hamata, Phorocera.

hambletoni, Ecpantheria.

hampei, Stephanoderes (Hypothenemus). Haplogonatopus vitiensis, parasite of Peregrinus maidis in Hawaii, 250.

Haplophyton cimicidum, toxicity of decoction of, to fruit-flies in Mexico, **397.**

Haploptilia (see Coleophora).

Haplosterna delegorguei (see Encosternum). Haplothrips cahirensis, predacious on Asterolecanium pustulans in Egypt, 227. harcyniae, Pissodes,

Harmologa fumiferana (Spruce Budworm), outbreak of, on Abies balsamea in Canada, 143, 313; in U.S.A., 313, 314; bionomics, food-plants and distribution of races of, 313, 314.

Harpalus rufipes, on strawberry in Britain,

hauseri, Macronaemia.

Hawaii, new mite on Litchi chinensis in, 111; miscellaneous pests in, 202, 217, 250, 251; Pseudococcus brevipes on pineapple in, 249, 345; keys to Lepidopterous larvae in, 264; beneficial insects and biological control in, 61, 250, 251, 374; pests intercepted in quarantine in, 223; summary of plant quarantines in, 145; risk of introduction of pests from New Caledonia into, 290.

hawaiiensis, Coccophagus. Hawthorn (see Crataegus). haytiellus, Crambus.

Hazel (see Corylus avellana).

Heat, uses of, against pests, 25, 72, 103, 229, 255, 286, 375, 403; apparatus for treating buildings with, 255; (vapour), treatment of fruit with, against *Anastrepha* spp., 396. (See Solar Heat and Water, Hot.)

Heather, pests of, in Britain, 232, 233. Heather-beetle (see *Lochmaea suturalis*). *Hedera helix*, new mite on, in California,

111. hedericola, Phytoptus.

helianthi, Frankliniella.

Heliothis armigera, in Bengal, 128; in Brazil, 278; in Queensland, 32, 87; in S. Rhodesia, 87, 126; in Russia, 306; in Tanganyika, 88, 89; in U.S.A., 57, 101, 211, 260, 261, 335, 336, 337; on cotton, 32, 57, 87, 88, 211, 261, 278, 306, 335, 336; on maize, 88, 89, 101,

260, 337; on other plants, 102, 126, 128; parasite of, 281; measures and experiments against, 32, 57, 87, 211, 260, 278, 279, 306, 335, 336.

Heliothis ononis, on flax in Canada, 143. Heliothis virescens, in S. America, 278; in U.S.A., 278, 280, 281; in W. Indies, on tobacco, on cotton, 278; 278, 280, 281; natural enemies of, 278, 281; measures and experiments against, 278, 280.

Heliothrips haemorrhoidalis, on pines in S. Africa, 83; bionomics of, on Cinchona and cacao in Belgian Congo, 363,

364; sprays against, 364.

Hellula phidilealis, bionomics and control of, on crucifers in Brazil, 95.

Hellula undalis, on cabbage in S. Rhodesia, 126.

hellulae, Bracon (Microbracon).

Helochara delta, transmitting virus disease

of vines in U.S.A., 172.

Helomyia lateralis (parasite of Eurygaster), bionomics and suggested utilisation of, in Russian Union, 301, 302.

Helopeltis, in W. Africa, 91, 196, 367; in Tanganyika, 89; on tea in Travancore, 246; on cacao, 196; on cotton, 89, 91, 367; measures against, 91.

Helopeltis bergrothi, on cacao, etc., in W. Africa, 196, 366.

Helopeltis orophila, bionomics and control of, on Cinchona, etc., in Belgian Congo, 36-38.

Helopeltis sanguineus, bionomics of, on cacao, cotton, etc., in W. Africa, 383; on guava and avocado in S. Rhodesia, 126.

Helophorus nubilus, cultural practices affecting, on wheat in Britain, 297.

helvolus, Metaphycus.

Hemadas nubilipennis, measures against, on blueberry in U.S.A., 402.

Hemerobius pacificus, predacious on Sais-

setia nigra in California, 374. Hemiberlesia rapax, on orange and vines

in Argentina, 4. Hemichionaspis theae (see Pinnaspis).

Hemileuca oliviae, parasite of, on grasses in U.S.A., 215.

Hemiptarsenus semialbiclavus, parasite of Phytomyza spicata in Fiji, 290.

hemipterus, Carpophilus.

Hemisarcoptes malus (coccisugus), attacking Coccids in France, Germany and U.S.A., 357.

hemisphaerica, Saissetia (see S. coffeae). Hemitarsonemus (see Tarsonemus).

Hemiteles, hyperparasite of Cydia pomonella in France, 355; parasite of Gilpinia hercyniae in Quebec, 224.

Hemiteles humeralis, hyperparasite Polychrosis viteana in U.S.A., 28.

Hemiteles tenellus, parasite of Coleophora laricella in Ontario, 143; hyperparasite of Polychrosis viteana in U.S.A., 28.

Henbane (see Hyoscyamus niger).

Henequen (see Agave fourcroides). Hepialus humuli, in harvested apples in Britain, 359; food-plants of, 359.

heracleana, Depressaria. heraclei, Brachycolus.

Depressaria hera-Heracleum lanatum, cleana on, in Br. Columbia, 333.

herbarium, Catorama.

Hercothrips fasciatus, on Argemone in Hawaii, 251, 252; possibly introduced on oranges from California, 252.

hercyniae, Gilpinia. hesperidum, Coccus.

Hesperophanes cinereus, in timber Spain, 167, 168; life-cycle of, 168.

hesperus, Lygus.

Hessian Fly (see Mayetiola destructor). Heterobostrychus aequalis, in timber in Ceylon, 300; on Eucalyptus in Mauri-

tius, 371. Heterobostrychus brunneus, on cassava in

Belgian Congo, 367.

Heteronychus licas, on maize in S. Rhodesia, 126.

sanctae - helenae, plants Heteronvchus attacked by, in N. S. Wales, 161; on vegetables in New Zealand, 93.

Heterotoma merioptera (see Capsus). Hexachlorcyclohexane (see Benzene Hexa-

chloride).

Hexachlorphenol, protecting wood against termites, 245.

Hexacola (parasite of Oscinella frit), not ovipositing on Drosophila, 75.

Hexadecyl Nitrile, mixture containing, not toxic to insects, 198.

Hexarthrum ulkei (see Rhyncolus).

2-n-Hexyl-pyridine Sulphate, effect of dust of, against Hylemyia brassicae, 276.

Hibiscus spp., experiments with Ecpantheria hambletoni and, in Brazil, 204; parasitism of cotton pests on, in India, 21, 390.

Hide Beetle (see Dermestes maculatus). hieroglyphica, Coccinella; Neokolla.

Hieroglyphus banian, on bamboos in India, 389.

hilaris, Bagrada.

himantopus, Athalia.

hippocastani, Melolontha.

hirtellus, Adoretus (see A. umbrosus); Collops; Ptinus. hirtipennis, Epitrix.

hirtum, Nicobium castaneum.

Histerids, survey of, associated with stored products, 135.

Histiostoma, predacious on Aulacophora foveicollis in Greece, 17.

histrionica, Murgantia.

Holcocera iceryaeella, on orange in Cali-

fornia, 113.

Holland, forest pests in, 7; Leptinotarsa decemlineata in, 157; miscellaneous pests in, 12, 253.

Hollyhock (Althaea), insects on, in India,

21, 390.

Homalodisca triquetra, bionomics of, on Agave fourcroides in Mexico, 131.

Homona coffearia, parasite of, on tea in Ceylon, 300.

homonae, Macrocentrus.

Honduras, Plaesius javanus imported into, against Cosmopolites sordidus, 261.

Honey, used for feeding insects, 144, 204,

215, 376, 390.

Hoop Pine (see Araucaria cunninghami). Hop Downy Mildew (see Pseudoperonospora humuli).

Hop Flea-beetle (see *Psylliodes attenuata*). Hop Powdery Mildew (see Sphaerotheca humuli).

philanthus, bionomics of, in Hoplia Britain, 155, 156.

Hoplocampa, benzene hexachloride against, on fruit trees in France, 331.

Hoplocampa minuta (fulvicornis), sprays against, on plum in Denmark, 380.

Hoplocampa testudinea (on apple), in Britain, 359; in Br. Columbia, 334; in Poland, 117; sprays against, 334, 359. Hops, pests of, in Britain, 359, 361;

Psylliodes attenuata on, in Germany, 231; pests of, in New York, 317.

Hordeum spp., Eurygaster integriceps on, in Uzbekistan, 301. (See Barley.)

Hormius pallidipes (see Parahormius). Horsehair, Anthrenus fasciatus damaging,

Horseradish, Chrysomelid on, in Den-

mark, 381. hortensis, Glischrochilus (Librodor).

horticola, Anomala (Phyllopertha).

howardi, Dysdercus. howitti, Aphodius.

humeralis, Hemiteles.

Humidity, influence of, on Glycyphagus destructor, 331; relations of insects to, 9, 13, 18, 20, 65, 73, 75, 76, 119, 123, 128, 135, 195, 261, 267, 377, 384, 406; regulation of, in insect rearing compartments, 216.

humuli, Hepialus; Phorodon.

Hungary, natural enemies of honeybees in, 6; miscellaneous pests in, 11, 122; lists of Hymenopterous parasites in, 18.

Hyacinth, effects of fumigants against Rhizoglyphus echinopus in imported bulbs of, in Sweden, 35; Aphids and virus diseases in, 1, 39...

hyacinthi, Rhizoglyphus (see R. echinopus).

Hyacinthus (see Hyacinth).

hyalinata, Diaphania.

Hyalomyia chilensis, introduction of, against Dysdercus in Porto Rico, 312.

Hyalopterus arundinis (Mealy Aphis), sprays against, in Br. Columbia and Denmark, 153, 380.

Hyalopterus pruni (see H. arundinis).

Hydnocera scabra, predacious on Cydia pomonella in California, 240.

Hydrochloric Acid, use of, to stimulate hatching of silkworm eggs, 324, 325; in formula for treating furs against

Tineola biselliella, 266, 267.

Acid, fumigation with, Hydrocyanic against Coccids, 125, 204, 227, 255, 256, 259, 271, 346, 364; liquid form of, lowering temperature fumigation tent, 346; new dosage tables for, 256; not harmful to Meta-phycus helvolus, 259; against Halticid on artichoke, 168; against Hylotrupes bajulus, 229; against pests in bulbs, 24; against pests of stored products, 127, 130, 231, 235, 236, 259, 295; not affecting vitamin B₁ in flour, 259; sorption of, by cereals, soils and foliage, 128, 236, 271; methods of generating and applying, 127, 130, 255, 256, 271; and injury to plants, 24, 168, 256. (See Calcium and Potassium Cyanides.)

Hydroecia immanis, on hops in New York,

Hydrogen Phosphide, fumigation with, against Calandra granaria, 229, 230.

Hydrogen-ion Concentration, effect of, on insecticides, 58, 396.

 α - bis (Hydroxyphenyl) $\beta\beta\beta$ - trichlorethane (see DDT Analogues).

Hylastes opacus, measures against, on conifers in S. Africa, 83, 84.

Hylastes palliatus, parasites of, in Finland, 12.

Hylastes rufibes, fumigation against, transmitting Dutch elm disease in U.S.A.,

Hylemyia antiqua (on onion), in Br. Columbia, 154; in California, 338; in Russia, 304; measures and experiments against, 154, 304, 338.

Hylemyia brassicae (on crucifers), in Br. Columbia, 154; in Russia, 304; in Switzerland, 288; in U.S.A., 100, 177, 178, 276; bionomics of, 178; measures and experiments against, 154, 178, 276, 288, 304.

Hylemyia cilicrura, on beans in Holland, 12; bionomics and control of, in U.S.A.,

179, 338.

Hylemyia coarctata, bionomics and control of, on cereals in Britain, 297.

Hylemyia floralis, bionomics of, on crucifers in Scotland, 121.

Hylemyia latipennis, parasite of Gilpinia latipennis in Quebec, 224.

Hylemyia sepia, on wheat in Britain, 297. Hylotrupes bajulus (in timber), in Germany, 18, 19, 228, 229, 255; in Spain, 167, 168; in U.S.A., 176; bionomics and physiology of, 18, 19, 167, 168; Clerid predacious on, 19, 229; measures and experiments against, 228, 229, 255. Hylurgopinus (see Hylastes).

Hylurgops (see Hylastes).

Hymenia recurvalis (fascialis), parasite of, 281.

Hymenoptera, Parasitic, olfactory responses in relation to rearing of, 281; rearing of, on unnatural hosts' 74, 75; importance of diapause in, in work on biological control, 105; lists of, in biological control, 105; Hungary, 18.

Hyoscyamus niger, Myzus persicae and virus disease of, in New Zealand, 247.

Hypera postica (see H. variabilis).

Hypera punctata, food-plants and disease of, in California, 241.

Hypera transsylvanica, measures against, on lucerne in Russia, 305, 306.

Hypera variabilis, on lucerne in U.S.A., 40, 41, 101, 206; other food-plants of, 41; natural enemies and biological control of, 41, 206; dust against, 101.

hyperici, Agrilus; Chrysomela (Chryso-

lina).

Hypericum perforatum, use of insects against, in New Zealand and Australia, 269, 321.

Hyperimerus, parasite of Clausenia purpurea in Virginia, 110.

Hyphantria cunea, measures against, on blueberry in U.S.A., 402. Hypnoides riparius, bionomics of, in

pastures in Britain, 92, 405.

hypnorum, Tachyporus.

Hypolixus truncatulus, Spathius critolaus reared on, in India, 21.

Hypothenemus (see Stephanoderes).

I.

Icerya montserratensis, natural enemies and biological control of, on Citrus in

Ecuador, 95, 96.

Icerya purchasi, in Belgian Congo, 364, 365; in France, 357; in India, 246, 384, 385, 391; in W. Indies, 110; on Citrus, 110, 246, 357, 364, 385; on other plants, 110, 357, 384, 385, 391; bionomics of, 385; natural enemies and biological control of, 110, 357, 364, 385.

Icerya tremae, on cacao in Belgian Congo

366.

iceryae, Cryptochetum. iceryaeella, Holcocera.

Ichneumon caudatus (see Ephialtes).

Idaho, Heliothis armigera on maize in, 260, 337; miscellaneous pests in, 220, 259, 284; Tettigoniids in, 113, 114.

Idiopterus formosanus, food-plants of, in Hawaii and California, 251, 338.

ignorata, Dasyneura. ignota, Empria.

ilicis, Paratetranychus.

Illeis cincta, feeding habits of, in India, 389. Illinois, Ctenolepisma urbana in, 106;

orchard pests in, 315, 316. illutana, Cydia (Laspeyresia).

imaginis, Thrips. imitata, Melanolophia

immaculata, Scutigerella.

immanata, Disstroma (Cidaria) citrata.

immanis, Hydroecia. impressiceps, Pheidole.

inara, Serrodes (see S. partita).

Inareolata (see Angitia)

incana, Scaptomyzella (Scaptomyza).

inconsequens, Taeniothrips.

inda, Euphoria.

India, pests of bamboo in, 388; cotton pests in, 20, 164, 362, 386, 388, 389, 390; forest pests in, 7, 390; fruit-piercing moths in, 386, 391; sugar-cane pests in, 390, 391, 406; pests of stored products in, 72, 73, 164, 300, 362; pests of timber in, 71, 72, 73, 371, 372; Schistocerca gregaria in, 136, 388; termites in, 73; miscellaneous pests in, 374, 384, 388, 391, 393; fungus-eating Coccinellids in, 389; natural enemies and biological control of insects in, 72, 164, 374, 385, 386, 389, 390, 391; Teleonemia scrupulosa imported into, against Lantana, 264; plant quarantine restrictions in, 111. (See also under the various Provinces.)

Indian Ink, used to mark insects, 390. Indiana, apple pests in, 218; wheat pests in, 103.

indica, Margaronia.

indicum, Syntomosphyrum (Melittobia). Indo-China, Pyrausta coclesalis on sugar-

cane in, 388. inermis, Stictocephala.

infausta, Aglaope.

infelix, Eucomys (Encyrtus).

Infra-red Rays, experiments with insects and, 53.

infuscata, Ormenis.

infuscatus, Limonius.

Infusorial Earth, against Gnorimoschema

operculella, 31. Inga jinicuil, Anastrepha spp. on, in Mexico, 395, 397.

Inositol, 257.

inquisitor, Ephialtes (Epiurus).

Insecticides, tests of organic compounds as potential, 60, 98, 100, 106, 107, 137,

176, 197, 198, 214, 245, 256, 275, 276, 279, 331, 344; bibliography on plants as sources of, 192; technique and analysis of tests of, 60, 64, 188, 197, 214, 328, 345; review of literature on, against soil pests, 393; possible increase of pests by routine application of, 254; general papers on, 57, 98-100, 264, 343. Insects, theories of causes of outbreaks of,

7, 13, 254; graphic method of indicating population of, 56; sense physiology in relation to control of, 253; tracheal system of, in relation to sprays, 252.

insidiosus, Orius.

insignicola, Physokermes (Kermes). instigator, Pimpla (Coccygomimus).

insulana, Earias.

insularis, Kalotermes (Calotermes); Mirax; Solubea.

integriceps, Eurygaster. intermedius, Dysdercus.

interpunctella, Plodia.

Inula viscosa, Dacus oleae on, in Italy, 118. Iodine, compounds of, as potential insecticides, **176.**

ionephela, Epichorista.

Iowa, pests of cereals in, 57, 101, 259; Cydia molesta in, 258; miscellaneous pests in, 175, 353.

Ipobracon amabilis, introduction against Diatraea saccharalis in Porto Rico, 312.

Ipomoea, Cylas formicarius on, in Louisiana, 216. (See Sweet Potato.)

Ips spp., parasites of, in Finland, 12; infesting weakened spruce in Germany, 14.

Ips bidentatus, 12.

Ips chalcographus, 12, 14.

Ips proximus, 12. Ips quadridens, 12.

Ips typographus, 12, 14.

Iraq, Dociostaurus maroccanus in, 199-201; Eurygaster integriceps in, 302.

Iridomyrmex detectus, measures against, in Victoria, 322.

Iris, effects of fumigation against pests of, in Washington, 24.

Isaria farinosa, infesting Operophtera brumata in Germany, 119.

ischioxanthos, Eurytoma.

Ischnus, parasite of Cydia pomonella in California, 240.

Ischnus polychrosidis, parasite of Polychrosis viteana in U.S.A., 28.

isipingoensis, Coccophagus.

Isoamyl Alcohol, Blepharoneura attracted by bait containing, 216.

Isophorone, 198.

Isotomurus, on Thuja seedlings in Denmark, 382.

Isotomurus palustris, 382.

italicus, Calliptamus (Caloptenus).

Italy, miscellaneous pests in, 6, 118, 120, 125; parasites of Dacus oleae in, 118; Exochomus quadripustulatus introduced into California from, 282.

Ivy (see Hedera helix).

Ixia, Myzus persicae and virus disease of, in U.S.A., 2.

J.

jaboticabae, Targionia (Aspidiotus). jacksoni, Pulvinaria.

janata, Achaea.

Japan, beneficial insects introduced into U.S.A. from, 208, 261, 374, 375.

Japanese Beetle (see Popillia japonica). Japanese Grass (see Zoysia japonica). japanica, Diestrammena.

japonica, Diestrammena (see D. japanica); Popillia.

Jarring, to dislodge insects from trees, 10,

Jasmine, Coccid on, in Brazil, 205. javanus, Plaesius.

Jobo (see Spondias).

juglans-regiae, Quadraspidiotus (Aspidiotus).

Jugoslavia, forest pests in, 5.

julianus, Strategus.

junci, Lixus.

juniperina, Contarinia.

Juniperus bermudiana, pests of, in Bermuda, 289, 290.

Juniperus communis hibernica, seasonal reaction of, to methyl bromide, 50.

Jussieua (Jussiaea), Helopeltis sanguineus on, in Nigeria, 383.

K.

Kale, Hylemyia floralis on, in Scotland, 121.

Kalotermes approximatus, fungi infesting, in Bermuda, 290.

Kalotermes brevis, tests of timber preservatives against, in Porto Rico, 243, 244.

Kalotermes brouni, in New Zealand, 234; characters of, 234, 235.

Kalotermes condonensis, introduced into New Zealand in timber, 234; characters of, 234, 235.

Kalotermes insularis, introduced into New Zealand in timber, 234; characters of, 234, 235.

Kalotermes tuberculatus, introduced into New Zealand in timber, 234; characters of, 234, 235.

kaltenbachi, Myzus.

Kansas, Cydia spp. in, 258, 275; miscellaneous pests in, 53, 114, 117, 220, 237, 353; Melitara dentata on Opuntia in, 241. Kaolin, experiments with, against pests of stored products, 31, 187, 327, 328; as a carrier for dusts, 185, 187; reducing toxicity of derris dust, 78.

Kapok (see Ceiba pentandra).

Karnyothrips flavipes, predacious on Asterolecanium pustulans in Egypt, 227.

Keiferia lycopersicella (Tomato Pinworm), parasites of, in Hawaii, 250; sprays against, in Virginia, 51; on egg-plant,

Kentucky, Conotrachelus nenuphar on peach and plum in, 315; Tyloderma fragariae on strawberry in, 262.

Kenya Colony, outbreaks of Laphygma exempta in, 245; parasite of Saissetia

nigra in, 374.

Kermes insignicola (see Physokermes). Kerosene, in mixtures against Coleoptera in timber, 84; as a solvent, 84, 339, 350. Kerosene Emulsion (see Oil Emulsions). Kieselguhr, against Gnorimoschema oper-

culella, 31.

kirbyi, Oiketicus. kirkpatricki, Bracon (Microbracon).

koehleri, Trichogramma.

Kok-saghyz (see Taraxacum kok-saghyz). kotinskyi, Microterys.

Krytox, 59.

kuchingensis, Microgaster. kuehniella, Ephestia. kuvanae, Ooencyrtus (Schedius).

labdacus, Spathius. labradus, Zizera.

Labrorhycus prismaticus, parasite Pyrausta nubilalis in Quebec, 138.

lacerdae, Pachymerus.

Lachenalia, Myzus persicae and virus

disease in, in U.S.A., 1.

Lachnosterna, in Canada, 140, 180; U.S.A., 61, 101, 179, 259, 262; plants damaged by larvae of, 101, 140, 259, 262; bionomics of, 140, 179; parasites of, 61, 180; experiment with Neoaplectana spp. and, 61; measures against, 141, 262.

Lachnosterna anxia, measures against, in

Canada, 141.

Lachnosterna calceata, bionomics of, in Texas, 270.

Lachnosterna micans, bionomics of, in U.S.A., 270.

Lachnosterna mindanaona, bionomics of, in Guam, 163, 164.

Lachnus, auct. (see Cinara).

lacteus, Coptotermes. lactucae, Pemphigus.

Laemophloeus minutus (in stored wheat), in Australia, 319, 320; in Oklahoma,

bionomics of, 320; against, 69, 319.

laetatorius, Diplazon.

laetivirens, Myllocerus.

laevifrons, Perilampus.

laevigata, Atta. laevis, Paniscus.

laeviusculus, Paratriphleps.

laeviventris, Chlorion (Palmodes).

Lagernaria vulgaris, experiment Aulacophora foveicollis and, in Greece,

Lamachus lophyri, parasite of Gilpinia hercyniae in Quebec, 224.

laminatus, Trichodes.

Lampides boeticus (see Cosmolyce).

lanigerum, Eriosoma.

Lantana, Ferrisia virgata on, in Egypt, 227; utilisation of Teleonemia scrupulosa against, in Queensland and Fiji, 32, 291, 321; risk of liberating T. scrupulosa against, in India, 264; leaves of, not protecting stored potatoes from Gnorimoschema operculella, 301.

Laphygma exempta, on cereals and grasses in E. Africa and S. Rhodesia, 86, 126, 245; parasites and biological control of, in Hawaii, 250; bionomics and outbreaks of, 245, 246;

against, 246.

Laphygma exigua, on maize and lucerne in S. Rhodesia, 126; parasites of, 126,

Laphygma frugiperda, in Argentina, 117; measures against, on cereals, grasses and cotton in Brazil, 93; on sugar-cane in Porto Rico, 243; parasite of, 281; tests of insecticides on, 339.

lapidum, Petrobia.

Larch (see Larix).

Larch Casebearer (see Coleophora laricella).

laricella, Coleophora (Haploptilia).

laricinellae, Chrysocharis.

Larix, Coleophora laricella on, in Ontario, 142.

Larix sibirica, pests of, in Finland and Siberia, 12, 400.

larvarum, Tachina (Exorista).

Lasioderma serricorne, in Fiji, 290; imported into Germany, 230; in Gold Coast, 66; in S. Rhodesia, 127; in U.S.A., 102; in stored cacao, 66; in dried vegetables, 102, 290; in drugs, 230; in tobacco, 127, 230; measures and experiments against, 66, 127.

Lasiohelea spp., possibly pollinating cacao in Trinidad, 78.

Lasius, toxicity of gammexane to, 257. Lasius niger, suggested use of Solenopsis fugax against, 16.

Laspeyresia (see Cydia).

lata, Phytolyma.

461 INDEX.

lateralis, Helomyia; Phytomyza (Napo- 1 myza); Strophosomus.

laticeps, Tarsonemus.

laticollis, Baris; Bryocoropsis.

latiferreana, Cydia (Melissopus).

latipennis, Hylemyia.

latiscapus, Prophyscus.

latus, Tarsonemus (Hemitarsonemus).

Latvia, Melolontha spp. in, 7.

Lauryl Thiocyanate, 44. (See Thiocyanates.)

Lawns (see Turf).

Lead, Dermestes damaging, 230.

Lead Arsenate, 343; against cotton pests, 87, 202, 210, 211, 335; against orchard pests, 3, 4, 23, 24, 43, 94, 119, 148, 149, 150, 151, 171, 179, 191, 210, 212, 222, 258, 273, 274, 284, 285, 314, 316, 334, 359, 360, 380; against pests of small fruits, 262, 360, 402; against tobacco pests, 45, 54, 280; against pests of vegetables, 158, 168, 185, 212, 217, 269, 289, 293, 323; treatment of stored potatoes with, against Gnorimoschema operculella, 187; against pests of other plants, 24, 28, 62, 104, 210, 238; against pests in soil and turf, 63, 137, 183, 262; testing of, on *Drosophila*, 144; in baits, 183, 280; in bait-sprays, 87, 162, 396; dusting with, 3, 4, 24, 43, 45, 54, 62, 150, 185, 210, 211, 212, 222, 262, 274, 293, 335, 360; carriers for, 24, 54, 185, 212, 262, 274, 293; oil-impregnated dusts of, 150, 316; formulae for, in sprays, 23, 24, 28, 39, 43, 54, 94, 104, 148, 150, 158, 168, 171, 185, 191, 210, 212, 217, 218, 238, 262, 269, 284, 289, 314, 316, 323, 334, 359, 402; adhesives and spreaders for, 28, 43, 62, 87, 150, 151, 171, 323; apparatus for applying atomised sprays of, 254; and aluminium sulphate, 62; and copper compounds, 28, 54, 62; and DDT, 171; and ferrous sulphate, 62; lime in sprays containing, 39, 43, 62, 148, 191, 314; effect of lime in sprays following, 39; and limesulphur, 23, 171; and nicotine, 43, 149, 150, 210, 274, 334, 359; and oils, 24, 28, 94, 185, 210, 212, 262, 284, 316, 334; "inverted" (dynamite) spray mixtures of oils and, 150, 151; and Paris green, 45; and phenothiazine, 258; and sulphur, 3, 39, 43, 150, 212, 222, 274; and thiocyanate, 149; and zinc sulphate, 191, 210, 314; addition of bee repellents to sprays containing, 212; and injury to plants, 39, 52, 105, 217, 314; chemical phases of injury to plants by, 39, 62, 314; plants absorbing arsenic from soil treated with, 63; basic form of, 314, 315; other insecticides compared with, 279, 289, 323, 324, 335, 336.

Leafhopper, Aster (see Macrosteles divisus).

Leafhopper, Grape (see Erythroneura comes).

Leafhopper, Potato (see Empoasca fabae). Leather, insects damaging, 72, 290.

Lecaniobius cockerelli, parasite of Saissetia nigra in Br. Guiana and W. Indies, 374.

lecaniorum, Eucomys.

Lecanium, Burm. (see Coccus).

Lecanium, auct. (see Eulecanium).

Lecidea, attracting Dociostaurus maroccanus, **200.** lecontei, Neodiprion.

lectularius, Cimex.

leda, Monomorium rothsteini.

Leea guineensis, Helopeltis bergrothi on, in Nigeria, 196.

leguminicola, Dasyneura.

Lema melanopa, dust against, on wheat in Uzbekistan, 303.

Lema trilineata californica, on solanaceous plants in Hawaii, 250.

Lemon, 291; pests of, in California, 97, 342, 350, 397; effect of, on parasite of Aonidiella aurantii, 92, 93; (fruit), Aonidiella aurantii reared on, 346.

leona, Anomis.

lepida, Parasa.

Lepidium alyssoides, Eutettix tenellus on, in New Mexico, 314.

Lepidoptera, book on, in Brazil, 401.

Lepidosaphes beckii (on Citrus), bionomics and control of, in Bermuda, 172; fungi infesting, in Belgian Congo, 364; in N. S. Wales, 248; characters distinguishing Unaspis citri from, 133.

Lepidosaphes ulmi, in Canada, 64, 332; in Europe, 357; in U.S.A., 353, 357, 401; on fruit trees, 332, 357; on small fruits, 64, 401; on Platanus, 353; comparison of injury to apple by Aphids and, 332; mites attacking, 357; spray against, 353.

Lepisma saccharina, in buildings in Canada and U.S.A., 107; value of laminated

paper wrapping against, 272.

Leptinotarsa decemlineata (Potato Beetle), in Belgium, 157, 158; legislation against introduction of, into Britain, 232; in Br. Columbia, 154; in France, 157, 158, 298, 331, 332; in Germany, 157, 158; in Holland, 157; in Ivory Coast, 165; in Luxemburg, 158; in Spain, 165, 166; in Switzerland, 125, 157; in U.S.A., 48, 154, 180, 207, 260; bionomics of, 158, 166, 298; experiments with species and hybrids of Solanum and, 159; tomato, 165, 166; measures and experiments against, 15, 48, 154, 158, 159, 180, 207, 260, 278, 279, 331, 332; surveys of data on, 157-159.

Leptocoris trivittatus, observations on, in

Utah, 106.

lebtogaster, Dacnusa.

Leptohylemyia (see Hylemyia).

Leptomastidea abnormis (parasite of Pseudococcus citri), experiments with P. comstocki and, in U.S.A., 109.

Leptomastix, failure to establish Japanese species of, against Pseudococcus comstocki in U.S.A., 209.

Leptomastix dactylopii, parasite of Pseudococcus citri in California, 109; experiments with P. comstocki and, 109.

leptothecus, Apanteles.

Leptothrips mali, predacious on Saissetia nigra in California, 374.

Leptura obliterata (see Strangalia).

Leptura rubra, humidity requirements of, in timber in Germany, 18.

Lespedeza, favouring Conotrachelus nenu-

phar on peach in Illinois, 317.

Lestodiplosis, predacious on Cecidomyiids and mites in Britain, 170, 308; occurrence of, in stored seed, 170.

Lethane 60, 49, 104, 148, 149. (See Thio-

cyanates.)

Lettuce, Agriotes obscurus on, in greenhouse in Britain, 405; Thrips tabaci on, in Fiji, 161; pests of, in Hawaii, 250, 251; pests of, in U.S.A., 39, 71, 112, 337, 338; insects and virus diseases of, **39**, **71**; as food for *Bombyx mori*, **325**.

Leucaena glauca, Ferrisia virgata on, in

Nigeria, 196.

Leucinodes elegantalis, bionomics and control of, on tomato in Brazil, 94; precautions against introduction of, into Argentina, 94.

leucophthalmus, Philaenus.

Leucopoecila albofasciata, outbreak of, on

golf course in Missouri, 219.

Leucoptera coffeella (on coffee), bionomics of, in Brazil, 311; value and utilisation of Mirax insularis against, in W. Indies, 97.

leucopterus, Blissus.

Librodor hortensis (see Glischrochilus).

licas, Heteronychus.

Light, causing loss of rotenone in derris dusts, 152, 153; reactions of insects to, 74, 123, 124, 165.

Light-traps, for Lachnosterna anxia, 141; ineffective against Polychrosis viteana and Dermestes maculatus, 27, 83.

ligneus, Carpophilus.

Lignin, as carrier for cubé dust, 283.

lignosellus, Elasmopalpus.

ligustici, Otiorrhynchus.

Ligustrum, Coccids on, in France, 5, 357. Ligyrus gibbosus (californicus), on Parthenium argentatum in California, 105.

Lilies (Lilium spp.), Aphids and virus diseases of, in U.S.A., 1, 2, 38, 174; mites on, in Washington, 24; effects of methyl bromide on, 24, 64.

lilii, Macrosiphum. Lilium (see Lilies).

Lima Beans, 61; pests of, in Hawaii, 250, 251; pests of, in U.S.A., 54, 79, 260; effects of soil treatments on, 56, 63.

limacina, Caliroa (Eriocampoides).

Lime (Citrus), relation of Anastrepha spp. to, in Mexico, 395, 397.

Lime, as a carrier for dusts, 3, 45, 59, 60, 112, 144, 150, 152, 212, 262, 317; addition of, to arsenical sprays, 39, 43, 62, 148, 176, 191, 212, 314, 315; addition of, to other sprays (see also Bordeaux Mixture), 58, 169, 212, 227, 271, 272, 317; tests and use of, against pests of stored products, 300, 312, 327, 372; as a repellent, 17, 371, 372; benzene hexachloride decomposed by, 257; reducing toxicity of cryolite to Drosophila, 144; forms of, 3, 39; formulae containing, 3, 39, 43, 45, 58, 60, 112, 148, 150, 152, 169, 176, 212, 227, 262, 314.

Lime-sulphur, unsatisfactory against Capsids, 66; against Coccids, 22, 98, 133, 169, 227, 248, 284, 285, 401; as repellent against Dendroctonus spp., 43; varying effects of, against Paratetranychus pilosus, 124, 208, 234, 360; against other mites, 35, 285, 286, 361; against Paratrioza cockerelli, 217; against Scirtothrips citri, 98; analysis and insecticidal action of, 169; formulae for, 35, 124, 133, 185, 217, 227, 234, 248, 284, 361, 401; formulae containing, 24, 171, 285; and DDT, 171; and lead arsenate, 24, 171; and nicotine, 285, 359; and oil emulsions, 22, 285; and sulphur, 98; spreader for, 248; and injury to plants 66, 105, 185.

Limestone (see Calcium Carbonate). Limneria pattoni (see Eulimneria).

Limonite, 372.

Limonius, measures against, in U.S.A., 23, 55, 56, 104, 260, 337; populations of, in relation to crops, 23.

Limonius californicus, in U.S.A., 23, 55, 56, 104, 337; plants attacked by, 23, 104, 337.

Limonius canus, plants attacked by, in U.S.A., 23, 104, 337.

Limonius infuscatus, plants attacked by, in California, 104, 337.

Lindorus lophanthae, predacious on Saissetia nigra in California, 374.

lineare, Dolichosoma.

lineata, Celerio.

lineatella, Anarsia.

lineaticollis, Antestia.

lineatum, Theridium; Trypodendron.

lineatus, Agriotes; Sitona.

lineolata, Diatraea.

lineolatus, Adelphocoris.

463 INDEX.

Liparis chrysorrhoea, auct. (see Nygmia phaeorrhoea).

Liparis dispar (see Lymantria).

Liposcelis granicola, sp. n., bionomics of, in Britain, 181.

Liriomyza solani (see Phytomyza).

Liriomyza strigata (see Agromyza). Lissonota parva, parasite of Coleophora

laricella in Ontario, 143. Listroderes obliquus, in U.S.A., 54, 261;

parasite imported against, 261; sprays

against, on tobacco, 54. Litchee (Litchi chinensis), pests of, in

Hawaii, 111, 250.

litchii, Eriophyes.

Lithocolletis felinelle, bionomics and control of, on *Platanus* spp. in U.S.A., **353.** Litomastix truncatella, parasite of Plusia

brassicae in Arizona, 112.

litura, Prodenia.

livornica, Celerio lineata.

Lixophaga diatraeae (parasite of Diatraea saccharalis), imported into U.S.A., 261; established in Florida, 260.

Lixophaga orbitalis, parasite of Cydia pomonella in U.S.A., 240; suggested utilisation of, in Canada, 241.

Lixus spp., on beet in Spain, 166.

Lizards, destroying noxious insects, 72, 356.

Lochmaea suturalis (Heather-beetle), in Britain, 232, 233; new fungus infesting, 232; bionomics and control of, 233.

Locust, Desert (see Schistocerca gregaria). Locust, Migratory (see Locusta migratoria migratorioides).

Locust, Moroccan (see Dociostaurus maroc-

canus).

Locusta marmorata, De Haan (see Dies-

trammena japanica).

Locusta marmorata, Harr. (see Scirtetica). Locusta migratoria migratorioides, Schistocerca gregaria possibly recorded as, in Belgian Congo, 366; in wheat-fields in Nigeria, 370; gammexane in against, 257.

locustarum, Eutrombidium (Trombidium)

(see E. trigonum).

Locusts and Grasshoppers, in E. Africa, 85, 87, 128; in W. Africa, 366, 370; in Arabia, 228; in Argentina, 66, 67, 120, 131, 252, 401; in Australia, 76, 321; in Bulgaria, 122; in Canada, 98, 144, 153, 154; in Ecuador, 5; in India, 136; in Iraq, 199-201; in S. Rhodesia, 126; in Spain, 166, 167; in U.S.A., 96, 101, 105, 174, 175, 193, 237, 259, 296, 341; phases of, 136, 166, 167, 201, 238, 388; breeding centres and 238, 388; breeding centres and ecology of, 136, 167, 200, 321; migrations of, 136, 200, 321, 366; effects of temperature, humidity and rainfall on, 76, 136, 167, 199, 322, 386, 388; diapause in eggs of, 76, 136, 167, 386; resistance of maize varieties to, 66, 67; Empusa grylli infesting, 122; insect enemies of, 201, 401; mite attacking, 193; protozoan parasites of, 296; mice feeding on eggs of, 96; measures and experiments against (see also under Baits), 87, 131, 201, 252, 257, 321; time schedules for surveys of, 341.

Loganberry, experiments with Thomasiniana theobaldi and, in Britain, 308.

Lolium, Metoponia rubriceps on, in New Zealand, 376.

Lonchocarpus spp., tests of toxicity of extracts of, to insects, 179, 197. (See Cubé.)

longana, Cnephasia.

longiceps, Aphaenogaster.

longicornis, Diabrotica; Paratrechina (Prenolepis).

longinoda, Oecophylla smaragdina.

longior, Tyrophagus (see T. putrescentiae). longipes, Anabrus.

longispinus, Pseudococcus (see P. adonidum).

longivalvata, Protopulvinaria.

longulus, Eulophus.

longus, Pachymerus.

lophanthae, Lindorus. lophyri, Lamachus.

Loquat (see Eriobotrya japonica). "Lorol" Thiocyanate, 44. (See Thiocyanate, anates.)

Louisiana, cotton pests in, 111, 261, 278; Ctenolepisma urbana in, 107; Cylas formicarius in, 216; Diatraea saccharalis in, 25, 26, 177, 260; Pantomorus in, 259; beneficial insects and biological control in, 26, 177, 216.

lounsburyi, Aspidiotiphagus.

Loxostege sticticalis, parasites America, 75.

loxostegei, Meteorus.

Loxotropa (parasite of Oscinella frit), attempted rearing of, on Drosophila, 75.

Lucerne, 92; Mocis repanda on, in Brazil, 93; Bruchophagus gibbus intercepted in seed of, in Brazil, 375; pests of, in Canada, 140, 153, 154, 339; pests of, in Czechoslovakia, 252; Aulacophora foveicollis on, in Greece, 17; Phytodecta fornicata on, in Rumania, 121; Laphygma exigua on, in S. Rhodesia, 126; pests of, in Russia, 305; pests of, in U.S.A., 22, 29, 40, 81, 100, 101, 106, 206, 241, 280, 285, 338, 339, 340; Capsids affecting seed production of, 29, 280; unfavourable to Limonius spp., 23; dinitro compound not injurious to,

Lucerne Meal, unsatisfactory carrier for cubé, 283.

Lucuma, fruit-flies infesting, in Mexico,

L'ucumaphila sagittata, bionomics of, in Mexico, 397.

ludens, Anastrepha.

luminosus, Pyrophorus.

lunulatus, Coccophagus (see C. scutellaris). Lupin, Metoponia rubriceps on, in New Zealand, 376.

luridula, Eurychilopterella.

luteipes, Azya.

luteola, Galerucella.

Lycidocoris mimeticus, on coffee in Fr. Togoland, 196.

lycopersicella, Keiferia.

Lycophotia margaritosa (see Peridroma

saucia).

Lyctus (in timber), in Australia and Britain, 181; in Germany, 18; bionomics and control of, in India, 71, 72; natural enemies of, 72, 181.

Lyctus parallelipipedus, in timber in

U.S.A., 176.

Lyctus planicollis, in timber in U.S.A., 176.

Lyda stellata, disease of, 7.

Lydella [stabulans var.] grisescens, relation of, to strains of Pyrausta nubilalis in Ontario, 139.

Lydinolydella metallica, imported into U.S.A. against Epilachna varivestis, 261.

Lygocerus, parasite of Clausenia purpurea

in Virginia, 109.

Lygus, in Czechoslovakia, 252; in Tanganyika, 88, 89; in U.S.A., 29, 101, 106; on beet, 101; on cotton, 88, 89, 101; on lucerne, 29, 101, 106, 252; effect of, on lucerne seed production, 29, 280; Collops predacious on, 106; measures against, 29, 252.

Lygus elisus, on lucerne in U.S.A., 101,

280

Lygus hesperus (in U.S.A.), on lucerne,

101, 280; dusts against, 63.

Lygus oblineatus, in Br. Columbia, 154; in U.S.A., 29, 63, 64, 101, 207, 280, 285, 316; on fruit trees, 154, 285, 316; on lucerne, 29, 101, 280; on other plants, 63, 207, 285; measures and experiments against, 29, 63, 64, 154, 207.

Lygus pabulinus, bionomics of, on strawberry in Britain, 360; on currant in Holland, 253; measures against, 253,

360.

Lygus sallei, on Parthenium argentatum in California, 105.

Lygus simonyi, factors affecting damage to cotton by, in Uganda, 88.

Lygus spinolae, sprays against, on vines in Austria, 117.

Lymantria dispar, in Czechoslovakia, 8; in Morocco, 310; bionomics of, in

in Portugal, 309, 310; in U.S.A., 4, 43, 53, 145, 238, 239, 310; on apple, 4; effects of defoliation of oaks by, 43, 309; tests and use of females and extracts of them for trapping males of, 8, 238, 239; insect enemies, disease and biological control of, 238, 310; quarantine fumigation against, 145; other measures against, 4, 53, 238, 239, 309.

Lymantria monacha, use of females for trapping males of, in spruce forests in Czechoslovakia, 8; Carabid predacious on, in Germany, 8; diseases of, 7, 8; pupal size and egg production in, 15.

Lysiphlebus, subgenus of Aphidius (q.v.),

Lysol, ineffective as repellent against *Melolontha*, **182**.

M.

Macaranga, Coccid on, in Br. Cameroons, 196.

Macremphytus varianus, on Cornus spp.

in U.S.A., 4.

Macrocentrus ancylivorus, in U.S.A., 24, 28, 47, 95, 105, 213, 218, 219, 240; introduced into Uruguay, 95; failure to establish, in Victoria, 322; use of, against Ancylis comptana, 262; use and value of, against Cydia molesta, 47, 95, 105, 213, 218, 219, 262, 322; attempted use of, against C. nigricana, 24; parasitising other Lepidoptera, 28, 47, 240; rearing of, on Gnorimoschema operculella, 218, 219; method of transporting, 219; toxicity of DDT to, 213.

Macrocentrus delicatus, hosts of, in New Jersey, 47.

Macrocentrus homonae, bionomics of, para-

sitising Homona coffearia in Ceylon, 300. Macrodactylus subspinosus, on blueberry in U.S.A., 402.

Macronaemia hauseri, parasite of, in Yunnan, 369.

Macrosiphum avenae (on cereals), natural enemies of, in Britain, 393, 394; in Nyasaland, 86.

Macrosiphum brachytarsus, on Cajanus cajan in Nyasaland, 86.

Tangan bhum facturas notus

Macrosiphum festucae, natural enemies of, on cereals in Britain, 393, 394.

Macrosiphum gei, auct. (see M. solanifolii).

Macrosiphum granarium (see M. avenae). Macrosiphum lilii, relation of, to virus diseases of ornamental plants in U.S.A., 1, 2, 38.

Macrosiphum onobrychis (in U.S.A.), on lucerne, 106, 206; on peas, 49, 100, 214,260,276,277,282,283; onvetch, 101; fungi infesting, 100; insects predacious

INDEX. 465

on, 106, 206; insecticides against, 49, 101, 214, 260, 276, 277, 282, 283.

Macrosiphum pisi (see M. onobrychis).

Macrosiphum rosae, new parasite of, in U.S.A., 96.

Macrosiphum solani, food-plants and populations of, in Britain, 358, 359; in U.S.A., 1, 2, 38, 40; and virus disease of potato, 2, 40, 358; not transmitting virus diseases of lilies, 1, 38; insecticides against, 2.

Macrosiphum solanifolii, in Australia, 34, 323; in Brazil, 288; in Britain, 358, 359; in Hawaii, 251; in U.S.A., 1, 2, 38, 40, 49, 96, 261; on potato, 2, 3, 40, 251, 288, 323, 358, 359; on other plants, 1, 38, 49, 251, 261, 358; relation of, to virus diseases, 1, 2, 38, 40, 323; new parasite of, 96; factors influencing populations of, 34, 358, 359; insecticides against, 2, 49, 288.

Macrosiphum sonchi, tests of mamey extract against, on Senecio confusus in

Porto Rico, 339.

Macrosteles divisus, food-plants and relation of, to virus disease in U.S.A., 71, 104, 173, 207, 378; forms of, 378; experiments with insecticides against, 104, 207.

macula-alba, Ceuthorrhynchus.

maculata, Empria.

maculatissima, Poecilophila (Porphyronota). maculatus, Dermestes; Neoplectrus.

maculipennis, Microplitis; Plutella.

maculiventris, Podisus.

Madras, Aphid transmitting mosaic of cardamom in, 392; food-plants and parasites of Pempherulus affinis in, 20-21; miscellaneous pests in, 246, 387,

Maerua, Anaphaeis aurota on, in Belgian

Congo, 368.

Maesa rufescens, Helopeltis orophila on, in Belgian Congo, 37.

Magdalis violacea, parasite of, in Finland,

Magnesite, against pests of stored products, 31, 187, 318, 319, 372; and sulphate, against cabbage nicotine pests, 322.

Magnesium Arsenate, spraying with, against Epilachna varivestis, 176.

Magnesium Oxide, experiments against pests of stored products, 228, 327; reducing toxicity of derris dust to Musca domestica, 78.

maidiradicis, Anuraphis. maidis, Aphis; Peregrinus.

Maine, outbreak and biological control of Gilpinia hercyniae in, 47; miscellaneous pests in, 3, 4, 354; orchard pests in, 3, 4, 42, 322; Aphids and virus disease of potato in, 2.

Maize, pests of, in E. Africa, 85, 86, 88, 89, 246; pests of, in Argentina, 66, 67, 242; pests of, in Brazil, 93; Pyrausta nubilalis on, in Canada, 137-139, 224; Aulacophora foveicollis on, in Greece, 17; Zabrus tenebrioides on, in Hungary, 11; pests of, in Mauritius, 371; Metoponia rubriceps on, in New Zealand, 376; Heteronychus sanctae-helenae on, in N.S. Wales, 161; increasing damage to cotton by Helopeltis in Nigeria, 383; pests of, in Pacific Is., 163, 250, 290; favouring natural enemies of Heliothis vivescens in Peru, 278; pests of, in S. Rhodesia, 126; pests of, in U.S.A., 3, 23, 26, 30, 49, 57, 101, 151, 177, 191, 237, 239, 259, 260, 337; treatment of ears of, against Heliothis armigera, 260, 337; varieties of, resistant to pests, 66, 67, 259; time of planting of, in relation to pests, 49, 89, 239; insecticide treatments not injurious to, 54, 56.

Maize (Stored), pests in, and their control, 25, 102, 131, 237, 312, 327, 361, 375; (cobs), ineffective as bait for cutworms, 140; (meal), in baits, 280, 370.

malacellus, Crambus.

Malachite Green, dust of, against Merodon equestris on Narcissus, 24.

Malachius, possibly predacious on Cydia

pomonella in France, 356. Malacocoris chlorizans, predacious on Paratetranychus pilosus in Britain, 234.

Malacosoma disstria, trees defoliated by, in New Brunswick, 155; in U.S.A., 43, 402; measures against, on blueberry, 402; rendering oak susceptible to Agrilus bilineatus, 43.

Malacosoma neustria, effect of diet on development of, in Bulgaria, 119.

Malaya, Eretmocerus serius obtained from, 261.

Aphelinus; Atractotomus; Leptomali, thrips; Psylla.

malivorella, Coleophora.

malus, Hemisarcoptes.

Mamestra brassicae, on Papaver somniferum in Hungary, 122; on crucifers in Russia, 304, 305; parasite of, 304, 305; tests of insecticides on, 197, 198, 304.

Mammea americana (Mamey), Anastrepha ludens on, in Mexico, 395; Myelois notabilis on, in Porto Rico, 339; insecticides obtained from, 338, 339.

Mandarin Orange (see Orange, Mandarin). Manganese Dioxide, reducing toxicity of derris dust, 78.

Mangifera indica (see Mango).

mangiferae, Aulacaspis cinnamomi (see A. cinnamomi); Cryptorrhynchus.

Mango, Coccids on, in Egypt, 227, 228; Toxoptera aurantii on, in Hawaii, 251; Anastrepha spp. on, in Mexico, 52,

53, 340, 395, 396; pests of, in New Caledonia, 161; Coccus hesperidum on, in Palestine, 326; A. mombin praeoptans on, in Porto Rico, 52, 397; Dacus ferrugineus tryoni on, in Queensland, 162; Ceratitis rosa on, in S. Rhodesia, 126; pests of, in Travancore, 387; (fruit), bacteria from, in bait for Anastrepha ludens, 396; sterilisation of, against fruit-flies, 396; (timber), limewash protecting, against Bostrychids,

mangoensis, Zizera labradus. Manihot utilissima (see Cassava). Manitoba, grasshoppers in, 144. Manninger's Apparatus, 252.

Mannitan Monolaurate, experiments with insecticides containing, 51.

mansueta, Campsomeris (Dielis).

Mantis religiosa, use of, against forest pests in Br. Columbia, 155.

Maple (Acer), Aphid on, in U.S.A., 353. Marasmia trapezalis, on sugar-cane in Porto Rico, 243.

Mareya spicata, Helopeltis bergrothi on, in Gold Coast, 196.

margaritosa, Lycophotia (see Peridroma saucia).

Margaronia indica, on pumpkin in Fiji, 290.

marginalis, Orthotylus. marginata, Mesogramma. marginatus, Eumerus. marginicollis, Cryptocephalus. marginiventris, Apanteles.

Marietta, hyperparasite of Saissetia nigra in Porto Rico, 374.

maritimus, Pseudococcus.

marmorata, De Haan, Diestrammena

(Locusta) (see D. japanica). marmorata, Harr., Scirtetica (Locusta).

maroccanus, Dociostaurus.

Marquesia, resistance of timber of, to termites in N. Rhodesia, 247.

marshalli, Chalcodermus.

Martius Yellow (see Dinitro-α-naphthol). Maruca testulalis, on beans in Fiji, 290; food-plants of, in Hawaii, 250.

marutae, Anuraphis (Aphis).

Maryland, Cotinis nitida in, 348; Popillia japonica in, 49, 50, 55, 347; pests of vegetables in, 49, 284.

Massachusetts, Hylemyia brassicae on crucifers in, 177; forest pests in, 43, 354; Lepismids in, 107; pests of stored sheep skins in, 82.

Mastrus aciculatus, parasite of Gilpinia

hercyniae in Quebec, 224.

Mastrus (Aenoplex) carpocapsae (parasite of Cydia pomonella), rearing of, in New Zealand, 269; method of rearing, on Ephestia in Ontario, 74; in U.S.A., 240, 241.

Mastrus pilifrons (Aenoplex betulaecola), parasite of Cydia spp. in U.S.A., 95, **240**; synonymy of, **240**.

materna, Othreis (Ophideres). mathesoni, Batrachedra.

mauritanicus, Tenebroides.

Mauritius, new Cecidomyiids on Citrus and Stenotaphrum in, 74; miscellaneous pests in, 371; sugar-cane pests and their biological control in, 370.

Mayetiola destructor, rearing and testing of, in relation to wheat varieties in

Indiana, 103.

maynei, Tragocephala. mayri, Telenomus. Meal Moth, Indian (see Plodia interpunctella).

Mealy Plum Aphis (see Hyalopterus arundinis).

Mealybug, Potato (see Phenacoccus solani). Mecistorhinus spp., bionomics of, cacao in Tinidad, 147.

Mecynotarsus, possibly predacious Cydia pomonella in France, 356.

mediata, Rhenea.

medicaginis, Aphis.

Medicago spp., Hypera variabilis on, in California, 41. (See Lucerne.)

Megaselia scalaris, in imported boot polish in Fiji, 161.

Melanagromyza (see Agromyza).

Melanaspis saccharicola, on sugar-cane and Pennisetum in Brazil, 205.

melanobatus, Erigorgus. melanocephala, Diabrotica. melanocephalus, Exochomus. melanochra, Stathmopoda.

melanoderes, Dysdercus. melanogaster, Drosophila.

Melanolophia spp., descriptions of larvae of, attacking forest trees in Canada, 180. melanopa, Lema.

Melanophila drummondi, on fire-killed Douglas fir in U.S.A., 40.

Melanophila fulvoguttata, drought favouring, on Tsuga in U.S.A., 43.

Melanoplus, review of data on important

species of, 174. Melanoplus bivittatus, in Saskatchewan,

153; in U.S.A., 174, 175; observations and survival of, on oviposition Iowa, 153, 175.

Melanoplus differentialis, in U.S.A., 96, 175; adult biology of, 175; absorption of water by egg of, 386; Peromyscus feeding on eggs of, 96; tests of arsenicals on, **278, 279.**

Melanoplus femur-rubrum, in Utah, 174. Melanoplus mexicanus, 216; outbreak of, in Br. Columbia, 154; in U.S.A., 174, 175; adult biology of, 175; egg-diapause in, **136.**

Melanoplus packardi, in Utah, 174.

melanothorax, Mesochorus.

melanotus, Coeloides.

Melia azedarach, Pseudococcus maritimus

on, in Egypt, 227.

Meligethes aeneus (Rape Beetle), in Britain, 257; apparatus for collecting, in Czechoslovakia, 252; in Denmark, 381; in Germany, 36; insecticides against, 36, 257, 381.

Melilotus (Sweet Clover), Lachnosterna on, in Canada, 140; pests of, in U.S.A., 29, 41, 96, 241, 285; use of, against

Limonius, 260; in bait for cutworms, 140. Melilotus hubamensis, Hypera variabilis reared on, 41.

Melissopus (see Cydia).

Melitara dentata (on Opuntia), bionomics and proposed utilisation of, in U.S.A., 241, 242.

Melittobia acasta, parasite of Cydia pomonella in France, 355.

Melittobia indica (see Syntomosphyrum).

mellerborgi, Polytus.

mellinipennis, Aegeria (Synanthedon).

mellipes, Euplectrus. mellitor, Bracon (Microbracon).

mellonella, Galleria.

Meloboris crassicornis, parasite of Oria musculosa in Britain, 308.

Melolontha, in Britain, 155; parasite of, in Denmark, 380; in Germany, 9, 10, 182; in Latvia, 7; in Poland, 9; in Russian Union, 159, 160; factors affecting abundance and distribution of species of, 9, 159; adult biology of, 10; diseases of, 7, 9; measures against, 7, 9, 10, 182.

Melolontha hippocastani, in Germany, 9, 10; in forests in Latvia, 7; distribution

of, in Russian Union, 159.

Melolontha melolontha (vulgaris); in Germany, 9, 10; in Latvia, 7; distribution of, in Russian Union, 159.

melolonthoides, Cochliotis.

Melon, pests of, in Greece, 16; pests of, in U.S.A., 21, 60, 179.

Melon Aphis (see Aphis gossypii).

Melophagus ovinus, mode of absorption of derris dust by, 383.

mendicus, Cleonus (Conorrhynchus). merceri, Contarinia.

Mercuric Chloride, solutions of, against *Hylemyia brassicae*, **178.**

Mercurous Chloride, uses of, against Hylemyia brassicae, 178, 276.

Mercury, experiments with vapour of, against pests of stored products, 164, 165, 363; solution of organic compound of, against *Hylemyia brassicae*, 178.

meridionalis, Ocnogyna boetica. meriopterus, Capsus (Heterotoma).

Merodon equestris, measures against, in Narcissus bulbs in Washington, 24.

Mescinia peruella, on cotton in Peru, 132. mesentina, Pieris (see Anaphaeis aurota).

Mesochorus, parasite of Plusia chalcites in Fiji, 290.

Mesochorus melanothorax, parasite of Euphorus sahlbergellae in W. Africa, 195. Mesogramma marginata, predacious on

Cinara in Bermuda, 290.

Mesohomotoma tessmanni, bionomics and relation of, to virus disease of cacao in Gold Coast, 66.

mesonotalis, Acropyga (Rhizomyrma).

Mesopeltis, introduced parasite of Saissetia oleae in California, 374; attempted use of, against S. nigra, 374.

Mesotrichia senior, bionomics of, on cassava in Belgian Congo, 367.

Metadinitrobenzene, as a contact insecticide, 198.

Metagonistylum minense (parasite of Diatraea saccharalis), in 'Florida, 260; attempted establishment of strains of, in Porto Rico, 26, 312.

Metal Shields, against termites, 73.

Metaleurodicus cardini (see Aleurodicus).

metallica, Lydinolydella.

Metanitraniline, 198.

Metapelma spectabilis, parasite of Cylas formicarius in Louisiana, 216.

Metaphycus flavus, parasite of Coccus hesperidum in Palestine, 326.

Metaphycus helvolus (parasite of Saissetia spp.), in California, 259, 374; introduced into N. S. Wales, 322; effect of insecticide treatments on, 259.

Metaphycus stanleyi (parasite of Saissetia spp.), introduced into California, 374; in Kenya, 374.

Metarbela sticticosta, on oak in Transvaal, 83.

Meteorus (parasite of Cydia pomonella), in France, 355; suggested introduction of, into Canada, 356.

Meteorus chrysophthalmus, 355.

Meteorus loxostegei, attempted rearing of, on hosts other than Loxostege, 75.

Methallyl Bromide and Chloride, fumigation with, against Calandragranaria, 229.

Methyl Alcohol, as solvent for benzene hexachloride, 257.

Methyl Bromide, fumigation of plants and planting material with, 24, 50, 64, 110, 111, 145; effects of, on plants, 24, 50, 64; against pests of fruit and vegetables, 111, 127, 214; against pests of other stored products, 127, 259, 343; not affecting vitamin B₁ in flour, 259; sorption of, by wheat, 128; reducing viability of wheat, 237, 259; determination of, 128, 368; fumigant mixture containing, 237.

Methyl Chloride, use of, for producing

aerosols, 43, 99, 261.

Methylated Spirit, in spray against Bagrada hilaris, 293.

4 - Methylcyclohexanone Semicarbazone, 99; toxicity of, to insects and plants, 279.

meticulosalis, Terastia.

468

Metoponia rubriceps, damage to plants by, in Australia and New Zealand, 269, 375. Mexican Bean Beetle (see Epilachna varivestis).

Mexican Fruit-fly (see Anastrepha ludens).

mexicanus, Melanoplus.

Mexico, pests of Agave in, 131; fruit-flies in, 52, 190, 216, 340, 395-397; miscellaneous pests in, 97, 147, 339, 352; beneficial insects and biological control in, 176, 251, 261, 395; introduction of beneficial insects into other countries from, 177, 251; modifications of regulations against importation of pests into U.S.A. from, 111, 286.

Lachnosterna Dendroctonus;

(Phyllophaga).

Mice, White-footed (see *Peromyscus*). Michigan, pests of blueberry in, 223, 401; orchard pests in, 314; Aphrophora spp. on pines in, 107.

Microbracon (see Bracon).

Microctonus epitricis, parasite of Epitrix hirtipennis in U.S.A., 41.

Microdus (see Agathis).

Microgaster kuchingensis, parasite of Pyrausta coclesalis in India, 389.

microgrammana, Cydia.

Micromyzus (see Idiopterus).

Microphanurus basalis, bionomics and utilisation of, against Nezara viridula in Pacific Is., 291.

Microplectron fuscipenne (parasite of Gilpinia hercyniae), liberation and establishment of, in Canada, 155, 224; factors affecting parasitism by, in Maine, 47; strain of, 155.

Microplitis brassicae, parasite of Plusia

brassicae in Arizona, 112.

Microplitis maculipennis (ophiusae), parasite of Achaea janata in India, 386.

Microterys consobrinus (see Encyrtus).

Microterys flavus, parasite of Saissetia nigra in Hawaii, 374.

Microterys frontatus (see Encyrtus).

Microterys kotinskyi, parasite of Saissetia nigra in Hawaii, 374.

Mictis profana, on orange in Fiji, 290. middletoni, Aphis.

Middlings (see Bran).

migratorioides, Locusta migratoria.

miliaris, Capnodis.

Milk (Dried), Trogoderma versicolor in, in California, 102; cheese mites reared on, 77; use of, in sprays, 148, 149, 334.

Milky Disease, Type A (see Bacillus popilliae).

Millettia, Coccid on, in Belgian Congo, 365.

mimeticus, Lycidocoris.

mindanaona, Lachnosterna (Ancylonycha).

minense, Metagonistylum.

Mineola vaccinii, on cranberry and blueberry in U.S.A., 23, 223, 401; insecticides against, 23, 223.

minerva, Draeculacephala.

Minnesota, Anabrus simplex in, 114; forest pests in, 107, 271; Capsids on lucerne and Melilotus in, 29; potato pests in, 207.

minuta, Frankliniella; Hoplocampa.

minutum, Trichogramma.

minutus, Dinoderus; Laemophloeus; Tetrastichus.

Mirax insularis, value and utilisation of, against Leucoptera coffeella in W. Indies,

miridicidus, Euphorus.

misera, Sarcophaga.

Mississippi, cotton pests in, 80, 260; miscellaneous pests in, 25, 69, 259.

Missouri, miscellaneous pests in, 107, 219, 258, 259.

Mites, attacking insects, 6, 17, 19, 164, 193, 328, 352, 356, 374; attacking other mites, 208, 397, 398; natural enemies of, 86, 135, 170, 208, 233, 361, 389, 397, 398; study of unidentified species of, damaging tomato, etc., in Ceylon, 299; survey of data on Tyroglyphid, in stored products, 368; effect of, on forest litter, 6; classification and new species of 111, 102 species of, 111, 193.

mjöbergi, Pheidole (Pheidolacanthinus). Mocis repanda, measures against, on cereals, grasses and lucerne in Brazil, 93; on sugar-cane in Porto Rico, 243.

modestus, Coccophagus.

Molasses, in baits, 27, 68, 134, 141, 201, 270; in bait-sprays for Trypetids, 134, 396, 397; in sprays against other insects, 87, 97.

molesta, Cydia (Grapholitha). molitor, Tenebrio.

Mombin, Yellow (see Spondias).

mombin praeoptans, Anastrepha.

Momordica balsamina, Ceratitis capitata on, in Hawaii, 250.

monacha, Corythaica; Lymantria.

Monieziella entomophaga (see phagus).

Monochaeta albicans, bionomics of, parasitising Operophtera brumata in Germany, 120.

Monodontomerus dentipes, utilisation of, against Neodiprion abietis in Br. Columbia, 155.

Monoethanolamine Oleate, as an emulsifier for oils, 108.

Monomorium carbonarium ebeninum, predacious on Diatraea saccharalis in Porto Rico, 243.

Monomorium rothsteini var. leda, measures against, collecting seeds of tobacco, etc., in Queensland, 369, 370.

Monotes, resistance of timber of, to

termites in N. Rhodesia, 247.

Montana, Cecidomyiid on clover in, 101; strain of Harmologa fumiferana on Pinus contorta in, 314; Tettigoniids in, 59, 114; parasite of Loxostege sticticalis in, 75; Epitrix tuberis not occurring in,

montrouzieri, Cryptolaemus.

Montserrat, Nezara viridula on Crotalaria in, 291.

montserratensis, Icerya. monuste, Ascia (Pieris).

mori, Bombyx.

Mormidea poecila (see Solubea).

Mormidea ypsilon, Solubea poecila recorded as, in Dutch Guiana, 147.

Mormon Cricket (see Anabrus simplex). Moroccan Locust (see Dociostaurus maroc-

canus).

Morocco, miscellaneous pests in, 170, 310, 356; beneficial insects and biological control in, 170, 310; Ovencyrtus kuvanae imported into Portugal from, 310.

mosellana, Sitodiplosis.

Mosquito Larvae, tests of plant extracts on, 65.

moultoni, Frankliniella.

Mountain Ash (see Eucalyptus regnans and Sorbus aucuparia).

Mucor hiemalis, infesting Kalotermes approximatus in Bermuda, 290.

Mucuna aterrima (see Stizolobium).

Mulberry, Eulecanium persicae on, in Egypt, 228.

multistriatus, Scolytus. munda, Cycloneda.

Murgantia histrionica, tests of dusts against, in New Jersey, 63.

muriceus, Phyrdenus.

musae, Dacus (Strumeta); Stenommatus. Musca domestica, experiments with parasites of other Diptera and, 75, 305; tests of insecticides, etc., on, 78, 137, 189, 245, 296, 331.

Muscina stabulans, 120.

musculosa, Oria.

musculus, Anthonomus.

museorum, Anthrenus.

Mushrooms, Sciara on, in Quebec, 142. Mustard, Colaphellus sophiae on, in Denmark, 381; relations of wireworms to, 260, 294.

Mustard Beetle (see Phaedon cochleariae). Mustard Oil (see Allyl Isothiocyanate). mutabilis, Ootheca.

mutica, Glypta.

Mvule (see Chlorophora excelsa).

Myelois ceratoniae, damage by, ascribed to Ephestia elutella, 232.

Myelois notabilis, on Mammea americana in Porto Rico, 339.

Myelophilus piniperda, parasites of, in Finland, 11, 12.

Mylabris dicincta, on leguminous crops in Nyasaland, 86.

Myllocerus laetivirens, bionomics and control of, in India, 388.

Mynah Bird (see Acridotheres tristis).

Myodochus serripes, on strawberry in Ohio, **263.**

myops, Oberea.

Myrciaria jaboticaba, Targionia jaboticabae on, in Brazil, 205.

myrmeleon, Eulophonotus.

Myrtus communis, Polychrosis botrana on, in Italy, **118.**

Mysore, Icerya purchasi on Acacia in, 391; Aphid transmitting mosaic of cardamom in, **392.**

Mytilococcus (see Lepidosaphes).

Myzus cerasi, sprays against, on cherry in

Br. Columbia, 153.

Myzus circumflexus, on papaya and rose in Hawaii, 251; relation of, to virus diseases of ornamental plants in U.S.A., 1, 2, 38.

convolvulii (see Macrosiphum Myzus solani).

Myzus festucae (see Macrosiphum).

Myzus kaltenbachi, parasite of, on oats in Britain, 77.

Myzus persicae, in Australia, 34, 323, 378; in Britain, 358; in Hawaii, 251; in New Zealand, 247; in Nyasaland, 85; in Porto Rico, 339; in S. Rhodesia, 126; in U.S.A., 1, 2, 38, 40, 44, 51, 70, 71, 96, 174, 206, 261; on beet, 206, 251; on cotton, 261; on crucifers, 2, 3, 39, 126, 251, 323, 339, 358, 359; on peach, 323; on potato, 34, 39, 126, 323, 358, 359; transmitting potato virus diseases, 2, 40, 323, 378; transmitting virus diseases of ornamental plants, 1, 2, 38, 71, 174; on other plants, 51, 85, 126, 251; relation of, to virus diseases of other plants, 70, 160, 247; bionomics and factors influencing populations of, 34, 323, 358; natural enemies of, 96, 206; measures and experiments against, 3, 44, 51, 339.

Myzus pseudosolani (see Macrosiphum solani).

N.

Naaki, 231, 372.

Nabis alternatus, Collops predacious on, in Utah, 106.

Nabis punctipennis, predacious on Heliothis virescens in Peru, 278.

Nadiasa concolor, on Pinus radiata in S. Africa, 83.

Nadiasa (Taragama) nyassana, on Aleurites and Bridelia in Nyasaland, 86.

Nadiasa (Taragama) repanda, bionomics of, in Spain, 170.

Nadiasa (Taragama) siva, food-plants of, in India, 393.

naevana, Rhopobota.

nana, Angitia; Lasiohelea.

nanulus, Neodiprion.

Naptha, against ants, 322; in mixtures against Coleoptera in timber, 84; uses of, against Tineola biselliella, 320, 321; as a solvent, 84, 321; forms of, 84, 320, 322.

Naphthalene, against Hylemyia brassicae, 178; soil treatment with, against Limonius spp., 23, 260; as repellent against Melolontha, 182; in sprays against Rhoecocoris sulciventris, 268; uses and value of, against Anthrenus and clothes moths, 72, 320, 321, 331; against pests in stored bulbs and potatoes, 35, 404; solution of, against termites in timber. 244.

Naphthenic Acid, in sprays, 273. Napomyza lateralis (see Phytomyza).

nararia, Natada.

Narcissus, treatment of bulbs of, against pests, 24; possible host of tulip virus disease, 39.

Narosa conspersa, on tea in Ceylon, 300. Natada nararia, parasites and disease of, on tea in Ceylon, 300.

natadae, Platyplectrus.

Nebraska, Anabrus simplex in, 114; Cydia molesta in, 258; pests of potato in, 216, 220; pests of stored grain in, 25. necopinus, Collops.

Necrobia rufipes, in sheepskins in Massa-

chusetts, 82, 83.

Nectria coccinea, associated with Cryptococcus fagi on beech in Canada, 354.

Nectria tuberculariae, use of conidial form of, infesting Lepidosaphes beckii in Belgian Congo, 364.

neglectus, Dysdercus (see D. sanguinarius). Nematodes, 362; infesting Lamellicorn

larvae, **61.**

Nematospora gossypii, method of spread

of, by Dysdercus, 357.

Nematus ribesii (on gooseberry and currant), bionomics of, in Britain, 360; in Denmark, 381; bird destroying, in Germany, 14; in Ontario, 64; insecticides against, 360, 381.

nemorum, Anthocoris; Phyllotreta.

nenuphar, Conotrachelus.

Neoaplectana spp., infesting Popillia japonica in U.S.A., 61; failure to infest Lachnosterna with, 61.

Neochmosis (see Cinara).

Neoclytus, in timber in U.S.A., 176.

Neodiprion abietis, liberation of Monodontomerus dentipes against, on conifers in Br. Columbia, 155.

Neodiprion dubiosus (see N. rugifrons). Neodiprion lecontei, on Pinus spp. in

Pennsylvania, 4.

Neodiprion nanulus, natural factors affecting, on Pinus banksiana in Canada, 8, 9.

Neodiprion rugifrons, natural affecting, on Pinus banksiana in

Canada, 8, 9.

Neodiprion sertifer, disease of, in Europe, Neodiprion swainei, natural factors affec-

ing, on Pinus banksiana in Canada,

8, 9. Neokolla spp., transmitting virus disease of vines in U.S.A., 172.

Neoplectrus maculatus, parasite of Limacodid in Ceylon, 300.

Neosyl, 66, 372.

Nerium oleander, Asterolecanium pustulans on, in Egypt, 227.

nero, Panoquina.

Netherlands Indies, Pyrausta coclesalis in, 388; Stenommatus musae in, 202.

neustria, Malacosoma.

Nevada, Anabrus simplex and its natural enemies in, 59, 115, 146, 368; foodplants of Eutettix tenellus in, 281.

New Brunswick, forest pests and their biological control in, 143, 154, 155, 398,

399.

New Caledonia, miscellaneous pests in, 161, 165, 290; parasite of Nezara imported into, 291; risk of introduction of pests into Hawaii from, 290.

New Hampshire, forest pests in, 354, 398. New Hebrides, Platyedra gossypiella in,

165.

New Jersey, Scolytus multistriatus on elm in, 313; Pyrausta nubilalis on maize in, 239; orchard pests in, 46, 105, 208, 213, 222, 258; Ancylis comptana on strawberry in, 47; vegetable pests in, 147, 221; Popillia japonica in, 63; beneficial insects and biological control in, 46, 105, 208, 213, 222; beneficial insects introduced into Washington from, 21,24.

New Mexico, Cydia pomonella on apple in, 68; biological control of Hemileuca oliviae on grasses in, 215; miscellaneous

pests in, 63, 114, 314, 353. New South Wales, Aphid and virus disease of carrot in, 369; Chortoicetes terminifera in, 321; pests of Citrus in, 248, 267, 268, 322; Gnorimoschema operculella on field and stored potatoes in, 184-187, 323; injurious Lamellicorns in, 161, 162; Tineola biselliella in baled wool in, 320; natural enemies and INDEX.

biological control of insects in, 248, 267, 268, 322; biological control of Hypericum perforatum in, 321; bean Bruchid imported into Fiji from, 290.

New York, forest pests in, 180, 313, 354; hop pests in, 317; miscellaneous pests in, 44, 179, 180; orchard pests in, 42, 58, 148, 149, 150, 179, 274; pests of vegetables in, 52, 179, 180, 206; parasite of Tineola biselliella in, 352.

New Zealand, Coleophora spissicornis on clover in, 163, 184; Myzus persicae transmitting virus disease of Hyoscyamus in, 247; miscellaneous pests in, 183, 269, 369, 375; pests of vegetables in, 93, 247, 269, 376; termites in, 234; mites in cheese and flax seed in, 77, 130, 269; parasites and biological control of insects in, 247, 269, 376; biological control of noxious plants in, 269, 321; insects imported into Fiji from, 161, 291.

Newfoundland, survey of forest pests in,

354.

Nezara viridula, food-plants of, in Fiji, 290, 291; in New Caledonia, 291; DDT against, in N. S. Wales, 322; parasites and biological control of, 291, 322.

Nezara viridula var. smaragdula, 290, 291. Nicandra physaloides, Lema trilineata californica on, in Hawaii, 250; Gnorimoschema operculella on, in N. S. Wales, 185; Myzus persicae on, in S. Rhodesia,

Nicobium castaneum var. hirtum, in timber in U.S.A., 176.

Nicotiana rustica, 85.

Nicotiana tabacum (see Tobacco).

Nicotine, 99; against Aphids, 36, 50, 54, 103, 180, 260, 293, 331, 336, 337, 338; against Capsids, 38, 66, 360, 361; against Cocids, 171, 374; against other Hemiptera, 21, 22, 264; against Cole-optera, 17, 168, 304, 315; against Contarinia pyrivora, 12; against Cydia pomonella, 43, 149, 150, 179, 273, 274, 359; against other Lepidoptera, 23, 28, 49, 139, 179, 223, 239, 260, 333, 338, 353, 360; against Paratetranychus pilosus, 124; against sawflies, 359, 360, 380; against thrips, 338, 364; dusting with, 21, 22, 43, 49, 50, 66, 103, 150, 171, 179, 239, 260, 274, 293, 304, 316, 333, 336, 337, 338, 360, 361; carriers for, 49, 239, 274; abrasive dust accelinsecticidal action of, 189; erating applied by aircraft, 338; application of, by vaporisation, 180, 360; fixed compounds of, 23, 28, 43, 49, 139, 149, 150, 179, 223, 239, 273, 274, 315; formulae containing, 23, 28, 36, 38, 43, 49, 50, 139, 149, 150, 171, 179, 239, 260, 273,

274, 293, 316, 336, 337, 338, 359, 360, 361, 364; and calcium arsenate, 337; and cryolite, 337, 338, 380; and dichlorethyl ether, 338; and lead arsenate, 43, 149, 150, 274, 359; and lime-sulphur, 359; and oil emulsions, 23, 149, 171, 179, 223, 273, 274; oil in dusts containing, 150; and pyrethrum, 260; and rotenone insecticides, 50, 66, 239, 260, 336, 338; and sulphur, 43, 49, 150, 239, 260, 274, 316; and thiocyanate, 149; and soap, 38, 124, 168, 264, 293, 380; addition of other agents to sprays containing, 149, 273, 274; nornicotines compared with, 54. Nicotine Cuprocyanide, experiments with,

47I

against Cydia pomonella, 179.

Nicotine Sulphate, against Aphids, 2, 50, 60, 153, 172, 180, 293, 317, 318, 322, 336; not harmful to natural enemies of Aphids, 60; against other Hemiptera, 28, 66, 142, 168; against Cydia pomonella, 149, 150, 179, 272-274; ineffective as repellent against mature larvae of C. pomonella, 274; soil treatment with mixture containing, against cocoon stages of *C. pomonella*, **108**; against other Lepidoptera, **23**, **28**, **179**, **185**, **210**, **239**, **262**, **322**, **339**; in bait-sprays for fruit-flies, 396; against sawflies, 263, 334; against thrips, 97, 284, 285; methods of testing, on Drosophila melanogaster, 188; dusting with, 60, 317, 322, 336; formulae containing, 23, 28, 60, 366, 97, 108, 142, 149, 150, 153, 172, 185, 210, 239, 262, 273, 274, 284, 293, 317, 322, 334, 336, 339, 396; and bentonite, 28, 149, 239, 272–274; and copper compounds, 2, 180, 317; and derris, 2; and lead arsenate, 149, 150, 210. 334: and lime, 60, 317; lime-sulphur, 285; and magnesite. 322; and oil emulsion, 23, 28, 108, 142, 149, 153, 179, 185, 210, 262, 268, 273, 284, 334; and sulphur, 150; and soap, 172, 185, 268, 293, 317; addition of other agents to sprays containing, 97, 149, 210, 273, 339.

niger, Chalcodermus; Lasius; Thysanus. Nigeria, cacao pests and associated insects in, **195**, **196**; Helopeltis on cotton in, 91, 383; Bruchid in field and stored nuts of Elaeis in, 165, 370;

miscellaneous pests in, 196, 370.

nigra, Saissetia.

nigrescens, Dysdercus.

nigricana, Cydia.

nigricans, Euxoa (Agrotis); Pityophthorus.

nigriceps, Cardiochiles. nigriteleus, Aphidius.

nigritus, Chilocorus. nigrofasciatum, Eulecanium (Lecanium). nigrofasciatus, Dysdercus. nigromaculatus, Exochomus (see E. melanocephalus.)

nigronervosa, Pentalonia.

Niphadolepis alianta, on tea in Nyasaland, 85.

nitidalis, Diaphania. nitidula, Cladis.

Nitriles, 44, 59, 104, 106, 128, 198, 295. Nitromethane, as a solvent, 189, 190. NNOP, NNOR and NNORP, 51.

nobbei, Eurytoma.

noctilio, Sirex.

Nomadacris septemfasciata, swarms and control of, in E. Africa, 85, 87; in S. Rhodesia, 126.

nordmannianae, Chermes.

Nornicotine, high toxicity of, to Aphids, 54; content of, in *Duboisia hopwoodii*,

54; identification of, in tobacco, 368.

notabilis, Myelois.

notativentris, Acerophagus (Pseudaphycus). Nothofagus cunninghami, susceptibility of, to borers in Victoria, 378.

Nothopanax guilfoylei, Aphis rumicis on, in Hawaii, 251.

Notodacus (see Dacus).

Nova Scotia, Rhagoletis pomonella on blueberry in, 144; Popillia japonica in, 137

Novius cardinalis (see Rodolia).

noxius, Anodontonyx. nubilalis, Pyrausta. nubilipennis, Hemadas. nubilus, Helophorus.

nuonus, Helophorus. nucum, Curculio.

Nudaurelia cytherea capensis, on pines in S. Africa, 83.

Nun Moth (see Lymantria monacha).

nüsslini, Chermes (Dreyfusia) (see C.

nordmannianae).

Nyasaland, cotton pests in, 85, 86, 89–91; miscellaneous pests in, 85–87; beneficial insects in, 85, 86; restrictions on importation of dahlias into, 85.

nyassana, Nadiasa (Taragama).

nyctelius, Panoquina.

Nygmia phaeorrhoea, in Britain, 360; in Bulgaria, 119; in Czechoslovakia, 8; in Portugal, 308; quarantine fumigation against, in U.S.A., 145; bionomics of, on fruit trees and oak, 119, 308, 360; use of females to attract males of, 8. nymphaeae, Rhopalosiphum.

Nysius ericae, outbreak of, in Br. Columbia, 154; derris ineffective against

adults of, 154.

O.

Oak, Metarbela sticticosta on, in S. Africa, 83; Boarmia roboraria on, in Britain, 359; Lepidoptera on, in Bulgaria, 119;

Icerya purchasi on, in France, 357; Dasychira pudibunda on, in Sweden, 19; pests of, in U.S.A., 43, 238, 270; Cnephasia longana ovipositing on, 342. Oak, Cork (see Quercus ilex var. suber).

Oats, 101; Toxoptera graminum on, in Argentina, 310; pests of, in Britain, 77, 199, 265, 294, 297, 394; Zabrus tenebrioides on, in Hungary, 11; Anodontonyx noxius on, in N. S. Wales, 161; Delphacodes striatellus and virus disease of, in Russian Union, 183; not attracted by Phyllobius pyri, 92; dinitro compound not injurious to, 54.

Oberea myops, bionomics and control of,

on blueberry in U.S.A., 402.

oblineatus, Lygus. obliqua, Allograpta. obliquus, Listroderes.

obliterata, Angitia (Inareolata); Strangalia (Leptura).

oblongus, Phyllobius. obscura, Schistocerca. obscurus, Agriotes.

obsoleta, Chloridea (see Heliothis armigera). obtectus, Bruchus (Acanthoscelides).

obtusus, Tachyporus.

occidentalis, Cuerna; Frankliniella.

ocellana, Spilonota (Tmetocera).

ocellatella, Gnorimoschema (Phthorimaea). ochraceus, Coccophagus.

Ochroma lagopus, Ferrisia virgata on, in Br. Cameroons, 196.

Ocnogyna boetica var. meridionalis, natural enemies of, in Spain and Morocco, 170. Octadecadienyl Nitrile, mixture con-

taining, not toxic to insects, 198.

Octadecyl Nitrile, mixture containing, not toxic to insects, 198. oculiperda, Thomasiniana.

Odoniella rubra, records of, in W. Africa, 196.

Odontionopa sericea (see Prasoidea).

Oecanthus brevicauda, on cotton and tobacco in Nyasaland, 86.

Oecophylla smaragdina, associated with Coccid on cacao in Belgian Congo, 366.

Oecophylla smaragdina longinoda, attempted utilisation of, against Capsids in W. Africa, 65, 195.

Ohio, orchard pests in, 191, 208, 258, 314; Pyrausta nubilalis in, 151; Anisolabis annulipes on greenhouse radish in, 67; strawberry pests in, 262, 263; natural enemies and biological control of pests in, 208, 209, 258, 262.

Oiketicus kirbyi, Apion simplex in nests of, in Argentina, 132.

Oil Emulsions, against Aleurodids, 51, 172; against Aphids, 51, 153, 253; against Capsids, 38, 66, 153, 253, 254; against Coccids, 22, 58, 98, 113, 133, 171, 172, 204, 211, 227, 228, 248,

INDEX. 473

249, 253, 258, 259, 268, 284, 285, 342, 346, 349, 350, 353, 364, 374; against other Hemiptera, 142, 253, 268, 285, 353; against ants, 371; against Coleoptera, 38, 168, 212, 319; against Cydia pomonella, 108, 148, 149, 179, **274, 284, 316;** not repellent to mature larvae of C. pomonella, 274; in mixtures for soil treatment against cocoon stages of C. pomonella, 108, 109; against other Lepidoptera, 23, 28, 51, 94, 97, 179, 185, 210, 212, 223, 254, 262; treatment of potato tubers with, against Gnorimoschema operculella, 186, 187; against mites, 124, 148, 172, 208, 233, 234, 253, 254, 285, 286, 404; against sawflies, 263, 334; soil treatment with, against termites, 74; against thrips, 285; destroying Metaphycus helvolus, 259; tests of toxicity of, to insect eggs, 15, 254; types of oils for, 23, 24, 58, 66, 97, 108, 113, 124, 142, 148, 149, 171, 210, 211, 223, 227, 233, 234, 254, 258, 262, 268, 273, 274, 284, 234, 254, 258, 262, 268, 273, 274, 284, 314, 334, 342, 346, 349, 350, 371, 374; emulsifiers for, other than soap, 51, 58, 108, 148, 149, 211, 334, 371; formulae for, 22, 23, 58, 94, 97, 108, 109, 113, 124, 133, 142, 148, 153, 171, 172, 185, 186, 187, 210, 211, 212, 227, 234, 249, 258, 262, 268, 284, 285, 334, 271, 374; formulae containing, 23, 24 371, 374; formulae containing, 23, 24, 28, 38, 94, 108, 109, 113, 142, 149, 153, 171, 185, 210, 211, 212, 227, 262, 273, 274, 284, 285, 216, 234, 249, 350; and 274, 284, 285, 316, 334, 349, 350; and aluminium sulphate, 273; and Bordeaux mixture, 58, 148, 268, 364; and calcium chloride, 404; and cryolite, 210, 284; and DDT, 113; and dichlorethyl ether, 108; and dinitro compounds, 15, 109, 153, 179, 233, 253, 258, 319; and Fermate, 148; and lead arsenate, 24, 28, 94, 185, 210, 212, 262, 284, 316, 334; "inverted" (dynamite) mixtures of lead arsenate and, 150, 151; and lime-sulphur, 22, 285; and nicotine, 23, 28, 108, 142, 149, 153, 171, 179, 185, 210, 223, 262, 268, 273, 274, 285, 334; and ortho- or paradichlorbenzene, 108, 227; and phenothiazine, 149; and potassium cyanide, 268; and pyrethrum, 38, 51, 108; and rotenone insecticides, 51, 113, 211, 212, 234, 258, 342, 346, 349; mutual solvents for derris and cubé extractives and, 349, 350; and tar distillates, 66; and thiocyanates, 149, 234; and zinc sulphate, 210, 364; materials reducing penetration of, into plant tissue, 253; and injury to plants, 66, 97, 105, 172, 227, 234, 334, 404.

Oil Palm (see Elaeis guineënsis).

Oils, as barriers against Blissus leucopterus,

57, 58; in treatments against Coleoptera in timber, 72, 84, 371; treatment of sheds with, against Depressaria heracleana, 333; apparatus for burning, to destroy Epilachna varivestis, 139; and insecticides, treatment of maize ears with, against Heliothis armigera, 260, 337; soil treatment with, against termites, 74; uses of, against Tineola biselliella, 320; double-nozzled spray apparatus for applying water and, 379; atomised sprays of insecticide in, 31, 38, 127, 254; in baits, 55, 140, 201, 259, 348; dusts impregnated with, 24, 150, 152, 158, 159, 276, 277, 283, 316, 317. (See Kerosene.)

Oils, Essential, in baits for *Polychrosis* viteana, 27; for treating wood against termites, 244; proprietary insecticide

containing, 51.

Oils, Vegetable, in sprays, **54**, **364**; uses of sulphonated, in sprays, **51**, **257**. (See Cottonseed, Sesame and Soy-bean

Oils.)

Oklahoma, pests of field and stored cereals in, 69, 70, 151, 237; Cydia molesta in, 258; Aphis gossypii and its natural enemies on melon in, 60; injurious Orthoptera in, 237, 270; Bufo spp. destroying cutworms in, 96.

oleae, Dacus; Parlatoria; Saissetia.

Oleander (see Nerium oleander).

Oleic Acid, uses of, in sprays, 273, 274, 364. oleivorus, Phyllocoptruta (Eriophyes, Phyllocoptes).

Olethreute's (see Argyroploce).

Oligonychus ulmi, auct. (see Parate-

tranychus pilosus).

Olive, Parlatoria oleae on, in California, 107; pests of, in Italy, 118; Teleonemia australis on, in S. Rhodesia, 126; injured by bait-sprays, 126.

Olive Fly (see Dacus oleae).

Olive Scale (see Parlatoria oleae).

oliviae, Hemileuca.

Olla abdominalis, predacious on Saissetia nigra in California, 374.

omnivorus, Psychophagus.

Omocena convergens, bionomics of, in Nyasaland, 86.

Omorgus tortricidis, parasite of Polychrosis viteana in U.S.A., 28.

Oncopeltus fasciatus, tests of fumigants on, 275; experiments with infra-red rays and, 53.

Oncopera, attempted introduction of Alomya debellator against, in Australia,

322.

Onion, Petrobia lapidum on, in Britain, 361; Hylemyia antiqua on, in Br. Columbia, 154; pests of, in Hawaii, 250, 251; Thrips tabaci on, in Tanganyika, 85; pests of, in U.S.A., 52, 70, 71,

180, 338; insects and virus diseases of, 70, 71; absorption of arsenic by, from soil, 63.

Onion Fly (see Hylemyia antiqua).

Onion Thrips (see Thrips tabaci).

onobrychis, Macrosiphum.

ononis, Heliothis.

Ontario, Epilachna varivestis on beans in, 144; pests of currants and goosberries in, 64: forest pests in, 8, 71, 142, 143, 313,314,354; miscellaneous pests in, 140, 141, 146, 354; Popillia japonica in, 137, 224; Pyrausta nubilalis in, 137, 138, 139; beneficial insects and biological control in, 139, 143, 181, 354.

Onychomys, destroying Chlorion laeviven-

tris in U.S.A., 146.

Ocencyrtus kuvanae, establishment of, against Lymantria dispar in various countries, 310.

Ootheca mutabilis, on leguminous crops in Nyasaland, 86.

opacus, Hylastes.

Gnorimoschema (Phthorioperculella, maea).

Operophtera brumata, in Germany, 35, 119; on apple in Switzerland, 394; bionomics of, 119; measures experiments against, 35, 253, 257.

Ophideres ancilla, fullonia and materna

(see Othreis).

Ophideres salaminia (see Eumaenas).

Ophiomyia (see Agromyza).

ophiusae, Microplitis (see M. maculipennis).

Ophonus pubescens (see Harpalus rufipes). Opilo domesticus, predacious on Coleoptera in timber in Germany, 19, 229.

Opius crawfordi, parasite of Anastrepha

ludens in Mexico, 395.

Opius tryoni, 290; parasite of Dacus in

Fiji, 161.

Opuntia, proposed utilisation of Melitara dentata against, in California, 242; Diaspis echinocacti on, in Nyasaland, 86.

Opuntia macrorrhiza, Melitara dentata on, in Kansas and Colorado, 241.

Orange, Hemiberlesia rapax on, in Argentina, 4; pests of, in Australia, 92, 268; pests of, in Bermuda, 172, 290; Cryptocephalid on, in Cuba, 116; Icerya montserratensis on, in Ecuador, 95; Mictis profana on, in Fiji, 290; fruit-piercing moths attacking, in India, 391; Anastrepha ludens on, in Mexico, 395; A. serpentina on, in Panama, 397; Lasiocampid on, in Spain, 170; pests of, in U.S.A., 98, 113, 211, 341: favouring Comperiella bifasciata, 92.

Orange, Mandarin or Tangerine, Paratetranychus citri on, in Bermuda, 172; Parlatoria ziziphus on, in Egypt, 228; development of Anastrepha ludens in, 395.

Orange Bug, Bronze (see Rhoecocoris sulciventris).

orbitalis, Lixophaga.

Orchids, Tenthecoris bicolor on, in Brazil, 134; pests of, in Hawaii, 250, 251.

Oregon, forest pests in, 40; pests of leguminous plants in, 100, 101, 259, 342, 343; Aphids and virus disease of Liliaceae in, 1; miscellaneous pests in, 220, 342; injurious Tettigoniids in, 113, 114, 115; parasite of Cydia pomonella in, 240.

Oregon Bait, 142.

Orgyia antiqua, on heather in Britain, 233. Oria musculosa, bionomics of, on cereals in Britain and Russia, 307, 308.

Oriental Fruit Moth (see Cydia molesta).

orientalis, Aonidiella.

Orius insidiosus, predacious on insects in

Utah, **263.**

Orius tristicolor, in Utah, 106, 263; predacious on insects, 263; predacious on, 106.

bionomics of, on Ormenis infuscata, Juniperus bermudiana in Bermuda, 289.

ornata, Solubea.

ornatum, Trogoderma. ornatus, Cheletogenes. ornithogalli, Prodenia.

Ornithogalum, Aphids and virus diseases of, in U.S.A., 1, 38.

orophila, Helopeltis.

Orthodichlorbenzene, in oil emulsion against Asterolecanium, 227; fumigation with, against pests of stored products, 83, 320, 321, 371; and kerosene, 321.

Orthol-K, 212.

Orthoptera, protozoan parasites of, 296. Orthotylus marginalis, predacious on Paratetranychus pilosus in Britain, 234.

oryzae, Calandra.

Oryzaephilus surinamensis, in U.S.A., 69, 102, 258; in Victoria, 318; in dried currants, 296; apparatus for removing, from tea, 74; not feeding on dehydrated vegetables, 102; in stored wheat, 69, 318; penetration of packing materials by, 258; fumigants against, 69, 296; tests and action of inert dusts on, 157, 319.

Oscinella frit, bionomics of, on cereals in Britain, 199, 297, 307; experiments on rearing parasites of, on other hosts, *75.*

ostreata, Pseudoparlatoria.

Othreis, measures against, in orchards in Queensland, 291.

Othreis ancilla, attacking fruits in India, 386.

Othreis fullonia (fullonica), attacking fruits in Queensland and India, 291, 386, 391; bionomics of, 291, 391.

Othreis materna (attacking fruits), bionomics of, in India, 386, 391; in Queens-

land, **291.**

Otiorrhynchus ligustici, on Papaver somniferum in Hungary, 122; baits for, in New York, 179.

Otiorrhynchus ovatus, on strawberry in

Ohio, 263.

Otiorrhynchus singularis, on fruit trees and bushes in Britain, 359, 360.

Otiorrhynchus sulcatus, on Taxus Britain, 171.

Ovaticoccus, n.n., for Gymnococcus, Dougl., 92.

Ovaticoccus agavium, type of genus, 92. ovatus, Otiorrhynchus (Brachyrrhinus). ovinus, Melophagus.

ovipennis, Cymatodera.

ovivorus, Telenomus (Phanurus).

Oxalic Acid, dinitro-o-cresol sprays acidified with, 58.

Oxycanus cervinatus, factors affecting development of, in pastures in New Zealand, 183, 184; baits for, 183.

Oxycarenus, cotton seed as manure, etc., favouring infestation of cotton by, in Uganda, 88.

Ρ.

pabulinus, Lygus.

Pachyceras xylophagorum (see Rhoptro-

Pachymerus lacerdae, bionomics and control of, in field and stored oil-palm nuts in Nigeria, 165, 370.

Pachymerus longus, in stored ground-nuts

in Nigeria, 370.

Pachyneuron altiscuta, reared from Coleophora laricella in Ontario, 143; parasite of Clausenia purpurea in Virginia, 110.

Pachyneuron coccorum, hyperparasite of Coccus hesperidum in Palestine, 326.

Pachyneuron syrphiphagum, hosts of, in Argentina, 117.

Pachypasa albofasciata, on Pinus radiata in S. Africa, 83.

Pachypasa capensis, on Acacia and pines in S. Africa, 83.

Pachypasa sericeofasciata, on Aleurites

and Bridelia in Nyasaland, 86. Pachypeltis, on Combretum in Br. Cam-

eroons, 196. Pachyrhizus erosus, tests of dust prepared from, against insects, 339.

pacificus, Hemerobius; Tetranychus.

packardi, Melanoplus.

Packing Materials, resistance penetration by insects, 102, 258, 272. Paleacrita vernata, rendering trees susceptible to other pests and disease in U.S.A., **43, 173.**

Palestine, Capnodis spp. in, 75, 384; Coccids and their natural enemies on Citrus, etc., in, 225, 325, 326; cinella septempunctata in, 225.

pallens, Pseudodacus (Anastrepha). palliatus, Hylastes (Hylurgops).

pallida, Gilpinia.

pallidipes, Parahormius (Hormius). pallidus, Acerophagus; Tarsonemus.

pallorana, Tortrix (Amelia).

Palm Oil, emulsion of, unsatisfactory against Coccids, 364.

Palmitic Acid, in sprays, 273.

Palmodes laeviventris (see Chlorion).

Palms (see Coconut and Elaeis).

paludosa, Tipula.

palustris, Isotomurus.

Panama, Anastrepha serpentina on orange in, 397; natural enemies and biological control of Coccids in, 313, 374; importation of Eretmocerus serius into Mexico from, 261.

panicea, Sitodrepa (Stegobium).

Paniscus, attacking Pieris rapae in New Zealand, 247.

Paniscus laevis, in Ceylon, 300.

Panolis flammea, outbreak of, in Ger many, 14.

Panoquina spp., on sugar-cane in Porto Rico, **243.**

(White-(Graphognathus) Pantomorus fringed Beetles), in U.S.A., 53, 111, 259: factors affecting infestation by, measures against, 53, 111; legislation against, 111.

Papaver (see Poppies). papaveris, Aphis; Dasyneura; Timaspis. Papaya, Aonidiella aurantii on, in Brazil,

Aphids on, in Hawaii, 251; Toxotrypana curvicauda on, in Mexico, 397; Dacus spp. on, in Queensland, 162; Aspidiotus combreti zizyphi on, in S. Rhodesia, 126; Pseudoparlatoria ostreata on, in W. Indies, 110, 243; D. xanthodes not developing on, 161.

Paper Disks, Tarred, to protect crucifers against Hylemyia brassicae, 178.

papuensis, Pristhesancus.

Parabromobenzenesulphonamide, 99.

Parabromo-N-ethylbenzenesulphonamide, toxicity of, to insects and plants, 279. (Parabromophenyl)-1,1,1-trichlo-

rethane (see DDT Analogues).

Parachlornitrobenzene, insecticidal action

(Parachlorphenyl) $\beta\beta$ -dichlorethylene (see DDT Analogues).

paracornutus, Phyllocoptes.

Paradichlorbenzene, in oil emulsion against Asterolecanium pustulans, 227; uses of, against borers in trees, 353, 365; in mixture against Coleoptera in against Dermestes and timber, 84; clothes moths, 83, 320, 331; against pests in stored bulbs and potatoes, 35, soil treatments with, against pests, 17, 108, 109; ineffective for fumigating ants' nests, 322; formulae containing, 108, 227, 353; injury to plants, 17, 35, 326. causing

Paraformaldehyde, reducing toxicity of

derris dust, 78.

476

Parahormius pallidipes, parasite of Keiferia lycopersicella in Hawaii, 250.

parallela, Aphrophora.

Parallelia algira, on castor in India, 386. parallelipipedus, Lyctus. paranensis, Schistocerca.

Paranitraniline, ineffective as an insecticide, 198.

Parasa lepida, parasite and control of, on tea in Ceylon, 300.

Parasavivida, on plum in S. Rhodesia, 126.

parasae, Apanteles.

Parasetigena silvestris (see Phorocera).

Parasites (of insects), rearing of, on unnatural hosts, **74**, **75**; book on, **357**. (See Hymenoptera, Parasitic.)

Paratetranychus, hibernation of, 328.

Paratetranychus bioculatus (on tea), in Madras, 246; in Nyasaland, 85.

Paratetranychus citri, sprays against, on Citrus in Bermuda, 172.

Paratetranychus coiti, on Platanus U.S.A., 352.

Paratetranychus ilicis, on Platanus

California, 352.

Paratetranychus peruvianus, possibly recorded as Tetranychus peruviensis on cotton in Peru, 132 (and note).

Paratetranychus pilosus (Fruit-tree Red Spider), in Britain, 233, 234, 360; in Canada, 334; in Holland, 253; in Switzerland, 124; in U.S.A., 4, 148, 208, 285, 334, 352; on Platanus, 352; bionomics of, 124, 233; natural enemies of, 208, 233, 234; effects of different sprays on increase and control of, 4, 124, 148, 208, 233, 234, 253, 254, 285, 334, 360; other factors affecting damage by, 334, 360.

Paratetranychus sacchari, injury caused by, on sugar-cane in Porto Rico, 243.

Paratetranychus ununguis, test of Dithane against, on spruce in U.S.A., 379.

Paratheresia diatraeae (see Theresia claripalpis).

Paratillus carus, predacious on Lyctus in Australia and Britain, 181.

Paratolyl Cyanide, action of, on Forficula and woodlice, 198.

Paratrechina longicornis, in Porto Rico, tests of insecticide treatments 339; against, 231, 339.

Paratrioza cockerelli, measures and experiments against, on potato and tomato

in U.S.A., 30, 216, 217.

Paratriphleps laeviusculus, predacious on Heliothis vivescens in Peru, 278.

Paria, measures against, on strawberry in Ohio, 262.

Paris Green, in baits, 68, 140, 183; in dusts, 45, 59, 63, 112; in sprays, 94, 287; apparatus for applying atomised sprays of, 254; test with, against Bruchid in stored cowpeas, 327; use of, against termites, 73; comparison of toxicities of other arsenicals and, 279; and arsenates, 45, 59, 63, 279; and Bordeaux mixture, 94; and pyrethrum, 63; and sulphur, 63, 64.

Parlatoria chinensis, distribution

control of, in U.S.A., 258.

Parlatoria oleae (Olive Scale), liberation of introduced parasite against, in California, 107.

Parlatoria ziziphus, sprays against, on Citrus in Egypt, 228.

Parsnip, Depressaria heracleana on, in Br. Columbia, 333.

Parsnip Webworm (see Depressaria heracleana).

Parthenium argentatum, pests of, in U.S.A., **104, 105, 259.**

partita, Serrodes. parva, Lissonota. parvula, Epitrix.

Passiflora, Coccid on, in Belgian Congo, 365; Dacus xanthodes reared on, in Fiji, 161; Anastrepha chiclayae on, in Mexico, 397.

Passiflora coerulea, Corythaica passiflorae

on, in Argentina, 263.

Passiflora edulis (Passion Fruit), Anastrepha pseudoparallela on, in Argentina, 4; Aphis gossypii on, in Hawaii, 251; Dacus ferrugineus tryoni on, in Queensland, 162.

Dacus (Chaetopassiflorae, Corythaica;

dacus).

Passion Fruit (see Passiflora edulis). pattoni, Eulimneria (Limneria).

Paullinia pinnata, Helopeltis sanguineus on, in Nigeria, 383.

Pavonia zeylanica, Pempherulus affinis on,

in India, 21.

Pea, Lepidoptera on, in Bengal, 128; Scaptocoris terginus on, in Brazil, 4; wireworms causing little damage to, in Britain, 266; pests of, in Canada, 141, 145, 154, 223; Prodenia litura on, in Fiji, 201; Bruchus pisorum in harvested, in France, 297; Macrosiphum solanifolii on, in Hawaii, 251; Cnephasia INDEX. 477

wahlbomiana on, in Holland, 12; pests of, in U.S.A., 22, 49, 100, 214, 260, 276, 277, 282, 283, 338; streak disease of cineraria transmitted to, 70; attacked by Phyllobius pyri, 92: injured by soil treatment with D-D mixture, 56.

Pea, Pigeon (see Cajanus cajan).

Pea Aphis (see Macrosiphum onobrychis). Pea Bruchid (see Bruchus pisorum).

Pea Moth (see Cydia nigricana).

Pea Weevil (see Sitona lineatus). Peach, pests of, in S. America, 94, 117, 397; pests of, in Australia, 322, 323; Caliroa limacina on, in Britain, 360; Lygus oblineatus on, in Br. Columbia, 153; Quadraspidiotus perniciosus on, in France, 125; Tortrix rosana on, in Holland, 12; Anastrepha spp. on, in Mexico, 395, 397; Dacus on, in New Caledonia, 161: Curimosphena fasciculosa on, in S. Rhodesia, 126; pests of, in U.S.A., 46, 54, 105, 213, 314, 315, 316; insecticides and injury to, 39, 54, 62, 314, 379.

Peach-tree Borer (see Aegeria exitiosa).

Pear, Caliroa limacina on, in Argentina, pests of, in Britain, 233, 360; 131; Psylla pyricola on, in Canada, 142, 153; pests of, in Denmark, 12, 382; pests of, in France, 125, 355; Anthonomus piri on, in Germany, 117; insects on, in India, 389, 391; Anastrepha spp. on, in Mexico, 395, 397; Bryobia pretiosa on, in Morocco, 356; Anuraphis piri on, in Sweden, 36; Anthonomus piri on, in Switzerland, 125; pests of, in U.S.A., 22, 153, 222, 240, 258, 284, 285; failure to rear Polychrosis viteana on fruit of, 27; spray destroying stumps and suckers of, 258.

Pear Psylla (see Psylla pyricola).

Pecan, pests of, in Texas, 209, 270, 346, 347.

Pecan Nut Casebearer (see Acrobasis caryae).

Pectinophora gossypiella (see Platyedra). pectinophorae, Chelonus.

Pectis papposa, Plusia brassicae on, in Arizona, 112.

Pediculoides ventricosus (attacking insects), in France, 356; in Germany, 19; in India, 164.

Pelecinus polyturator, bionomics of, in Canada, 180.

pellio, Attagenus. pellionella, Tinea.

Peltophorum, resistance of timber of, to termites in N. Rhodesia, 247.

Pempheres affinis (see Pempherulus). pempheridis, Entedon.

pempheriphila, Euderus.

Pempherulus affinis (Cotton Stem Weevil), in India, 20, 21, 164; in Siam, 20; bionomics of, 20; parasites of, 21, 164.

Pemphigus betae, on beet in Alberta, 144; food-plants and control of, in California, 338.

Pemphigus lactucae, possibly transmitting virus disease of lettuce in U.S.A., 39.

Penetrol, as an emulsifier, 65.

Penicillium, infesting Epitrix hirtipennis in U.S.A., 41.

pennipes, Trichopoda.

Pennisetum purpureum, Melanaspis saccharicola on, in Brazil, 205.

Pennsylvania, miscellaneous pests in, 4, 146, 276, 401.

Pentabromophenol, solutions of, protecting wood against termites, 245.

Pentachlorphenol, solution of, against Coleoptera in timber, 84; tests and uses of, against termites, 69, 244, 245, 272; of little value for treating packing material against insects and fungi, 272; and trichlorbenzene, 69.

Pentalonia nigronervosa, on esculenta in Hawaii, 251; Colocasia probable vector of cardamom mosaic in India, 392; on banana, 392.

Penthestes atricapillus septentrionalis, destroying insects in Utah, 263.

Pentilia egena (predacious on Coccids), in Argentina, 133; in Brazil, 133, 205.

Pepper (see Capsicum and Piper nigrum). Pepper Tree (see Schinus molle).

Peranabrus scabricollis, distribution and feeding habits of, in U.S.A., 114.

Perchlorethylene, as solvent for benzene hexachloride, 257.

peregrinator, Calosoma.

Peregrinus maidis, parasite of, on maize in Hawaii, 250.

peregrinus, Tetracnemus.

Perichares coridon, on sugar-cane Porto Rico, 243.

Peridroma saucia, on cabbage in Br. Columbia, 154; parasite of, 281.

Perilampus laevifrons, P. tristis recorded as, **356.**

Perilampus tristis, hosts of, in France, 355: recorded as P. laevifrons, 356.

Perilitus coccinellae, parasite of Coccinellids in Yunnan, 369.

Periplaneta americana, value of laminated paper wrapping against, 272; insecticide tests on, 64, 347, 368.

Periplaneta australasiae, value of laminated paper wrapping against, 272.

perniciosus, Pseudococcus; Quadraspidiotus (Aonidiella, Aspidiotus, Comstockaspis).

Peromyscus maniculatus osgoodi, feeding on grasshopper eggs in N. Dakota, 96.

Peronea comariana, bionomics and control of, on strawberry in Britain, 360.

Peronospora tabacina, spray against, on tobacco in U.S.A., 54.

Persia, Eurygaster integriceps on cereals in, 302.

persicae, Eulecanium; Myzus.

Persimmon (*Diospyros*), polyphagous insects feeding on, **237**, **270**.

persuasoria, Rhyssa.

pertinax, Coelostethus (Dendrobium).

Pertipan, 182.

Peru, cotton pests in, 132, 278; miscellaneous pests in, 5, 131, 397; natural enemies of *Heliothis virescens* in, 278; plant-quarantine restrictions in, 286; beneficial insects imported into other countries from, 261, 312.

peruella, Mescinia. peruviana, Acaulona.

peruviana, Acauiona.

peruvianus, Paratetranychus (Tetranychus). peruviensis, Tetranychus.

Petrobia lapidum, measures against, on onion in Britain, 361.

Petrol, fumigation with, against insects in stored seeds, 371, 375; soil treatment with, against Cotinis nitida, 345.

Petroleum Ether, as a solvent, 244. Phacopteron, on cacao in Belgian Congo, 366. Phaedon armoraciae var. concinnus, bionomics of, on horse-radish in Denmark, 381.

Phaedon cochleariae, on crucifers in Britain, 257, 405; bionomics of, 405; gammexane against, 257.

Phaeogenes, parasite of Coleophora laricella in Ontario, 143.

Phaeogenes epinotiae, 143.

phaeorrhoea, Nygmia.

Phanuropsis semiflaviventris, parasite of Mecistorhinus in Trinidad, 147.

Phanurus ovivorus (see Telenomus). phaseoli, Agromyza (Melanagromyza).

Phaseolus vulgaris (see Beans).

Phasia, parasite of Eurygaster integriceps in Russia, 303.

Pheidolacanthinus (see Pheidole).

Pheidole spp., measures against, collecting seeds of tobacco, etc., in Queensland, 369, 370.

Pheletes (see Limonius).

Phenacoccus aceris (on apple), sprays against, in Br. Columbia, 153, 154; in U.S.A., 3.

Phenacoccus gossypii, on lima bean and egg-plant in Hawaii, 250.

Phenacoccus solani, biological control of, on potato sprouts in insectaries in California, 61.

Phenanthrene, treatment of timber with, against termites, 244.

Phenol (see Carbolic Acid).

Phenothiazine, spraying with, against

Lepidoptera, 28, 149, 258, 322, 323, 324; spraying with, against other insects, 180, 191; dust of, not protecting bulbs from Merodon equestris, 24; tests of toxicity of, to insects, 198, 260, 368; micronised form of, 28, 258, 260; formulae containing, 24, 28, 149, 258, 323; and bentonite, 180; and lead arsenate, 258; and oil emulsion, 149; wetting agents for, 149; residues of, on apples, 149; Parthenium argentatum not injured by dusts containing, 105. Phenothiazone, effects of, on Periplaneta

americana, 368.

phidilealis, Hellula.

Philaenus leucophthalmus, bionomics and control of, on strawberry in Ohio, **262**. philanthus, Hoplia.

Philippines, parasite of Saissetia nigra in, 374.

Philothalpus analis, predacious on Anastrepha ludens in Mexico, 395.

Phleum pratense, Diabrotica undecimpunctata on, in California, 30.

Phonoctonus, predacious on Dysdercus in Tanganyika, 88.

Phonoctonus principalis, predacious on Dysdercus in Belgian Congo, 367.

Phoracantha semipunctata, on Eucalyptus in S. Africa, 83.

Phorbia (see Hylemyia).

Phorocera hamata, parasite of Gilpinia hercyniae in Quebec, 223.

Phorocera silvestris, introduced parasite of Lymantria dispar in U.S.A., 238.

Phorodon humuli (on hops), in Britain, 361; insecticides against, in New York, 317, 318.

Phosphorus, on cacao in Belgian Congo, 366.

Phthalic Glyceryl Alkyd Resin, uses of, in sprays, 54, 148, 149, 211.

Phthalonitrile, dusting with, against *Protoparce* spp., **59**.

Phthorimaea (see Gnorimoschema).

phthorimaeae, Chelonus.

phycidis, Habrocytus.

Phyllobius oblongus, resistance of, to derris dusts, 381.

Phyllobius pyri, bionomics of, attacking wheat, etc., in Britain, 92.

Phyllocoptes destructor (on tomato), dust against, in Colorado, 219; in Hawaii, 250; on Solanum nodiflorum, 250.

Phyllocoptes oleivorus (see Phyllocoptruta). Phyllocoptes paracornutus, sp. n., on almond in California, 111.

Phyllocoptes schlechtendali, bionomics and control of, on pear in Washington, 285.

Phyllocoptruta oleivorus (on Citrus), spray against, in Bermuda, 172; cultural practices not affecting infestation by, in Florida, 341.

Phyllodecta vitellinae, 198.

Phyllopertha (see Anomala).

Phyllophaga (see Lachnosterna).

Phyllotreta, dusts against, on rape in Denmark, 381.

Phyllotreta nemorum (Turnip Flea-beetle), benzene hexachloride against, in Britain, 256.

Phylloxera vitifoliae (vastatrix), races of, on vines in Switzerland, 124; evolution of races of, 118.

Phymatodes dimidiatus, in building timber

in Br. Columbia, 287.

Phyrdenus divergens, bionomics and control of, on solanaceous plants Brazil, 287.

Phyrdenus muriceus, bionomics and control of, on solanaceous plants in S. America and U.S.A., 287.

Physalis peruviana, mites on, in Ceylon,

Physokermes insignicola, introduced Coccinellid predacious on, on Pinus radiata in California, 282.

Phytalus smithi (see Clemora).

Phytocoris tiliae, predacious on Paratetranychus pilosus in Britain, 234.

Phytodecta fornicata, bionomics of, on lucerne in Rumania, 121.

Phytolyma lata, bionomics and control of, on Chlorophora excelsa in Uganda, 329. phytolymae, Psyllaephagus.

Phytomyza lateralis, hot water treatment of chicory against, in Belgium, 120.

Phytomyza solani, on tomato in Denmark, 381.

Phytomyza spicata, parasite of, on maize in Fiji, 290.

Phytophaga destructor (see Mayetiola). Phytophthora infestans, on potato Ú.S.A., 206, 207; DDT against, 207. effectiveness

Phytoptus hedericola, sp. n., on ivy in

California, 111. Picea (see Spruce).

Picea abies (excelsa), pests of, in Finland,

Picea pungens, injured by fumigation with methyl bromide, 50.

piceae, Chermes.

piceus, Attagenus; Mecistorhinus.

picipes, Rhabdopterus.

picta, Ceramica.

pictus, Cirrospilus; Poekilocerus (Poeci-

Pieris, toxicity of gammexane to, 257. Pieris brassicae, dusts against, on cabbage in Denmark, 381; experiments with Ernestia consobrina and, in Russia, 305; tests of insecticides on, 197, 198.

Pieris mesentina (see Anaphaeis aurota).

Pieris monuste (see Ascia).

Pieris vapae (on crucifers), in Australia,

322; in Fiji, 290; in New Zealand, 247, 376; in Russia, 305; in U.S.A., 48, 49, 338, 348; attraction of Verbena bonariensis for adults of, 348; parasites and biological control of, 247, 322; seldom attacked by Ernestia consobrina, 305; dusts against, 48, 49, 322, 338; tests of insecticides on, 197, 198, 278, 279; comparison of food-consumption of Plutella maculipennis and, 376.

Pig Bristle, Anthrenus fasciatus damaging,

Pigeon Pea (see Cajanus cajan).

Pigs, use of, against insect pests, 141, 341. pilifrons, Mastrus.

pilleriana, Sparganothis.

pilosus, Paratetranychus. Pimento (see Capsicum).

pimpinellae, Anthrenus.

Pimpla aequalis, parasite of Polychrosis viteana in U.S.A., 28.

Pimpla instigator, bionomics of, parasitising Dasychira pudibunda in Sweden,

Pimpla sanguinipes, parasite of Cydia pomonella in California, 240.

Pimpla terebrans (see Ephialtes).

pinastri, Sphinx.

Pine, pests of, in S. Africa, 83, 84, 324; pests of, in Canada, 9, 313, 314; pests of, in Finland, 11; pests of, in Germany, 6, 14; Xyleborus camphorae on, in Mauritius, 371; Sirex noctilio on, in New Zealand, 269; Sphinx pinastri on, in Poland, 6; pests of, in U.S.A., 4, 43, 107, 282, 286, 287, 313, 314, 398; billets of, as traps for Hylastes opacus,

Pine, Jack (see Pinus banksiana). Pine, Monterey (see Pinus radiata).

Pine, Mugho (see Pinus mugo). Pine, Pitch (see Pinus rigida).

Pine, Ponderosa (see Pinus ponderosa).

Pine, Red (see Pinus resinosa). Pine, Scots (see Pinus sylvestris).

Pine, Shortleaf (see Pinus echinata).

Pine, White (see *Pinus strobus*).

Pine Geometrid (see Bupalus piniarius). Pine Oil, not protecting wood from termites, 244; spray formula containing,

Pine Sphingid (see Sphinx pinastri).

Pine Tar, unsatisfactory as barrier against

Blissus leucopterus, 57.

Pineapple, Pseudococcus brevipes on, in Belgian Congo, 365; types of injury to, by P. brevipes in Hawaii, 249, 345; Batrachedra on, in Porto Rico, 97; Diaspis bromeliae on, in Queensland,

pinguis, Agromyza (Ophiomyia). pini, Diprion; Pissodes.

piniarius, Bupalus.

piniperda, Myelophilus (Blastophagus). Pinnaspis aspidistrae, on Annona and Citrus in Argentina, 4, 133; characters distinguishing Unaspis citri from, 133. Pinnaspis theae, 246.

Pinus (see Pine).

Pinus banksiana, pests of, in Canada and U.S.A., 8, 107, 313, 314.

Pinus contorta, strain of Harmologa fumiferana on, in Montana, 314.

Pinus echinata, Dendroctonus frontalis and blue-stain fungi in, in U.S.A., 286, 287.

Pinus insignis (see P. radiata). Pinus mugo, sawflies on, in Pennsylvania, 4. Pinus patula, Pachypasa capensis on, in

S. Africa, 83.

Pinus ponderosa, pests of, in U.S.A., 314, 398; forms of, resistant to Retino-

diplosis, 398.

Pinus radiata, pests of, in S. Africa, 83; Physokermes insignicola on, in California, 282; Sirex noctilio on, in New Zealand, 269.

Pinus resinosa, Aphrophora spp. on, in

U.S.A., 107.

Pinus rigida, Dendroctonus spp. on, in U.S.A., 43.

Pinus strobus, Neodiprion lecontei on, in Pennsylvania, 4.

Pinus sylvestris, pests of, in Canada and U.S.A., 4, 314.

Pinus taeda, Xyleborus camphorae on, in Mauritius, 371.

Piophila casei, fumigation against, in cheese factories in Wisconsin, 343.

Piper cubeba, tests of extracts of, against Culex fatigans, 65.

Piper guineense, Odoniella rubra on, in Fr. Togoland, 196.

Piper nigrum, experiments with, as an insecticide, 65.

piri, Anthonomus; Anuraphis (Yezabura). pisi, Bruchus (see B. pisorum); Macrosiphum (see M. onobrychis).

pisorum, Bruchus.

Pissodes harcyniae, on spruce in Finland, 7, 12; parasites of, 12.

Pissodes pini, parasites of, in Finland, 11, 12.

Pissodes validirostris, parasites of, in Finland, 11, 12.

Pistacia, Capnodis cariosa on, in Palestine, 75.

Pisum sativum (see Pea).

Pitch, preparation of soap from, as adhesive for calcium arsenate, **202**.

Pittosporum, Icerya purchasi on, in France, 357.

Pityogenes (see Ips).

Pityophthorus nigricans, on Parthenium argentatum in California, 105.

plaesiosema, Gnorimoschema.

Plaesius javanus (predacious on Cosmo-

polites sordidus), imported into Honduras, 261; establishment of, in Tahiti, 202.

Plagiodera versicolora, in Canada, 71; on willow and poplar in U.S.A., 71.

Plagiognathus albatus, on Platanus in U.S.A., 353.

Plagiognathus arbustorum, predacious on Paratetranychus pilosus in Britain, 234. Plagionotus floralis, measures against, on

lucerne in Russia, 305.
planaris, Corythaica (see C. passiflorae).

planicollis, Lyctus.

Plant Pest Legislation, in Britain, 232; in U.S.A., 111, 145, 193, 286; summaries of, in other countries, 111, 145, 146, 286, 328.

Plants, studies of, as sources of insecticides, 48, 54, 65, 106, 135, 179, 192, 338. platani, Rhyncaphytoptus; Stomacoccus. platanoides, Drepanosiphum.

Platanus spp., pests of, in U.S.A., 352,

353.

platensis, Aphidius. platycephalus, Egius.

Platyedra gossypiella (Pink Cotton Bollworm), in Fr. W. Africa, 165; in Algeria, 165; in Brazil, 375; in Belgian Congo, 367; in India, 390; in Mexico, 286; in Nyasaland, 85; in Pacific Is., 165; in Porto Rico, 97; in Sudan, 87; in Tanganyika, 88; in Texas, 261, 286; legislation against, in U.S.A., 111, 286; intercepted in seed cotton in U.S.A., 100; resistance of Gossypium thurberi to, 87; bionomics of, 261; parasites and biological control of, 261, 367; not parasitised by Rogas testaceus, 390; treatment of cottonseed against, 286, 375.

Platynota flavedana, on strawberry in Ohio, 263.

Platyplectrus natadae, parasite of Limacodids in Ceylon, 300.

Platypus subgranosus, factors influencing infestation of fire-killed timber by, in Victoria, 377.

Platypus wilsoni, in fire-killed Douglas fir in U.S.A., 40.

Plecoptera reflexa, new parasite of, on Dalbergia sissoo in India, 390.

plecopterae, Rogas.

Plesiocoris rugicollis, on apple in Britain, 234.

plesiotypus, Aenoplex (see Mastrus pilifrons).

Plodia interpunctella, in Canada, 225, 403; in Germany, 230, 231; in Porto Rico, 312; in U.S.A., 102, 258; in stored cereals, 225, 312, 403; effects of damage to wheat by, 403; in other stored products, 102, 230, 231; bionomics of, 231; measures and experiments against,

INDEX. 481

225, 231, 312; penetration of packing materials by, 258.

Plum, Quadraspidiotus perniciosus on, in Argentina, 132; pests of, in Britain, 233, 360; Nygmia phaeorrhoea on, in Bulgaria, 119; pests of, in Denmark, 380, 381, 382; pests of, in India, 388, 391: Anoplognathus on, in N. S. Wales, 162; pests of, in Nyasaland, 86; Parasa vivida on, in S. Rhodesia, 126; Paratetranychus pilosus on, in Switzerland, 124; pests of, in U.S.A., 222, 315; Myzus persicae overwintering on wild, in Maine, 2; Thomasiniana theobaldi not developing on, 308:

Plum Aphis, Mealy (see Hyalopterus arundinis).

Plum Curculio (see Conotrachelus nenu-

phar).

brassicae, in Hawaii, 250; in U.S.A., 48, 49, 112; on cabbage, 48, 49; on lettuce, etc., 112, 250; bionomics of, 112; dusts against, 48, 49, 112.

Plusia chalcites, parasites of, in Fiji, 290. Plutella maculipennis (on crucifers), in S. Africa, 293; in Fiji, 269, 290, 291; in New Zealand, 269, 376; in Porto Rico, 339; in U.S.A., 49, 260; effect of nutrition on adults of, 376; parasites and biological control of, 269, 291, 376; insecticides against, 49, 260, 293, 339; comparison of damage by Pieris rapae and, 376.

Poa bulbosa, Dociostaurus maroccanus attracted to, 200.

Podagrica puncticollis, repellent against, on cotton in Sudan, 87.

Podisus, predacious on Gilpinia hercyniae in Quebec, 224.

Podisus maculiventris, on peach in Illinois,

Podonectria coccicola, forms of, infesting Lepidosaphes beckii in Belgian Congo,

poecila, Solubea (Mormidea).

Poecilocerus (see Poekilocerus).

Poecilophila maculatissima, on apple and plum in Nyasaland, 86.

Poekilocerus pictus, on bamboos in India,

Poland, pests of apple in, 117; forest pests in, 6, 9; organisation of plant protection in, 5.

politus, Chilocorus. Pollard (see Bran).

Pollination, relation of insects to, 78, 245.

polychrosidis, Ischnus.

Polychrosis botrana (Vine Moth), investigations on, in Germany, 13, 14; foodplants of, in Italy, 118; in Spain, 298.

Polychrosis viteana, bionomics and control of, on vines in U.S.A., 27, 28, 258. Polygonum, host of Macrocentrus spp. on, in New Jersey, 47.

Polyhedral Disease, of Lepidoptera, 7, 8,

Polyommatus boeticus (see Cosmolyce).

Polyscelis, failure to rear, on Drosophila, *75.*

polytoma, Gilpinia. polyturator, Pelecinus.

Polytus mellerborgi, on banana in Fiji, 202. Pomegranate, Anastrepha ludens on, in

Mexico, 395. pometaria, Alsophila.

pomi, Aphis (Doralis).

pomonella, Cydia (Carpocapsa); Rhago-

pomorum, Anthonomus.

Popillia japonica, in Canada, 137, 224; in U.S.A., 49, 50, 55, 58, 63, 79, 180, 347, 402; factors affecting damage to maize by, 49; on other plants, 180, 402; tests and use of Bacillus popilliae against, 79, 280; treatments of plant roots and soil against, 50, 58, 63, 137, 393; repellent sprays against, 180; baits for, 55, 347.

Poplar, Capnodis miliaris on, in Palestine,

75. (See Populus spp.)

Poppies (Papaver spp.), Xylena exoleta on, in Denmark, 382; pests of, in Hungary,

Poppy, Prickly (see Argemone alba var. glauca).

Population Studies (of insects), 44, 56, 129, 254, 293, 296, 405.

Populus (see Poplar).

Populus deltoides, Teragra guttifera on, in S. Africa, 83.

Populus grandidentata, Malacosoma disstria on, in Br. Columbia, 155.

Populus nigra italica, Plagiodera versicolora on, in U.S.A., 71.

Populus tremula, Stilpnotia salicis on, in Sweden, 32.

Populus tremuloides, pests of, in N. America, 155, 271.

Porizonidea (Porizon), imported into U.S.A. against *Listroderes obliquus*, **261.** porosa, Capnodis.

Porotermes adamsoni, introduced into New Zealand in timber, 234; characters

of, 234, 235.

Porphyronota maculatissima (see Poecilophila).

Porthetria (see Lymantria).

Porto Rico, cotton pests in, 97, 278, 312; identity of Tingid on egg-plant in, 263; Anastrepha mombin praeoptans on mango in, 52, 397; miscellaneous pests in, 97, 110, 243, 312, 339; new Pentatomid on rice in, 147; sugar-cane pests in, 26, 97, 243, 312; termites in, 243, 244; beneficial insects and biological control

in, 26, 97, 110, 243, 312, 374; beneficial insects and toad imported into other countries from, 261, 313; summary of plant quarantines in, 145; pests intercepted in quarantine in, 223; plants as sources of insecticides in, 338.

Portugal, Lepidoptera on cork oak in, 309, 310; Quadraspidiotus perniciosus on fruit trees in, 125; natural enemies and biological control of Lymantria dispar in, 310; legislation against introduction of Leptinotarsa decemlineata into Britain from, 232.

postica, Hypera (see H. variabilis).

postvittana, Tortrix.

Potash Fish-oil Soap (see Soap).

Potassium Cyanide, in sprays against Rhoecocoris sulciventris, 268.

Potassium Oleate, as a wetting agent for

sprays, 169.

Potato, Gnorimoschema operculella on, in S. Africa, 326; pests of, in Argentina, 4; pests of, in Australia, 31, 33, 34, 93, 184-187, 322, 323, 378, 404; Leptinotarsa decemlineata on, in Belgium, 157, 158; pests of, in Brazil, 264, 287, 288, 289; pests of, in Britain, 266, 358, 359; pests of, in Canada, 139, 154; mites on, in Ceylon, 299; Aphids on, in Belgian Congo, 365; Galeruca tanaceti on, in Denmark, 381; Epilachna vigintioctopunctata on, in Fiji, 290; pests of, in France, 157, 158, 298, 331, 357; L. decemlineata on, in Germany, 157, 158; pests of, in Hawaii, 250, 251; L. decemlineata on, in Holland, 157; pests of, in India, 300; L. decemlineata on, in Luxemburg, 158; Myzus persicae on, in S. Rhodesia, 126; Mamestra brassicae on, in Russia, 305; L. decemlineata on, in Spain, 165, 166; L. decemlineata on, in Switzerland, 125, 157; pests of, in U.S.A., 2, 22, 23, 30, 39, 48, 51, 61, 146, 173, 180, 206, 207, 214, 217, 220, 260; insects and virus diseases of, 2, 33, 39, 173, 323, 365, 378; experiments with hybrids of Solanum demissum and, against L. decemlineata, 159; relation of wireworms to, 23, 266, 294; Coccids reared on sprouts of, 61, 109; insecticides causing injury to, 22, 159; (tubers), restrictions on importation of, into Britain against L. decemlineata, 232; Popillia japonica imported into Canada in, 137; regulations relating to importation of, into U.S.A. from Mexico, 111; Gnorimoschema operculella in, 31, 146, 186, 212, 300, 326, 404; (dehydrated), insects in, in California,

Potato Beetle, Colorado (see Leptinotarsa decemlineata).

Potato Flea-beetle (see *Epitrix cucumeris*).

Potato Flea-beetle, Eastern (see Epitrix tuberis).

Potato Flea-beetle, Western (see Epitrix subcrinita).

Potato Leafhopper (see Empoasca fabae). Potato Mealybug (see *Phenacoccus solani*).

Potato Psyllid (see Paratrioza cockerelli). Potato Tuber Moth (see Gnorimoschema

operculella).

potentillae, Capitophorus.

praetiosa, Bryobia.

Praon aguti, sp. n., hosts of, in U.S.A., 96. Prasoidea sericea, on Pinus radiata in S. Africa, **83.**

prelli, Chermes.

Prenolepis longicornis (see Paratrechina). pretiosum, Trichogramma.

pretiosus, Tetracnemus. Prickly Ash (see Zanthoxylum clavaherculis).

Prickly Pear (see Opuntia).

Prickly Poppy (see Argemone).

principalis, Phonoctonus. prismaticus, Labrorhycus.

Pristhesancus papuensis, predacious on Rhoecocoris sulciventris in N. S. Wales,

Pristiphora californica, bionomics and control of, on pear in Washington, 285. Pristiphora geniculata, aerosol against, on Sorbus aucuparia in Quebec, 142.

Pristomerus vulnerator (parasite of Cydia pomonella), bionomics of, in France, 355; suggested introduction of, into Canada, 356.

Privet (see Ligustrum).

Probits, for expressing toxicity of insecticides and Bacillus popilliae, 188, 277, 280, 345.

Proceras argyrolepidus, on maize and

sorghum in Tanganyika, 85.

Process sacchariphagus, utilisation of Trichogramma australicum against, on sugar-cane in Mauritius, 370.

Prochiloneurus clavatus, parasite of Pseudococcus perniciosus in Nyasaland, 86.

Prociphilus betae (see Pemphigus). Prodenia litura, life-cycle of, on cauli-

flower in Bengal, 363; food-plants and parasites of, in Fiji, 201, 202; on flax in Nyasaland, 86.

Prodenia ornithogalli, parasite of, 281.

productus, Criocephalus. profana, Mictis.

prolixus, Rhodnius.

Prontaspis citri (see Unaspis).

Propane, use of, for producing aerosols,

Prophyscus latiscapus, gen. et sp. n., parasite of Quadraspidiotus perniciosus in Argentina, 132.

2-n-Propyl-pyridine, as a fumigant, 275.

Propylene Dichloride (see Dichlorpropane).

Prospattella berlesei, parasite of Aulacaspis cinnamomi in Egypt, 228.

Prosthogonimus, relation of dragonflies to, 5.

Protaphidius (Coelonotus), subgen. of Aphidius, 96.

Protoparce quinquemaculata, dusts against, on tomato in Virginia, 59.

Protoparce sexta, measures against, on tomato and tobacco in U.S.A., 59, 215, 216; pupal stage of, 215, 216; adult morphology of, 192.

Protopulvinaria longivalvata, on jasmine and avocado in Brazil, 205.

Protozoa, in Orthoptera, 296.

provancheri, Sagaritis.

proximus, İps. pruinosa, Sericesthis.

pruinosum, Eulecanium (Lecanium).

Prune, Tetranychus on, in Britain, 360; Hyalopterus arundinis on, in Br. Columbia, 153.

pruni, Hyalopterus (see H. arundinis).

prunifoliae, Rhopalosiphum.

Prunus spp., Paratetranychus pilosus on, in Britain, 233; Anastrepha spp. on, in Mexico, 397; Capnodis on, in Palestine, 75; toxicity of Bordeaux mixture to, 62. (See Cherry, Peach, Plum, etc.)

Psallus ambiguus, predacious on Paratetranychus pilosus in Britain, 234.

Psallus seriatus, dusts against, on cotton in Texas, 261.

Pselactus spadix (see Codiosoma).

Pseudaonidia trilobitiformis, food-plants of, in Belgian Congo, 365.

Pseudaphycus, establishment of, against Pseudococcus comstocki in U.S.A., 79, 208, 209; bionomics of, 209.

Pseudaphycus notativentris (see Acerophagus).

pseudobrassicae, Rhopalosiphum.

Pseudococcus, biological control of, on vines in California, 214.

Pseudococcus adonidum, in California, 59, 109; in Nyasaland, 86; food-plants of, 59, 86; natural enemies and biological control of, 59, 86, 109.

Pseudococcus bakeri (see P. maritimus).

Pseudococcus brevipes (on pineapple), in Belgian Congo, 365; types of damage by, in Hawaii, 249, 345.

Pseudococcus citri, 196; in California, 109; on Citrus in Palestine, 225; natural enemies and attempted biological control of, 109, 225.

Pseudococcus comstocki, natural enemies and biological control of, on apple in U.S.A., 78, 109, 208, 209, 222, 258; new predator of, in Uzbekistan, 328.

Pseudococcus exitiabilis, food-plants and

relation of, to virus disease of cacao in W. Africa, 66, 196.

Pseudococcus gahani, parasite of, in California, 109.

Pseudococcus longispinus (see P. adonidum).

Pseudococcus maritimus, 214; effect of Acerophagus pallidus on rearing of, on potato sprouts in California, 61; foodplants of, in Egypt, 227.

Pseudococcus perniciosus, natural enemies and control of, on Aleurites in Nyasa-

land, 86.

Pseudococcus virgatus (see Ferrisia).

Pseudodacus pallens, on Bumelia spiniflora in Mexico and Texas, 397; systematic position of, 397.

pseudomagnoliarum, Coccus. pseudoparallela, Anastrepha.

Pseudoparlatoria ostreata, Chilocorus cacti predacious on, on papaya in W. Indies, 110, 243.

Pseudoperonospora humuli, spray against, on hops in New York, 317.

pseudosolani, Myzus (see Macrosiphum solani).

Pseudotsuga taxifolia, Melanolophia imitata on, in Br. Columbia, 180; pests of, in U.S.A., 40, 314; Cnephasia longana ovipositing on, 342.

pseudotsugae, Dendroctonus.

psidii, Pulvinaria.

Psila rosae, experiments with carrot varieties and, in Sweden, 35; DDT against, in Switzerland, 288.

Psophocarpus tetragonolobus, mites on, in Ceylon, 299.

Psychophagus omnivorus, parasite of Gilpinia hercyniae in Quebec, **224**.

Psylla costalis, on apple in Switzerland, 125.

Psylla mali, sprays against, on apple in Europe, 35, 117, 253.

Psylla pyricola (on pear), in Br. Columbia, 142, 153; in Washington, 22, 258, 285; measures against, 22, 142, 258.

Psyllaephagus phytolymae, parasite Phytolyma lata in Uganda, **329.**

Psyllid Yellows, Paratrioza cockerelli causing, in potato and tomato in Colorado, 30.

Psylliodes attenuata (Hop Flea-beetle), tests of inert dust against, in Germany, 231.

Pterandrus rosa (see Ceratitis).

Pterocarpus angolensis, resistance of timber of, to termites in N. Rhodesia, 247.

Pteromalus azureus (see Rhopalicus).

Pteronidea (see Nematus). Pteronus (see Nematus).

Ptinids, key to larvae of, associated with stored products in Britain, 135.

Ptinus brunneus (see P. hirtellus).

INDEX. 484

Ptinus fur, experimentally infesting dehydrated vegetables, 102.

Ptinus hirtellus, in timber in U.S.A., 176. Ptinus tectus, infesting drugs in Germany,

230; action of inert dusts on, 157. pubescens, Ophonus (see Harpalus rufipes). pudibunda, Dasychira.

pugnax, Solubea. pulchra, Tragocephala.

pulla, Stictiella.

Pulvinaria jacksoni, Coccinellids predacious on, on Aleurites in Nyasaland,

Pulvinaria psidii, on Ficus and mango in Egypt, 228.

Pulvinaria vitis, spray against, on Platanus in U.S.A., 353.

pulvinariae, Coccophagus.

Pumice Dust, experiments with, against insects, 131.

pumila, Agathis (Microdus).

Pumpkin, Margaronia indica on, in Fiji, 290; less affected than squash by Anasa tristis in U.S.A., 21, 179.

punctata, Hypera. punctatum, Anobium.

punctatus, Bracon (Microbracon).

puncticollis, Podagrica. punctipennis, Nabis. punctipes, Geocoris.

punctiventris, Cleonus (Bothynoderes).

Punjab, cotton pests in, 296, 391; soil fauna of cotton fields in, 362; Acridids in forests in, 389.

purchasi, Icerya.

purpurea, Clausenia.

pustulans, Asterolecanium.

putrescentiae, Tyrophagus.

Pycnoderes quadrimaculatus, new parasite of, introduced from Mexico into Hawaii, 251.

pygmaeus, Bracon (Microbracon); Cephus. Pyrausta coclesalis, bionomics and dis-

tribution of, in Asia, 388.

Pyrausta nubilalis (European Corn Borer), in Canada, 137-139, 224; in U.S.A., 3, 48, 49, 101, 151, 239, 259, 317, 402; on plants other than maize, 48, 139, 317, 402; bionomics and strains of, 3, 138, 139, 151, 224; parasites and biological control of, 138, 139, 259, 281; other measures against, 3, 48, 49, 138, 139, 224, 239.

Pyrax, 283, 335.

Pyrax ABB, 49, 207.

Pyrene, tests with, for treating timber against termites, 244.

Pyrethrins, content of, in pyrethrum preparations, 3, 30, 51, 63, 64, 104, 108, 127, 142, 180, 189, 190, 274.

Pyrethrum (including extracts), against Aphids, 50, 51; against Capsids, 29, 37, 38, 64, 66, 153, 260; against Jassids, 28, 104, 180; against other Hemiptera, 21, 63, 64, 179, 260, 353, 374; against Coleoptera, 3, 31, 38, 84, 104, 125, 168, 180, 231, 233, 402; against Diptera, 4, 142, 255; against Lepidoptera, 31, 51, 108, 112, 119, 127, 223, 231, 238, 260, 274, 333; repellent effect of mixtures containing, on full-grown larvae of Cydia pomonella, 274; in oil, treatment of maize ears with, against Heliothis armigera, 260, 337; against sawflies, 142, 360; action of, on Thermobia domestica, 340; against thrips, 51, 52, 104; aerosols of, 142, 189, 238, 261; dusting with, 3, 4, 21, 29, 30, 37, 52, 63, 66, 84, 104, 112, 119, 142, 153, 168, 180, 223, 231, 233, 260, 333, 360, 402; carriers for, 37, 112, 180; bentonite reducing toxicity of, 282; proprietary dusts of, 30, 63, 64, 153, 180; proprietary spray mixtures containing, preparation of concentrates of, 189, 332; and nicotine, 260; and oil emulsions, 38, 51, 108, 274; atomised sprays of oil or water containing, 31. 38, 127, 255; and Paris green, 63; and rotenone insecticides, 28, 51, 104, 119, 180, 231, 233; and soap, 51, 125, 168; other wetting agents for, 51, 255; and sulphur, 63, 168, 260; and thiocyanates, 30, 104.

Pyrethrum Plants, fungus associated with injury to, by Thrips tabaci in Tanganyika, 85.

pyri, Eriophyes; Phyllobius.

pyricola, Psylla.

Pyridine, in baits for Trypetids, 216, 340; treatment of stored potatoes with dust containing, against Gnorimoschema operculella, 31, 187.

Pyridine Derivatives, tests and uses of, as

insecticides, 275, 276.

Pyrilla, method of estimating populations of, on sugar-cane in India, 390. pyrivora, Contarinia.

Pyrocide, 30, 63, 153, 180.

Pyroderces rileyi, feeding-habits and control of, on orange in California, 113. Pyrophorus luminosus, imported

U.S.A. against Lamellicorns, 261. Pyrophyllite, as a carrier for dusts, 48,

49, 207, 283, 335.

Quadraspidiotus juglans-regiae, spray against, on Platanus in U.S.A., 353. Quadraspidiotus perniciosus, in Argentina, 132; in Austria, 117; in Canada, 64, 332; in Caucasus, 125; in France, 5, 125, 356, 357; in India, 391; in Italy, 125; in Spain and Portugal, 125; risk of spread of, into Switzerland, 125; in U.S.A., 22, 58, 284, 353; on fruit trees, 5, 22, 58, 125, 132, 284, 356, 357, 391; comparison of injury to apple by Aphids and, 332; on currant and gooseberry, 64; on other plants, 5, 353; natural enemies of, 132, 357; sprays against, 22, 58, 284, 353; fumigation of nursery stock against, 125.

quadricollis, Cathartus. quadridens, Ips (Pityogenes).

quadridentata, Ascogaster.

quadrigemina, Chrysomela (Chrysolina). quadrimaculatus, Anopheles; Pycnoderes. quadripustulatus, Exochomus.

quadriseriata, Ctenolepisma.

Quarantine, pests intercepted in: in Hawaii and Porto Rico, 223; in U.S.A., 100, 223.

Quartz Dusts, experiments with, against insects, 131; as diluent for rotenone dusts, 283.

quasi-ingrami, Forcipomyia.

Quassia, and soap, spraying with, against *Hoplocampa* spp. and Aphids, **334**, **380**. *Quaylea whitteri* hyperparasite of *Sais*-

Quaylea whitteri, hyperparasite of Saissetia nigra in California and Hawaii, 374.

Quebec, forest pests in, 8, 71, 142, 143, 223, 313, 354; Pyrausta nubilalis on maize in, 138, 224; miscellaneous pests in, 142, 224; Cydia nigricana on peas and beans in, 139, 141, 144, 145, 223; Lachnosterna in, 141, 181; Popillia japonica in, 137; beneficial insects and biological control in, 138, 143, 181, 223, 224.

Queensland, pests of Araucaria cunninghami in, 32; Heliothis armigera on cotton in, 32, 87; fruit pests in, 32, 162, 248, 291; pests of stored products in, 31, 320, 404; ants on seedling tobacco in, 369; natural enemies and biological control of insects in, 32, 292, 320; establishment of Teleonemia scrupulosa against Lantana in, 32, 321.

Quercus (see Oak). Quercus ilex var. suber, Lepidoptera on,

in Portugal, 309, 310.

Quercus stellata, Lachnosterna spp. on, in Texas, 270.

Ouince, Cydia molesta on, in Brazil, 94; Caliroa limacina on, in Britain, 360; Lepidoptera on, in France, 355, 356; Anastrepha spp. on, in Mexico, 395, 397; Thomasiniana theobaldi not developing on, 308.

quinquefasciatus, auct., Culex (see C.

fatigans).

quinquemaculata, Protoparce.

Quisqualis, Anua coronata on, in India, 386.

R.

radicicola, Trifidaphis.

Radish, Athalia rosae on, in Denmark, 381; Crocidolomia binotalis on, in Fiji, 290; Aphids on, in Hawaii, 251; pests of, in U.S.A., 67, 177, 178, 276; injured by pyridine derivatives, 276.

Railway Waggons, fumigation of, against

pests of stored products, 259.

Rainfall, relation of locusts and grass-hoppers to, 76, 167, 199; relation of other insects to, 34, 37, 66, 182, 183, 233, 292, 303.

Raisins (see Fruit, Dried).
rapae, Ceuthorrhynchus; Pieris.

rapax, Hemiberlesia.

Rape, Hellula phidilealis on, in Brazil, 95; Meligethes aeneus on, in Czechoslovakia, 252; pests of, in Denmark, 381; M. aeneus on, in Germany, 36; thrips on, in Sweden, 36.

Rape Beetle (see Meligethes aeneus).

Raphanus raphanistrum, Myzus persicae on, in U.S.A., 2.

Raphanus sativus (see Radish).

Raphanus sativus longipinnatus (see Daikon).

Rapid Plant Bug (see Adelphocoris rapidus).

rapidus, Adelphocoris.

Raspberry, pests of, in Britain, 308, 360;

Nysius ericae on, in Br. Columbia,

154; pests of, in Denmark, 381, 382. Rats, destroying *Diparopsis castanea*, 91. rectangulata, Chloroclystis.

rectilinea, Aegocera. rectirostris, Anthonomus.

Red Arrow, 51.

reflexa, Plecoptera. religiosa, Mantis.

repanda, Mocis; Nadiasa (Taragama).
Resin, doubtful value of spray containing,

against Rhoecocoris sulciventris, **268.** Resin Midge (see Retinodiplosis).

Retama monosperma, Lasiocampid on, in Spain, 170.

Reticulitermes flavipes, experiments with laminated paper wrappings and, 272.

Retinodiplosis, bionomics of, on Pinus ponderosa in California, 398.

retusus, Gnathotrichus.

Reviews:— Bawden (F. C.), Plant Viruses and Virus Diseases, 192; da Costa Lima (A.), Lepidoptera of Brazil, 401; García Mercet (R.), Parasites of injurious insects, 357; Gough (H. C.), A Review of the Literature on Soil Insecticides, 393; Hinton (H. E.), A Monograph of the Beetles associated with stored Products, Vol. I., 330; Lepesme (P.), Les Coléoptères des denrées alimentaires et des produits

Mallis industriels entreposés, 403; (A.), Handbook of Pest Control, 318; Paillot (A.), L'Abeille. Anatomie, maladies, ennemis, 328; Risbec (J.), Observations sur les insectes des plantations en Nouvelle-Calédonie, 161; Ruiz Castro (A.), The Insect Fauna of Grape-vines in Spain, 298; Schneider-Orelli (O.), A practical Course on Entomology, 379; Smith (C. F.), The Aphidiinae of North America, 96; Thompson (W. R.), A Catalogue of the Parasites and Predators of Insect Pests, 128; Society of American Foresters, Important Tree Pests of the Northeast, 31.

Rhabdopterus picipes, bionomics and control of, on blueberry and cranberry in

U.S.A., 402.

Rhaconotus, parasite of Pempherulus affinis in India, 21.

Rhagoletis cingulata, sprays against, on cherry in Washington, 23.

Rhagoletis fausta, sprays against, on

cherry in Washington, 24.

Rhagoletis pomonella, in Nova Scotia, 144; in U.S.A., 3, 4, 42, 150, 191, 222, 401; on apple, 3, 42, 144, 150, 191, 222; alternative host-fruits of apple race of, 222; on blueberry, 4, 144, 401; measures against, 3, 4, 150, 191, 222; use of Drosophila for testing insecticides against, 144; methods of timing seasonal cycles of, 42.

rhamni, Aphis.

Rhamnus cathartica, winter food-plant of Aphis rhamni in Britain, 358.

Rhaphidopalpa (see Aulacophora).

Rhenea mediata, on Trichilia emetica in Nyasaland, 86.

Rhigopsidius tucumanus, on potato in Argentina, 4.

Rhizobius ventralis, predacious on Saissetia nigra in California, 374.

Rhizoglyphus echinopus (hyacinthi), intercepted in Sweden, 35; in Washington, 24; fumigation of bulbs against, 24, 35.

Rhizomyrma (see Acropyga).

Rhizopertha dominica (in stored cereals), in Australia, 129, 318, 319, 320, 372; in India, 165; in Oklahoma, 69, 70; bionomics of, 320; measures and experiments against, 69, 70, 165, 318, 319, 372; experimentally infesting dehydrated carrot and potato, 102.

Rhodesia, Northern, termites in, 247.
Rhodesia, Southern, cotton pests in, 87;
miscellaneous pests in, 126, 386; Coleoptera in timber in, 84; virus diseases of plants in, 85, 126; use of Pentatomid as food in, 327; plant quarantines in, 286; regulations against importation of dahlias into Nyasaland from, 85.

rhodesiaensis, Bemisia.

Rhodnius prolixus, tests of inert dusts on, 188.

Rhododendron, pests of, 118.

Rhoecocoris sulciventris, bionomics and control of, on Citrus in N. S. Wales, 267. Rhopalicus azureus, hosts of, in Finland, 12.

Rhopalicus suspensus, hosts of, in Finland, 12.

Rhopalocampta forestan, on Canavalia ensiformis in Nyasaland, 86.

Rhopalosiphum nymphaeae, food-plants of, in Hawaii and Fiji, 251, 290;

Syrphid predacious on, 290.

Rhopalosiphum prunifoliae, natural enemies of, on cereals in Britain, 393, 394; effect of DDT on, on apple in Indiana, 218.

Rhopalosiphum pseudobrassicae, on cruci-

fers in Hawaii, 251.

Rhopobota naevana, sprays against, on cranberry in U.S.A., 23.

Rhoptrocerus xylophagorum, hosts of, in Finland, 12.

Rhubarb, Trachea atriplicis on, in Denmark, 381.

Rhyncaphytoptus platani, bionomics of, on Platanus orientalis in California, 352.

Rhynchagrotis anchocelioides, measures against, on blueberry in U.S.A., 402. Rhynchites, spray against, on fruit trees

in France, 331.

Rhyncolus culinaris, humidity requirements of, in timber in Germany, 18.

Rhyncolus ulkei, in timber in U.S.A., 176. Rhyssa persuasoria, establishment of, against Sirex noctilio in New Zealand, 269.

Ribes (see Currant and Gooseberry).
Ribes alpinum, Thomasiniana on,
Germany, 308.

ribesii, Nematus (Pteronidea, Pteronus).

ribis, Capitophorus.

Rice, distribution of Solubea spp. on, in America, 147; other pests of, in Brazil, 4, 93; pests of, in Fiji, 290; pests of, in Mauritius, 371.

Rice (Stored), Coleoptera in, 103, 236; sorption of fumigants by, 236.

Rice Weevil (see Calandra oryzae).

Richardia, Corythaica monacha on, in Brazil, 263.

Ricin, 106.

Ricinine, and related compounds, toxicity of, to Cydia pomonella, 106.

Ricinus communis (see Castor).

rileyi, Pyroderces.

riparius, Hypnoidus (Cryptohypnus).

roboraria, Boarmia.

Rodolia cardinalis (predacious on Icerya spp.) suggested introduction of, into

Belgian Congo, 365; establishment of, in Ecuador, 96; utilisation of, in France, 357; establishment and bionomics of, in India, 385; utilisation of, in Porto Rico, 110.

Rodolia fumida (roseipennis), in Ceylon, 385; bionomics of, in India, 385.

Rogas plecopterae, sp. n., bionomics of, parasitising Plecoptera reflexa in India, 390.

Rogas rufocoxalis, parasite of Plusia brassicae in Arizona, 112.

Rogas testaceus, bionomics of, in India, 389, 390.

rosa, Ceratitis (Pterandrus). rosaceana, Tortrix (Cacoecia).

rosae, Athalia; Eurytoma; Macrosiphum; Psila.

vosana, Tortrix (Cacoecia).

rosaphidis, Aphidius.

Rose, thrips on, in S. Australia, 129; Aonidiella aurantii on, in Brazil, 204, 205; pests of, in Fiji, 290; Myzus circumflexus on, in Hawaii, 251; pests of, in India, 385, 393; pests of, in U.S.A., 96, 107; experiment with Thomasiniana theobaldi and, 308.

Rose Chafer (see Macrodactylus subspinosus).

roseipennis, Rodolia (see R. fumida).

roseus, Anuraphis. roseveari, Aprostocetus.

Rotenone, 99; against Aphids, 51, 101, 214, 260, 336; against Coccids, 349, 350, 374; against Jassids, 28, 104; against other Hemiptera, 51, 63, 64, 353; against Coleoptera, 79, 100, 101, 176, 260; against Lepidoptera, 51, 119, 197, 212, 223, 262, 338; against Rhagoletis pomonella, 191; against thrips, 104; testing and modes of action of, on insects, 189, 197, 340; in dusts, 63, 64, 79, 100, 101, 119, 176, 191, 212, 214, 223, 260, 262, 336, 338; in sprays, 28, 51, 104, 223, 349, 350, 353, 374; formulae containing, 51, 63, 79, 101, 104, 212, 260, 336, 338; and calcium arsenate, 336; and other derris extractives, 212, 350; and mannitan monolaurate, 51; and nicotine, 260, 336, 338; addition of, to oil emulsions, 349, 350; and pyrethrum, 28, 51, 104, 119; and sulphur, 63, 260, 336; "activated" form of, 336; proprietary insecticides containing, 51, 212. Cubé, Derris, Lonchocarpus and Tephrosia.)

rothsteini, Monomorium.
rouxi, Habrolepis.
rubecula, A-panteles.
rubens, Ceroplastes.
rubeo, Cremastus.
ruber, Deraeocoris (Capsus).

rubi, Anthonomus.
rubicola, Agrilus.
rubidum, Sphaeroderma.
rubra, Leptura; Odoniella.
rubriceps, Metoponia.
rubrocinctus, Selenothrips.
rufa, Formica; Solenopsis geminata.
ruficeps, Stolotermes.
ruficollis, Dysdercus; Ephialtes.
ruficornis, Cerotoma.
ruficrus, Apanteles.
rufimanus, Bruchus.

rufimanus, Bruchus. rufipes, Eurytoma; Harpalus; Hylastes (Hylurgopinus); Necrobia.

rufiscutellaris, Glypta. rufocoxalis, Rogas.

rufo-pratensis, Formica rufa. rufovillosum, Xestobium. rugicollis, Plesiocoris.

rugifrons, Neodiprion. rugosus, Carabus.

Rumania, *Phytodecta fornicata* on lucerne in, **121.**

Rumex, Capnodis tenebricosa on, in Palestine, 75.

Rumex obtusifolius, Macrosiphum solanifolii on, in Britain, 358; experiments with Aphis viburni and, 383.

rumicis, Aphis. ruralis, Voria. rusci, Ceroplastes.

Russia (U.S.S.R.), Cleonus punctiventris on beet in, 166, 302; pests and virus disease of cereals in, 183, 302, 307; distribution of Melolontha spp. in, 159; miscellaneous pests in, 125, 304–306; Glycyphagus destructor in stored grain in, 330; insects pollinating Taraxacum hok-saghyz in, 245; natural enemies and biological control of insects in, 302, 303, 304, 306. (See Siberia and Asia, Central.)

Russian Thistle (see Salsola pestifer). rusti, Coccophagus.

rustica, Dexia; Tachina.

rutgersi, Acropyga (Rhizomyrma).

Rye, Hylemyia coarctata on, in Britain, 297; pests of, in Denmark, 382; Zabrus tenebrioides on, in Hungary, 11; (rolled), as bait for cutworms, 140.

Rye-grass (see Lolium).

S.

saccharalis, Diatraea. sacchari, Aphis; Paratetranychus (Tetranychus). saccharicola, Melanaspis. saccharina, Eldana; Lepisma. sacchariphagus, Proceras. 488 INDEX.

Sagaritis provancheri, parasite of Heliothis virescens in Virginia, 281; list of hosts of, 281.

Sagaritis websteri, parasite of Plusia brassicae in Arizona, 112.

sagittata, Lucumaphila.

Sahlbergella singularis, on cacao in W. Africa, 65, 66, 195, 365, 366; bionomics and attempted biological control of, 65, 195; insecticides against, 66.

Sahlbergella theobroma (see Distantiella).

sahlbergellae, Euphorus.

St. Augustine Grass (see Stenotaphrum secundatum).

St. Croix, Heliothis virescens on cotton in, **278**.

St. John's Wort (see Hypericum per-

foratum).

Saissetia coffeae (hemisphaerica), in Palestine, 326; on Cajanus cajan in Porto Rico, 243; natural enemies of, 243, 326.

Saissetia nigra, bionomics and control of, in California, 373, 374; food-plants, natural enemies and distribution of,

373, 374.

Saissetia oleae, in Australia, 322; California, 61, 259, 374; in Palestine, 326; parasites and biological control of, 259, 322, 326, 374; biological control of mealybug affecting rearing of, 61; measures against, on Citrus,

saissetiae, Anysis; Coccophagus.

salaminia, Eumaenas (Ophideres).

salicis, Stilpnotia.

Salicylaldehyde, not toxic to insects, 198. Salix (see Willow).

Salix caprea, Geometrid on, in Britain, 360.

Salix vitellina, Cavariella aegopodii not transmitting carrot virus to, in Victoria, 292.

sallei, Lygus.

Sallow (see Salix caprea).

Salsola pestifer, Eutettix tenellus on, in U.S.A., 281.

Salt (Sodium Chloride), ineffective against Bruchid in cowpeas, 327; solutions containing, against Tineola biselliella, 266, 267; increasing toxicity of derris dust to Musca domestica, 78; increasing efficiency of cutworm baits, 140.

saltator, Halticus.

sambuci, Asterolecanium pustulans (see A. pustulans).

José Scale (see Quadraspidiotus perniciosus).

sanctae-helenae, Heteronychus.

sanctus, Sympherobius.

Sand, use of, to protect stored products from pests, 131, 300.

sanguinarius, Dysdercus.

sanguinea, Cycloneda. sanguineus, Helopeltis.

sanguinipes, Pimpla (Ephialtes).

sanscrita, Halyzia.

Santo Domingo, new Pentatomid on rice in, 147.

Saperda candida, methods of timing seasonal cycles of, on apple in Maine,

Sapindus, Serrodes partita on, in India,

Saponins, toxicity of, to fruit-flies, 397. Sapote, Black (see Diospyros ebenaster).

saratogensis, Aphrophora. Sarcophaga, parasite of Helopeltis orophila

in Belgian Congo, 37.

Sarcophaga caridei, parasite of Schistocerca paranensis in Argentina, 401.

Sarcophaga misera, parasite of Helopeltis orophila in Belgian Congo, 37.

Sarcophaga tuberosa, parasite of Anabrus simplex in Nevada, 146; Bembicid destroying, 146.

Sargentia greggii, Anastrepha ludens on,

in Mexico, 52, 395.

Saskatchewan, Cephus cinctus in, 98; grasshoppers in, 98, 144, 153; Heliothis ononis on flax in, 144.

Sassafras Oil, in baits for Polychrosis viteana, 27.

Satin Moth (see Stilpnotia salicis).

saucia, Peridroma. sauteri, Dinarmus.

Saw-toothed Grain Beetle (see Oryzaephilus surinamensis).

Sawdust in bait, 145, 154, 179, 183, 259; protecting stored potatoes from Gnorimoschema operculella, 300.

Sawfly, Apple (see Hoplocampa testudinea).

savi, Chlorochroa.

scabra, Hydnocera.

scabricollis, Peranabrus.

scalaris, Megaselia.

Scale, Beech (see Cryptococcus fagi).

Scale, Black (see Saissetia oleae).

Scale, California Red (see Aonidiella aurantii).

Scale, Citricola (see Coccus pseudomagnoliarum).

Scale, Florida Red (see Chrysomphalus ficus).

Scale, Olive (see Parlatoria oleae).

Scale, San José (see Quadraspidiotus perniciosus).

Scale, Scurfy (see Chionaspis furfura).

Scale, Sycamore (see Stomacoccus platani). Scale, White Citrus (see Unaspis citri).

Scale, White Wax (see Ceroplastes destruc-

Scale Insects, natural enemies of, 59, 61, 79, 86, 92, 95, 96, 109, 110, 132, 133, 208, 209, 214, 227, 248, 259, 313, 322,

325, 326, 328, 364, 365, 368, 374, 385, 389; ants associated with, 65, 66, 147, 214, 365; and plant diseases, 66, 147; comparison of injury to apple by Aphids and, 332; classification and new species of, 92, 166, 328.

Scaptocoris terginus, food-plants of, in

Argentina and Brazil, 4.

Scaptomyzella (Scaptomyza) incana, cauliflower in Denmark, 381.

Scenopinus fenestralis, predacious Tineola biselliella in U.S.A., 351.

Schedius kuvanae (see Ooencyrtus).

Schinus molle, Anoplognathus on, N. S. Wales, 162; Capnodis cariosa on, in Palestine, 75.

Schinus terebinthefolius, Myllocerus laeti-

virens on, in India, 388.

Schistocerca americana, attacking cotton

in Ecuador, 5.

Schistocerca gregaria, activities and control of, in Arabia, 228; in Belgian Congo, **366:** Gryllotalpa feeding on eggs of, in cages in Egypt, 226; in India, 136, 388; in Nigeria, 370; in Tanganyika, 85; migrations of, 85, 366, 370; factors affecting development of, 366, 386, 388; absorption of water by egg of, 386; methods of distinguishing phases and phase transformation in, 136, 201,

Schistocerca obscura, bionomics of, in

Oklahoma, 237.

Schistocerca paranensis, in Argentina, 66, 67, 120, 401; in Ecuador, 5; resistance of varieties of maize to, 66, 67; oviposition habits of, 120; parasite of, 401. schlechtendali, Phyllocoptes.

schwartzi, Anuraphis.

Sciara, aerosol against, in mushroom houses in Quebec, 142.

Scirtetica marmorata, Harr., 146.

Scirtothrips aurantii, on Citrus in S. Rhodesia, 126.

Scirtothrips citri, measures against, on Citrus in California, 97, 98.

scitula, Eublemma.

Scolytus multistriatus, factors affecting infestation and spread of Ceratostomella ulmi by, on elm in U.S.A., 43, 173, 313.

scrophulariae, Anthrenus.

scrupulosa, Teleonemia.

Scurfy Scale (see Chionaspis furfura).

scutellaris, Coccophagus.

scutellata, Sturmia.

scutellatus, Gonipterus.

Scutellista cyanea, parasite of Saissetia nigra in California and Hawaii, 374; hosts of, in Palestine, 326.

Scutigerella immaculata, dichlorethylether emulsion against, on vegetables in California, 338.

Scymnus, predacious on Sipha flava in Argentina, 133; predacious on Saissetia nigra in California, 374.

Scyphophorus acupunctatus, bionomics of, on Agave fourcroides in Mexico, 131.

Seed-corn Maggot (see Hylemyia cilicrura). segetum, Agrotis (Euxoa).

Seius, sulphur reducing control of Paratetranychus pilosus by, in Ohio, 208.

Selenaspidus silvaticus, parasite of, on Trichilia emetica in S. Africa, 107.

Selenothrips rubrocinctus (on cacao), in Belgian Congo, 366; not transmitting virus disease in Gold Coast, 66.

Semesan, 178.

semialbiclavus, Hemiptarsenus.

semiflavidus, Anastatus.

semiflaviventris, Phanuropsis.

semipunctata, Phoracantha.

semirugosus, Allodorus.

senatoria, Anisota.

senilis, auct., Ceromasia (see Lydella stabulans grisescens).

senior, Mesotrichia (Xylocopa). sepia, Hylemyia (Chortophila). septemfasciata, Nomadacris.

septempunctata, Coccinella.

seriatus, Psallus.

Serica brunnea, in grass in Britain, 155.

sericea, Prasoidea (Odontionopa). sericeofasciata, Pachypasa.

Sericesthis pruinosa, damaging plant roots in N. S. Wales, 161.

sericeum, Trombidium (see Eutrombidium trigonum).

serius, Eretmocerus.

serpentina, Anastrepha.

serricorne, Lasioderma.

serripes, Myodochus.

Servodes partita (inara), bionomics of, in India and S. Rhodesia, 386.

sertifer, Neodiprion. servus, Euschistus.

Sesame Oil, as an activator for aerosols, 99, 142.

Sesamia epunctifera, bionomics of,

wheat in Nyasaland, 86. Sesamia fuscifrontia, bionomics of, on

wheat in Nyasaland, 86. Sesamia vuteria, on maize and rice in

Mauritius, 371.

sexannulatus, Cryptus. sexmaculata, Chilomenes.

sexmaculatus, Tetranychus.

sexta, Protoparce.

Sheepskins, insects infesting, in Massachusetts, 82.

sheldoni, Eriophyes. Shrew (see Sorex).

Siam, Pempherulus affinis in, 20.

Siberia, Enarmonia diniana desertana and its parasites on larch in, 400-401.

Sida spp., insects on, in S. America, 204, 263; parasitism of Pempherulus affinis on, in India, 21.

signata, Cryptognatha. signataria, Melanolophia.

signatus, Anthonomus.

Silica Dusts, tests with, against pests of stored products, 31, 231, 372; use of, in testing contact insecticides, 197; danger of, to man, 372; proprietary forms of, 231, 372.

Silk, Tineola biselliella not developing on,

351.

Silkworm (see Bombyx mori).

Silvanus surinamensis (see Oryzaephilus). silvaticus, Selenaspidus (Aspidiotus). silvestris, Phorocera (Parasetigena).

similaris, Epitrix.

similis, Collops; Diprion.

simonyi, Lygus.

simplex, Anabrus; Apion; Cuspicona; Taeniothrips.

Sinea diadema, predacious on noxious insects in Utah, 206.

singularis, Otiorrhynchus; Sahlbergella. sinicus, Adoretus.

Sinocalamus (see Bamboos).

Sinoxylon, bionomics and control of, in timber in India, 71, 72, 371.

Sinoxylon anale, in timber in India, 371. Sinoxylon brazzai, on cassava in Belgian Congo, **367.**

Sinoxylon conigerum, on Eucalyptus and Terminalia in Mauritius, 371.

Sinoxylon sudanicum, natural enemies of, in India, 21, 164; bionomics of, on cotton, 164.

Sipha flava, bionomics of, on sugar-cane in Argentina and Porto Rico, 133, 243; bibliography of, 133.

Sirex, on Abies in Switzerland, 194.

Sirex noctilio, biological control of, on Pinus radiata in New Zealand, 269.

siro, auct., Tyroglyphus (see Tyrolichus casei).

Sitodiplosis mosellana, infestation by, as basis for forecasting wheat yield in Britain, 293.

Sitodrepa panicea, products infested by, in California and Germany, 102, 230.

Sitona (on lucerne), apparatus for collecting, in Czechoslovakia, 252; in Russia,

Sitona cylindricollis, Bufo cognatus destroying, on Melilotus in N. Dakota, 96. Sitona flavescens, fungus infesting, in

Britain, 232.

Sitona lineatus, on leguminous crops in Washington, 22, 24, 100, 101; overwintering of, 24; dusts against, 22; toxicity of gammexane to, 257.

Sitones (see Sitona).

Sitophilus (see Calandra).

Sitotroga cerealella, measures and experiments against, in stored cereals in S. America, 131, 375; use of eggs of, for rearing Trichogramma spp., 125, 370.

siva, Nadiasa (Taragama). sjaelandicus, Corymbites. slossonae, Agrothereutes.

smaragdina, Oecophylla. smaragdula, Nezara viridula.

Smartweed (see Polygonum).

smithi, Clemora (Phytalus); Diversinervus.

Smodicum cucujiforme, in timber in U.S.A., 176.

Smynthurus viridis, study of growth of population of, on clover in W. Australia, 129.

Snapdragon (see Antirrhinum).

Snowberry (see Symphoricarpus) Soap, spraying with, 38, 97, 268, 317; as an emulsifier, 227, 304; other uses of, in mixed sprays, 28, 38, 45, 51, 97, 124, 125, 168, 172, 185, 202, 264, 268, 293, 307, 317, 334, 364, 365, 380; formulae containing, 45, 51, 97, 168, 185, 187, 227, 268, 293, 304, 307, 317, 364, 380; types of, 28, 45, 51, 168, 202,

Soda, Caustic (see Sodium Hydroxide).

Soda Ash (see Sodium Carbonate).

Sodium Aluminium Sulphate, in lead arsenate sprays, 323.

Sodium Antimonyl Hydroxyacetate, and sugar, in sprays against Thrips tabaci, 52, 180.

Sodium Antimonyl Lactophenolate, and sugar, in sprays against Thrips tabaci, 52.

Sodium Arsenate, in bait for fruit-flies, 134.

Sodium Arsenite, in baits, 60, 134, 140, 145, 201, 270; in bait-spray for cotton stainers, 88; and sulphur, dusting with, against Lygus oblineatus, 63; spraying with, against Nomadacris and Laphygma, 87, 246; aeroplane spraying with, against Eurygaster integriceps, 303; as preservative for timber against termites, 247: mode of action of, on Periplaneta americana, 64; question of risk of, to domestic animals, 60, 201.

Sodium Carbonate, uses of, in sprays, 108, 307, 365; in preparation of pitch soap, 202.

Sodium Chloride (see Salt).

Sodium Cyanide, for generating hydrocyanic acid gas (q.v.), 127, 130.

Sodium Dinitro-o-cresylate, in dormant sprays against orchard pests, 58, 153, 179; and oil emulsion, 153; less ovicidal than dinitro-o-cresol, 15.

Sodium Fluoaluminate (see Cryolite).

Sodium Fluoride, against Bruchid

cowpeas, 327; materials treated with, against Anthrenus, 73; treatment of timber with, against pests, 84, 299; in baits and bait-sprays, 134, 142, potato injured by sprays con-343; taining, 185.

Sodium Fluosilicate, in baits, 59, 134, 140, 179, 201; in bait-sprays, 17, 134, 162, 256, 260; plants injured by dusts and sprays of, 105, 185; treatment of stored seeds with, against pests, 327, 328; question of danger of, to mammals and birds, 56, 59, 60, 328.

Sodium Hydroxide (Caustic Soda), in

sprays, 268, 272.

Sodium Lauryl Sulphate, in derris dusts and sprays, 50, 139.

Sodium Nitrite, aeroplane spraying with, against Eurygaster integriceps, 303, 304. Sodium Oleate, as a wetting agent for

sprays, 169.

Sodium Sulphoricinate, as a wetting agent for sprays, 169.

Sodium Tetradecyl Sulphate, 51.

Sogata furcifera, on rice in Fiji, 290.

Soil, studies on Arthropod fauna of, 129, 362; effects of type of, on insect pests, 265, 266, 369; effect of termites on, 329; effect of lead arsenate in, on plants, 63.

solana, Empoasca.

solani, Macrosiphum; Phenacoccus: Phytomyza (Liriomyza).

solanifolii, Macrosiphum.

Solanum, lists of species of, susceptible or resistant to Leptinotarsa decemlineata,

Solanum auriculatum, Cuspicona simplex

on, in New Zealand, 93.

Solanum demissum, experiments with hybrids of potato and, against Leptinotarsa decemlineata, 159.

Solanum gilo, insects on, in Brazil, 264,

Solanum melongena (see Egg-plant).

Solanum nigrum, Gnorimoschema oper-culella on, in N. S. Wales, 185.

Solanum nodiflorum, Phyllocoptes destructor on, in Hawaii, 250.

Solanum pterocaulon, Gnorimoschema operculella on, in N. S. Wales, 185.

Solanum sodomeum, Cuspicona simplex on,

in New Zealand, 93.

Solanum torvum, Prodenia litura on, in Fiji, 201; Corythaica passiflorae on, in Costa Rica, 263.

Solanum tuberosum (see Potato).

Solanum verbascifolium, 196.

Solar Heat, against Platyedra gossypiella in cottonseed, 375.

Solenopsis fugax, question of use of, against harmful ants, 16.

Solenopsis geminata, in houses in Germany, 16; oil emulsion against, in Mauritius, 371; suggested encouragement of, against Anthonomus grandis in Texas, 122.

Solenopsis geminata var. diabola, destroying insects in Arizona, 177.

Solenopsis geminata rufa, destroying tomatos in Hawaii, 250.

Solidago, pests of, in U.S.A., 259.

solstitiale, Amphimallon (Amphimallus).

Solubea, revision of, 147.

Solubea insularis, on rice in Mexico, 147. Solubea ornata, sp. n., on rice in W. Indies,

Solubea poecila, on rice in S. America and Trinidad, 147; systematic position of,

Solubea pugnax, on cereals and grasses in U.S.A., 147.

Somalia, organisation of locust control in,

sonchi, Macrosiphum.

Sonchus oleraceus, Aphis sacchari on, in Hawaii, 251.

Sooty Mould, associated with Coccids, 59, 79, 171, 228; associated with Phorodon humuli, 317.

sophiae, Colaphellus (Colaphus).

Sorbus aucuparia, Pristiphora geniculata on, in Quebec, 142.

sordidus, Aphanus; Cosmopolites. Sorex, destroying Chlorion laeviventris in U.S.A., **146.**

Sorghum, pests of, in E. Africa, 85, 88, pests of, in U.S.A., 102, 191; dinitro compound not injurious to, 54. Sorghum (Stored), Tribolium destroying

eggs of Corcyra cephalonica in, 393. Soy Bean, Pyrausta nubilalis on, in Ontario, 139.

Soybean Flour, in baits, 140, 179; as an adhesive for sprays, 43, 51, 272.

Soybean Oil, in sprays, 2, 149, 273; causing loss of rotenone in stored derris dusts, 152.

spadix, Codiosoma (Pselactus).

Spain, pests of beet in, 120, 166; Dociostaurus maroccanus in, 166, 167; Leptinotarsa decemlineata in, 165, 166; miscellaneous pests in, 167, 168, 170, 310; orchard pests in, 125, 169; vine pests in, 298; beneficial insects and biological control in, 125, 170, 310, 357.

Spalangia, parasite of Dacus in Fiji, 161. Spalangia drosophilae, hosts and rearing

Sparaxis, Aphid transmitting virus to Tritonia from, in U.S.A., 2.

Sparganothis pilleriana, on vines Spain, 298.

Sparganothis sulfureana, on strawberry in Ohio, **263.**

Sparrows, destroying Hylemyia floralis, 121.

Spathius, parasite of Sinoxylon in India, 164.

Spathius critolaus, hosts and biology of, in India, 21, 164.

Spathius exarator, parasite of Anobium punctatum in Germany, 19.

Spathius labdacus, 164; parasite of Pempherulus affinis in India, 21.

Spatulifimbria castaneiceps, on tea in Ceylon, 300.

speciosa, Diabrotica.

spectabilis, Metapelma.

Sphaeroderma rubidum, bionomics and control of, on artichoke in Spain, 168.

Sphaerostilbe, infesting Unaspis citri in N. S. Wales, 248.

Sphaerotheca humuli, on hops in New York, 317.

Sphingomorpha chlorea, piercing fruits in India and S. Rhodesia, 386.

Sphinz pinastri, on pine in Poland, 6. Spicaria javanica, utilisation of, against

Icerya purchasi in W. Indies, 110. spicata, Phytomyza.

spiculatus, Ergates.

Spiders, destroying noxious insects, 28, 30, 112, 298, 320.

Spilochalcis spp., parasites of Coleophora laricella in Ontario, 143.

Spilonota ocellana (on apple), insecticides against, in New York, 179; in Switzerland, 394.

Spinach, Hylemyia cilicrura on, in California, 338.

spinarum, Athalia (see A. rosae).

spinolae, Lygus.

spinosula, Guerinia (see Tachina rustica). spissicornis, Coleophora.

splendens, Chrysoplatycerus.

Spondias, Anastrepha spp. on, in Mexico, 52, 53, 397.

Spondias mombin, form of Anastrepha mombin praeoptans on, in Mexico, 52; Helopeltis sanguineus on, in Nigeria, 383.

Spondylis buprestoides, humidity requirements of, in timber in Germany, 18.

Spray Kast, 149.

Sprays, methods and value of atomisation of, on crops, 15, 254; tracheal system of insects in relation to, 252; study of deposition and retention of, 116, 252; particle size specifications for powders for, 42.

Spruce, 194; pests of, in Canada, 143, 155, 180, 223, 313, 398; pests of, in Czechoslovakia, 8; pests of, in Finland, 7, 11, 12; pests of, in Germany, 14; Dasychira pudibunda on, in Sweden, 19; pests of, in U.S.A., 47, 313, 379,

398; methyl bromide causing injury to, 50.

Spruce, Colorado (see Picea pungens).

Spruce, Norway (see Picea abies).

Spruce Budworm (see Harmologa fumiferana).

spurius, Apanteles. sputator, Agriotes.

Squash, Aphis maidis on, in Hawaii, 251; pests of, in U.S.A., 21, 179, 212; as trap crop for Anasa tristis, 21.

Squash Bug (see Anasa tristis). stabulans, Lydella; Muscina.

stanleyi, Metaphycus.

Staphylococcus flaccidifex, 81.

Starch, relation of *Dinoderus minutus* to, in bamboo, **312.**

Stathmopoda, predacious on Unaspis citri in N. S. Wales, 248.

Stathmopoda melanochra, in Australia, 385; predacious on Icerya purchasi in India, 385.

Statistica Methods, application of, to entomological problems, 282.

Stearic Acid, in fixed nicotine sprays, 273. Steganoptycha (see Enarmonia).

Stegobium (see Sitodrepa).

steinwedeni, Acaphylla.

stellata, Lyda.

Stenocarus fuliginosus, outbreaks and control of, on poppy in Hungary, 122.

Stenodiplosis geniculati, 170.

Stenodiplosis stenotaphri, sp. n., on Stenotaphrum dimidiatum in Mauritius, 74.
Stenommatus musae, distribution of, on

banana, 202.

stenotaphri, Stenodiplosis.

Stenotaphrum dimidiatum, new Cecidomyiid on, in Mauritius, 74.

Stenotaphrum secundatum, experiment with Crambus haytiellus and, in Florida, 115.

Stephanoderes, on cassava in Belgian Congo, 367.

Stephanoderes hampei, bionomics of, in Brazil, 203.

Sterculia spp., Mesohomotoma tessmanni on, in Gold Coast, 66; Dysdercus spp. on, in Nyasaland, 91; Pseudococcus exitiabilis on, in Fr. Togoland, 196.

sticticalis, Loxostege. sticticosta, Metarbela.

Stictiella pulla, destroying Sarcophaga tuberosa in Nevada, 146.

Stictocephala inermis, bionomics and control of, on pear in Washington, 285.

Stictococcus, experiment with ant and, on cacao in Gold Coast, 65; list of species of, 166.

Stictococcus formicarius, ant associated with, on cacao in Belgian Congo, 365.

Stictococcus guineensis, sp. n., on coffee in Spanish Guinea, 166.

stigma, Chilocorus.

stigmatera, Agathis (Bassus).

Stilobezzia, on cacao in Trinidad, 78.

Stilpnotia salicis, bionomics of, on aspen in Sweden, 32.

Stizoides unicinctus, relation of, to Anabrus simplex in U.S.A., 146.

Stizolobium aterrimum, 383.

Stolotermes ruficeps, characters of, in New Zealand, 234.

Stomacoccus capsulatus, bionomics of, on Platanus wrightii in Arizona, 353.

Stomacoccus platani, bionomics and control of, on Platanus spp. in U.S.A. and Mexico, 352.

Strangalia obliterata, in fire-killed Douglas fir in U.S.A., 40.

Strategus julianus, bionomics of, on Agave

fourcroides in Mexico, 131.

Strawberry, Capitophorus fragaefolii on, in Argentina, 4; pests of, in Britain, 117, 181, 358, 359, 360; pests of, in Denmark, 381, 382; pests of, in N.S. Wales, 161, 162; mite on, in Switzerland, 117; pests of, in U.S.A., 46, 96, 117, 191, 262, 263, 379; spray causing injury to, 379.

Strawberry Leafroller (see Ancylis comp-

Streptanthera, Aphid transmitting virus to Tritonia from, in U.S.A., 2.

striata, Anastrepha.

striatellus, Delphacodes (Delphax). strigata, Agromyza (Liriomyza). strobilella, Cydia (Laspeyresia).

Stromatium fulvum (unicolor), bionomics of, in timber and furniture in Spain, 167, 168.

Strophosomus lateralis, on heather Britain, 233.

Strumeta (see Dacus).

Sturmia, utilisation of, against Gilpinia hercyniae in Br. Columbia, 155; parasite of Anaphaeis aurota in Belgian Congo, 368; parasite of Laphygma exigua in S. Rhodesia, 126.

Sturmia scutellata, introduced parasite of Lymantria dispar in U.S.A., 238.

stylifer, Lasiohelea.

subcoleoptrata, Alophora.

subcrinita, Epitrix.

subgranosus, Platypus.

subobscura, Drosophila.

subspinosus, Macrodactylus. Sudan, Anglo-Egyptian, cotton pests in,

Sudan Bollworm (see Diparopsis cast-

anea). sudanicum, Sinoxylon.

Sugar, in baits, 27, 179, 216, 340, 396; in bait-sprays for fruit-flies, 162, 190, 256; in sprays and bait-sprays against other insects, 17, 52, 88, 97, 180, 185, 260; used for feeding insects, 17, 144, 215, 231, 390.

Sugar-cane, Sipha flava on, in Argentina, 133; Melanaspis saccharicola on, in Brazil, 205; pests of, in India, 390, 391, 406; Pyrausta coclesalis on, in Indo-China and Java, 388; pests of, in Mauritius, 370; Heteronychus sanctae-helenae on, in N. S. Wales, 161; pests of, in Porto Rico, 26, 97, 243, 312; Cochliotis melolonthoides on, in Tanganyika, 85; pests of, in U.S.A., 25, 116, 177, 260, 261.

Sugar-cane Aphis, Yellow (see Sipha

Sugar-cane Moth Borer (see Diatraea saccharalis).

sulcatus, Gnathotrichus; Otiorrhynchus,

sulciventris, Rhoecocoris.

sulfureana, Sparganothis.

Sulphite Lye, benzene hexachloride dis-

persed in water by, 257.

Sulphur, against Capsids and Pentatomids, 29, 53, 54, 260, 261; against Coccus pseudomagnoliarum, 98; against Paratrioza cockerelli, 30, 216, 217; against mites, 148, 172, 219, 285, 286, 299, 307; increasing infestation of apple by Paratetranychus pilosus, 4, 208; against Scirtothrips citri, 98; not protecting stored potatoes and maize from pests, 31, 327; dusting with, 4, 29, 30, 63, 98, 216, 217, 219, 260, 261, 299, 307, 316; in mixed dusts, 3, 43, 49, 59, 63, 64, 78, 101, 112, 150, 168, 211, 222, 239, 240, 260, 261, 274, 316, 335, 336; increasing toxicity of derris dust to Musca domestica, 78; protecting rotenone in dusts containing nicotine, 240; as a fumigant (see Sulphur Dioxide); spraying with, 4, 30, 39, 172, 208, 216, 217, 285, 286, 299; in mixed sprays, 39, 43, 98, 150, 172, 212, 227, 307, 317; applied by aircraft, 307; forms of, 78, 208, 261, 274, 327; formulae containing, 3, 30, 39, 43, 48, 59, 63, 64, 78, 150, 168, 172, 217, 219, 222, 227, 239, 260, 274, 307, 316, 317, 336; and injury to plants, 78, 105.

Sulphur Dioxide, fumigation experiments

with, 26, 27, 231.

Sulphuric Acid, and sodium cyanide, for generating hydrocyanic acid gas (q.v.),

Sunflower, Thrips angusticeps on, in Denmark, 381.

Sunn Hemp (see Crotalaria juncea).

Supella supellectilium, value of laminated paper wrapping against, 272.

superbus, Adelphocoris. superstitiosus, Dysdercus.

surinamensis, Oryzaephilus (Silvanus). suspensus, Rhopalicus.

suturalis, Lochmaea.

swainei, Neodiprion.

Sweden, forest pests in, 19, 20, 32; miscellaneous pests in, 35, 36; natural enemies of Dasychira pudibunda in, 20.

Swedes, Aphids and virus disease of, in Australia, 323; Hylemyia floralis on, in Britain, 121; Nysius ericae on, in Br. Columbia, 154; pests of, in Denmark, 381.

Sweet Clover (see Melilotus).

Sweet Potato, pests and virus disease of, in E. Africa and Belgian Congo, 329, 330; pests of, in Fiji, 290, 291; Aphids on, in Hawaii, 251; Cylas formicarius on, in Louisiana, 216; pests of, in Mauritius, 371; weevil on, in S. Rhodesia, 126.

Sweet-potato Leaf-roller (see Tricho-

ptilus wahlbergi).

Switzerland, Chermes spp. on Abies spp. in, 194; orchard pests in, 124, 125, 394; Tarsonemus pallidus on strawberry in, 117; vegetable pests in, 125, 157, 288; Phylloxera on vines in, 124.

Sycamore (see Platanus).

Sycamore Scale (see Stomacoccus platani). sycophanta, Calosoma.

sydneyensis, Anarhopus.

Sylepta derogata, on cotton in Belgian Congo, **367.**

Sylvia communis, destroying Nematus

ribesii in Germany, 14.

Sympherobius, factors affecting utilisation and value of, against mealybugs in Palestine, 225, 226.

Sympherobius amicus, 225. Sympherobius sanctus, 225.

Symphoricarpus, damaged by Anabrus simplex in U.S.A., 115.

Symphoricarpus racemosus, Clysiana ambiguella breeding on, in Germany, 13.

Synanthedon (see Aegeria).

Syntomopus americanus, parasite of Agro-

myza virens in California, 104. Syntomosphyrum indicum, liberation of, against fruit-flies in Fiji, 291.

syrphiphagum, Pachyneuron.

Syrphoctonus agilis (see Diplazon).

Syrphus balteatus, predacious on Aphids in Britain, **359.**

Syrphus corollae var. vitiensis (predacious on Rhopalosiphum nymphaeae), parasite of, in Fiji, 290.

Systates, spray against, on Cinchona in Belgian Congo, 38.

T-gas, 231.

tabaci, Bemisia; Faustinus (Collabismodes) (see F. cubae); Thrips.

Tabacol, insecticidal and fungicidal value of, 252.

Tachina larvarum, introduced parasite of Lymantria dispar in U.S.A., 238.

Tachina rustica, parasite of Gilpinia hercyniae in Quebec, 224.

Tachycines asynamorus, records and confusion of, with Diestrammena in N. America, 146.

Tachyporus spp., bionomics of, predacious on Capitophorus fragariae in Britain,

Taeniocampa gracilis, bionomics and control of, on apple in Estonia, 119.

Taeniothrips cardamomi, not transmitting virus disease of cardamom in India, 392. Taeniothrips gladioli (see T. simplex).

Taeniothrips inconsequens, sprays against, on pear in Washington, 285.

Taeniothrips simplex, measures against, on Gladiolus in Quebec, 224.

Taeniothrips xanthius, on greenhouse plants in Hawaii, 250.

Tagetes minuta, not protecting stored maize from Calandra oryzae, 327.

Tahiti, banana pests in, 202; biological control of Cosmopolites sordidus in, 202. Talc, as a carrier for dusts, 22, 24, 45,

49, 78, 79, 112, 178, 180, 212, 239, 262, 293, 334,

Tamarindus indica (Tamarind), Aonidiella

aurantii on, in Brazil, 204. Tamarix, Icerya purchasi on, in France, 357.

tanaceti, Galeruca (Adimonia).

Tanganyika Territory, cotton pests in, 88, 89; miscellaneous pests in, 85, 88, 89, 245; virus disease of sweet potato in, 330.

Tannic Acid, increasing toxicity of derris dust to Musca domestica, 78.

Tar, Water-gas, as barrier against Blissus

leucopterus, 57.

Tar Distillates, in mixture for brushing almond trees against Aglaope, 169; as dormant sprays against orchard and vine pests, 35, 117, 124, 169, 233, 234, 253, 254; ineffective against Paratetranychus pilosus, 233, 234, 254; tests and uses of, in other sprays, 66, 227, 228, 268, 304; soil treatment with, against Contarinia pyrivora, 12; in dusts against pests of vegetables, 276. 304; wheat bags dipped in, against pests, 319; ineffective as repellent against Melolontha, 182; not repellent to honey-bees, 6; characters associated with toxicity of, to insect eggs, 254; effects of, on plants, 66, 276, 304; types of, 35, 117, 124, 304; formulae for, 12, 117, 276, 304; formulae containing, 169, 268; emulsifiers for, 304; and naphthalene, 168; and potassium cyanide, 168; and quicklime, 169.

Taragama (see Nadiasa).

Taraxacum kok-saghyz, insects pollinating, in Russian Union, 245; pests of, in U.S.A., 259.

Targionia jaboticabae, on Myrciaria jaboticaba in Brazil, **205.**

Tarnished Plant Bug (see Lygus oblineatus).

Taro (see Colocasia esculenta).

Tarsonemus bakeri, mite predacious on, in California, 398.

Tarsonemus bancrofti, measures against, on sugar-cane in Florida, 260.

Tarsonemus fragariae (see T. pallidus).

Tarsonemus laticeps, fumigation against, on bulbs in Washington, 24.

Tarsonemus latus, on tomato in Hawaii, **250.**

Tarsonemus pallidus, in Britain, 360; in Holland, 12; in Switzerland, 117; in U.S.A., 43, 263; on strawberry, 117, 263, 360; on other plants, 12, 43; measures and experiments against, 12, 43, 360.

Tartar Emetic, in bait-sprays for fruitflies, 162, 190, 397; in sprays against thrips, 52, 97, 180; resistance of Scirtothrips citri to, 97; and sugar, 52, 97, 162, 180, 185, 190; and molasses, 397; formulae containing, 52, 180, 397; plants not injured by, 105, 185.

Tasmania, Chrysomela hyperici imported into, against Hypericum perforatum,

321.

Taxomyia taxi, on Taxus in Britain, 171.

Taxus, pests of, in Britain, 171.

Taxus cuspidata, reactions of, to methyl bromide, 50.

taxus, Aonidiella.

Tea, pests of, in Ceylon, 294, 300; Helopeltis orophila on, in Belgian Congo, 37; pests of, in India, 246; pests of, in Nyasaland, 85, 86.

Tea (Stored), apparatus for removing Oryzaephilus surinamensis from, 74.

Teak, risk to, of releasing Teleonemia scrupulosa in India, 264.

Technomyrmex detorquens (albipes), associated with mite on litchee in Hawaii, 250.

tectus, Ptinus.

tegalensis, Chionaspis (Aulacaspis).

telarius, Tetranychus.

Telenomus, parasite of Rhoecocoris sulciventris in N. S. Wales, 268.

Telenomus mayri, parasite of Stilpnotia salicis in Sweden, 32.

Telenomus ovivorus, parasite of Coleophora laricella in Ontario, 143.

Teleonemia australis, bait-spray for, on olive in S. Rhodesia, 126.

Teleonemia scrupulosa (attacking Lantana), utilisation of, in Fiji, 291; risk to teak of liberating, in India, 264;

establishment of, in Queensland, 32, 321.

Temperature, effect of, on formation of hypopi by Glycyphagus destructor, 330, 331; relation of locusts and grasshoppers to, 76, 238, 322, 388; relation of other insects to, 7, 13, 18, 52, 73, 74, 75, 89-90, 108, 123, 128, 135, 159, 180, 184, 185, 261, 267, 291, 324, 325, 361, 377, 384, 392, 395, 396; effects of constant and varying, on development of insects, 160, 220; methods of expressing effect of, on insect development, 7, 187, 188; apparatus to regulate variable, 216.

tenebricosa, Capnodis.

Tenebrio molitor, experimentally infesting dehydrated vegetables, 102; distribution of DDT analogue in tissue of, 347; tests of inert dusts on, 157; effects of infra-red rays on, 53.

tenebrioides, Zabrus. tenebrionis, Capnodis.

Tenebroides mauritanicus, possibly predacious on Cydia pomonella in France, 356; measures against, in stored products in Queensland and Oklahoma, 31, 69; not feeding on dehydrated vegetables, 102; distribution of DDT analogue in tissue of, 347.

tenella, Galerucella.

tenellus, Eutettix; Hemiteles.

Tennessee, insect pests in, **25**, **107**, **344**. Tenthecoris bicolor, Tephrosia extract against, on orchids in Brazil, **134**, **135**.

Tephrosia toxicaria, extract of, against Tenthecoris bicolor in Brazil, 135.

Teragra guttifera, measures against, on Populus deltoides in S. Africa, 83.

Terastia meticulosalis, on Erythrina in Madras, 246.

terebrans, Dendroctonus; Ephialtes (Pimpla).

tergata, Euaraba.

terginus, Scaptocoris.

Terminalia, Sinoxylon conigerum on, in Mauritius, 371.

Terminalia catappa, form of Anastrepha fraterculus on, in Mexico, 397.

terminalis, Euproctis. terminifera, Chortoicetes.

Termites, of New Zealand, 234; rendering soil favourable for *Chlorophora excelsa* in Uganda, 329; attacking timber, 69, 73, 234, 243, 244, 247, 290, 298, 318; timbers resistant to, 247; ants predacious on, 122; working in wood occupied by ants, 69; fungi infesting, 290; measures and experiments against, 69, 73, 243, 244, 247, 272, 298, 318.

tessellatus, auct., Corymbites (see C. sjaelandicus).

tessmanni, Mesohomotoma.

testaceipes, Aphidius (Lysiphlebus).

testaceus, Rogas.

testudinea, Hoplocampa.

testulalis, Maruca.

Tetracnemus, species of, not reproducing on Pseudococcus comstocki, 109.

Tetracnemus peregrinus, 109; utilisation of, against Pseudococcus adonidum in California, 59.

Tetracnemus pretiosus, 109.

Tetracrium rectisporum (see Podonectria coccicola).

Tetradecyl Nitrile, mixture containing, not toxic to insects, 198.

Tetramethylthiuram, compounds of, not toxic to Mamestra brassicae, 198.

Tetranaphthene, toxicity of, to Cydia

pomonella, 215.

Tetranychus, creolin emulsion against, in greenhouses in Russia, 304; survey of hibernation of, 328.

Tetranychus althaeae (see T. telarius).

Tetranychus bimaculatus (in U.S.A.), on carnation, 379; on Platanus, 352; on strawberry, 263, 379; spray against, 379; use of, for testing spray deposits, 61.

Tetranychus bioculatus (see Paratetranychus).

Tetranychus crataegi, possibly on prune in

England, 360.

Tetranychus pacificus (in U.S.A.), bionomics and control of, on pear, 285, 286; on Platanus, 352.

Tetranychus peruvianus (see Paratetrany-

chus).

Tetranychus peruviensis, Paratetranychus peruvianus possibly recorded as, on cotton in Peru, 132 (and note).

Tetranychus sacchari (seeParatetranychus). Tetranychus sexmaculatus, spray against, on grapefruit in Bermuda, 172; on Platanus spp. in California, 352.

Tetranychus telarius, on cotton in Azerbaijan, 306; on tomatos in Britain, 404; on greenhouse plants in Kansas, 54; on currant and gooseberry in Ontario, 64; measures against, 54, 306, 404; tests of fumigants on, 275.

Tetranychus willamettei (in U.S.A.), sprays against, on pear, 286; on Platanus, 352.

Tetrastichus, parasite of Neodiprion spp. in Canada, 9; parasite of Rogas plecopterae in India, 390.

Tetrastichus minutus, parasite of Clausenia purpurea in Virginia, 110.

tetricus, auct., Anodontonyx (see A noxius).

Tetropium castaneum, on spruce in Finland, 7.

Tetropium fuscum, on spruce in Finland, 7. texana, Anomis.

texanus, Chelonus.

Texas, cotton pests in, 57, 122, 210, 211, 261, 286, 335; Cydia molesta in, 258; fruit-flies in, 52, 395, 397; miscellaneous pests in, 41, 147, 259, 270; pests of pecan in, 209, 270, 346, 347; beneficial insects and biological control in, 122, 177, 261; Chilocorus cacti introduced into Porto Rico from, 110.

Textiles, insects damaging, 72, 107, 136, 163, 229, 232, 351, 403; testing of treatments against pests of, 190, 267. Thea bisoctonotata, bionomics of, in India.

389.

theae, Pinnaspis (Hemichionaspis).

Thecla echion (see Tmolus). theobaldi, Thomasiniana.

theobroma, Distantiella (Sahlbergella).

Theresia claripalpis, introduction of, into Porto Rico against Diatraea saccharalis. 312.

Theridium lineatum, predacious on Leptinotarsa decemlineata in France, 298.

Thermobia domestica, in buildings in Canada and U.S.A., 107; penetration of laminated paper wrapping by, 272; latent effect of pyrethrum and rotenone dusts on, 340.

Thermodes Hot-air Process, 255.

Thespesia rogersii, Dysdercus spp. on, in Nyasaland, 91.

Thiocyanates, against Coleoptera, 30, 49, 104, 260; against Lepidoptera, 49, 148, 149, 198; against Aphids and thrips, 44; against mites, 44, 234; not acting as stomach poisons, 198; aerosols of, 44; in dusts, 30, 49, 104; in sprays, 148, 149, 234; emulsifier for, 148; and acrylontrile, 44; and lead arsenate, 148, 149; and nicotine, 149; and oil emulsions, 149, 234; and pyrethrum, 30, 104; causing injury to plants, 44, 49, 149.

Thionol, effects of, on Periplaneta americana, 368.

Thomasiniana, on Ribes alpinum in Germany, 308.

Thomasiniana crataegi, 308. Thomasiniana oculiperda, 308.

Thomasiniana theobaldi, investigations on, in Britain, 308.

Thosea, on tea in Ceylon, 300.

Thrips, on cotton in Peru, 132.

Thrips angusticeps, plants attacked by, in Denmark, 381; on beans in Holland, 12; on crucifers in Sweden, 36.

Thrips imaginis, study on populations of, on rose in S. Australia, 129.

Thrips tabaci, in Fiji, 161; in Tanganyika, 85; in U.S.A., 44, 52, 70, 104, 174, 180, 338; on onion, 52, 85, 180, 338; fungus associated with, on pyrethrum, 85; on other plants, 44, 70, 104, 161, 174; relation of, to virus diseases, 70, 174;

measures and experiments against, 44, 52, 104, 161, 180, 338.

Thrips, Citrus (see Scirtothrips citri). Thrips, Onion (see Thrips tabaci).

Thuja occidentalis, Isotomurus on seedlings of, in Denmark, 382; reactions of, to methyl bromide, 50.

Thuja orientalis, reactions of, to methyl

bromide, 50.

Thuja plicata, Phymatodes dimidiatus in timber of, in Br. Columbia, 287.

thujafolia, Cinara (Dilachnus).

Thyraeella collaris (parasite of Plutella maculipennis), establishment and value of, in New Zealand, 269, 376; imported into Pacific Is., 269, 291; effects of temperature and humidity on, 376, 377.

Thyreophagus entomophagus, attacking Lepidosaphes ulmi in U.S.A. and Europe, 357; in stored products, 357.

Thysanoptera, predactious on Coccids, 227, 374; *Orius* spp. attacking, 263; relation of, to plant diseases, 70, 174.

Thysanus, hyperparasite of Saissetia nigra

in Porto Rico, 374.

Thysanus elongatus, parasite of Clausenia purpurea in Virginia, 109.

Thysanus niger, parasite of Clausenia purpurea in Virginia, 109.

Tigridia, Aphids and virus disease of, in U.S.A., 2.

tiliae, Phytocoris.

Timaspis papaveris, on Papaver somni-

ferum in Hungary, 122.

Timber, Coleoptera in, 18, 71, 84, 167, 176, 181, 228, 229, 255, 287, 300, 369, 371, 377, 403, 406; termites in, 234, 243, 244, 247, 298; *Xylocopa* in, 290, 372; treatment of, against pests, 72, 84, 243, 244, 247, 298, 371, 372.

Timothy Grass (see *Phleum pratense*).

Tinea granella, measures against, in

stored grain in Quebec, 225.

Tinea pellionella, in W. Australia, 163;

in India, **72.**

Tineola biselliella, in Australia, 163, 320; in Germany, 230; in U.S.A., 82, 102, 350; infesting drugs, 230; not infesting dehydrated foodstuffs, 102; damaging furs and fabrics, 163, 266, 350; infesting sheepskins and baled wool, 82, 320; bionomics of, 83, 351; natural enemies and disease of, 320, 352; measures and experiments against, 83, 257, 266, 320; procedures for rearing and testing treatments against, 190.

tingitiphagus, Anaphes.

Tinospora, Othreis spp. on, in India, 386, 391.

Tipula czizeki, bionomics of, on vegetables in Germany, 182.

Tipula paludosa, in Germany, 182. tipuliformis, Aegeria (Conopia).

Tmetocera ocellana (see Spilonota).

Tmolus echion, on Capsicum in Hawaii, 250

Toads (see Bufo spp.).

Tobacco, Faustinus spp. on, in Brazil, 288; mites on, in Ceylon, 299; Heliothis virescens on, in Cuba, 278; Xylena exoleta on, in Denmark, 382; Nezara on, in Fiji, 291; Criocerid on, in Hawaii, 250; pests of, in Nyasaland, 85, 86; cutworms on, in Ontario, 140; ants in seed-beds of, in Queensland, 369; pests of, in S. Rhodesia, 126; pests of, in U.S.A., 41, 45, 54, 215, 278, 280, 281, 344; insects and virus diseases in, 160, 247; not infested by Corythaica passiflorae, 264; injured by xanthone, 46; investigation of nornicotine types of, 368.

Tobacco (Stored), pests in, and their control, 100, 127, 230, 232; (seed),

Tribolium confusum in, 127.

Tobacco (as an insecticide), preparation and uses of sprays containing, 293, 364, 365; rendering soap sprays ineffective against Rhoecocoris sulciventris, 268; not protecting stored potatoes from Gnovimoschema operculella, 187; as a carrier for dusts, 54; question of nornicotine in, 368.

Tobacco Flea-beetle (see Epitrix hirti-

pennis).

Tobacco Hornworm (see Protoparce sexta). toddaliae, Ceroplastes.

Togoland, British, cacao pests in, 195, 196.

Togoland, French (see Africa, French

West).

Tomato, Scaptocoris terginus on, in Argentina, 4; Gnorimoschema operculella on, in Australia, 185; pests of, in Brazil, 94, 263, 264, 287, 288; Tetranychus telarius on, in Britain, 404; on, in Ceylon, 299; Phytomyza solani on, in Denmark, 381; Tipula czizeki on, in Germany, 182; Aulacophora foveicollis on, in Greece, 17; pests of, in Hawaii, 250, 251; Leptinotarsa decemlineata on, in Ivory Coast, 165; pests of, in New Zealand, 93, 269; Myzus persicae on, in S. Rhodesia, 126; susceptibility of varieties of, to L. S. Rhodesia, 126; decemlineata in Spain, 166; pests of, in U.S.A., 30, 49, 51, 59, 219, 220, 270; effects of insecticide treatments on, 49, 56, 94; virus disease of, in cineraria, 70.

Tomato (Fruit), precautions against introduction of *Leucinodes elegantalis* into Argentina in, **94.**

Tomato Pinworm (see Keiferia lycoper-sicella).

Tomato Russet Mite (see Phyllocoptes destructor).

Tomato Stem-borer (see Gnorimoschema plaesiosema).

tomentosus, auct., Byturus (see B. urbanus). Tomocera californica, parasite of Saissetia nigra in Hawaii, 374.

Tonga Is., Thyraeella collaris imported into, from Fiji, 291.

tortricidis, Omorgus (Campoplex).

tortricis, Anachaetopsis; Copidosoma.

Tortrix citrana, on orange in California, 113.

Tortrix pallorana, on strawberry in Ohio, 263.

Tortrix postvittana, on vines, etc., in Victoria, 322.

Tortrix rosaceana, on strawberry in Ohio, 263.

Tortrix rosana, on peach and Vaccinium in Holland, 12.

Tortrix viridana, bionomics of, on cork oak in Portugal, 309.

Toxoptera aurantii, on Citrus in Argentina, 4; experimentally transmitting virus disease of cacao in Gold Coast, 66; on Calophyllum and mango in Hawaii, 251; on tea in Nyasaland, 86; natural enemies of, 4, 86.

Toxoptera graminum (on cereals), bionomics and control of, in Argentina, 310, 311; in Oklahoma, 70; in Tanganyika, 85; susceptibility of varieties of barley to, 70.

Toxotrypana curvicauda, fruits infested by, in Mexico and Texas, 397.

Trachea atriplicis, on rhubarb in Denmark, 381.

tragica, Elodia (Arrhinomyia).

. Tragocephala spp., measures against, on cacao in Belgian Congo, 365.

transsylvanica, Hypera. trapezalis, Marasmia.

Traps, suggested use of coloured, for Aleurodids, 392; for Cosmopolites sordidus, 365; for moths, 13, 238, 292. (See Light-traps.)

tremae, Icerya.

Trenches, against Cleonus punctiventris, 302; against Lepidopterous larvae, 94, 104, 246, 402.

Trialeurodes vaporariorum, sprays against, on egg-plant in Virginia, **51**.

Triaspis vestiticida, imported into U.S.A. against Anthonomus grandis, 261.

Tribolium, in stored grain in Algeria, 403; interfering with rearing of Corcyra cephalonica in Bombay, 393; fumigation against, in flour mills in U.S.A., 235, 236.

Tribolium castaneum, in stored groundnuts in Queensland, 31; in stored wheat in Victoria, 129, 130, 293, 318, 319; nutritive requirements and strains of, 235, 293; measures and experiments against, 130, 157, 295, 318, 319.

Tribolium confusum, in stored tobacco seed in S. Rhodesia, 127; in stored cereals and flour in Victoria, 129, 130, 318; experimentally infesting dehydrated foodstuffs, 102; effect of type of food on, 235; penetration of packing materials by, 258; tests and use of fumigants against, 26, 27, 103, 104, 130, 236, 275; action of mercury vapour on eggs of, 165; tests of inert dusts on, 157; toxicity of pyridine derivatives to, 275; effect of infra-red rays on, 53.

Tribolium destructor (see Aphanotus). Trichilia emetica, Selenaspidus sylvaticu

Trichilia emetica, Selenaspidus sylvaticus on, in S. Africa, 107; Rhenea mediata on, in Nyasaland, 86.

1,1,1-Trichlor-2, 2-bis (p-chlorphenyl) ethane (see DDT).

Trichloracetonitrile, experiments with, as a fumigant, 104, 128, 295, 296; and acrylonitrile, 104; and carbon dioxide, 296; determination of, 128.

Trichlorbenzene, and pentachlorphenol, soil treatment with, against termites,

69.

Trichlorethylene, and ethylene dichloride, fumigation with, against pests of stored grain, 318, 319; and dibutyl phthalate, as solvent for derris resins, 211.

Trichodes laminatus, destroying eggs of Dociostaurus maroccanus in Iraq, 201.

Trichogramma (parasite of Lepidopterous eggs), utilisation of, in Russian Union, 306; in U.S.A., 28, 241; effect of rearing conditions on value of, 306.

Trichogramma australicum, rearing and utilisation of, against Proceras sacchari-

phagus in Mauritius, 370.

Trichogramma koehleri, in Argentina, 117.
Trichogramma minutum (parasite of Lepidopterous eggs), in India, 389, 393; in Porto Rico, 97, 243; in Spain, 125; in U.S.A., 26, 95, 112, 177; mass rearing of, 125, 393; field utilisation of, 26, 97, 125.

Trichogramma pretiosum, rearing and utilisation of, against Cydia pomonella

in Spain, **125.**

Trichomalus trujilloi, considered distinct from Dibrachys cavus (q.v.), 95.

Trichomma enecator, parasite of Cydia pomonella in France, 355; suggested introduction of, into Canada, 356.

Trichopoda pennipes, introduction of, into Australia against Nezara viridula, 322; utilisation of, against Anasa tristis in U.S.A., 21.

Trichoptilus wahlbergi, bird destroying, on sweet potato in Mauritius, 371.

Triethanolamine Oleate, reducing penetration of oil into plant tissue, 253.

Trifidaphis radicicola, on beans in California, 338.

Trifolium (see Clover).

Trigonella foenum-graecum, relation of wireworms to, in California, 260.

trigonum, Eutrombidium.

triguttata, Carneocephala.

trilineata, Lema.

trilobitiformis, Pseudaonidia.

Trinidad, insects on cacao in, 78, 147; miscellaneous pests and beneficial insects in, 147, 176, 313; Coccinellids imported into, 313.

tripterus, Mecistorhinus. triquetra, Homalodisca.

Trissolcus euschisti, parasite of Coleophora laricella in Ontario, 143.

tristicolor, Orius.

tristigmus, Euschistus.

tristis, Anasa; Perilampus.

tritici, Contarinia.

Tritonia crocata, Myzus persicae and virus disease of, in U.S.A., 2.

Triumfetta rhomboidea, parasitism of Pempherulus affinis on, in India, 21.

trivittatus, Leptocoris.

Trogoderma granarium, in drugs in Germany, 230.

Trogoderma ornatum, synonymy of, 330. Trogoderma versicolor, in dried eggs and milk in California, 102; resistance of, to fumigants, 295.

Trogophloeus elongatulus, on cucumber in Holland, 12.

troilus, Anthicus.

Trombidium, attacking Leptinotarsa decemlineata in France, 298.

Trombidium locustarum (see Eutrombidium trigonum).

Trombidium sericeum (see Eutrombidium trigonum).

trujilloi, Trichomalus (see Dibrachys cavus). truncatella, Litomastix (Copidosoma).

truncatulus, Hypolixus.

tryoni, Dacus (Strumeta) ferrugineus; Opius (Biosteres). trypetes, Chilo.

Trypodendron lineatum (bivittatum), in fire-killed Douglas fir in U.S.A., 40.

Tsuga, drought favouring infestation of, by Melanophila fulvoguttata in U.S.A., 43.

Tubercularia coccicola (see Nectria tuberculariae).

tuberculatus, Kalotermes (Calotermes).

tuberis, Epitrix.

tuberosa, Sarcophaga. tucumanus, Rhigopsidius.

Tulip, Aphids and virus diseases of, in U.S.A., **2, 38.**

Tung (see Aleurites).

Turanodinia coccidarum, gen. et sp. n., predacious on Pseudococcus comstocki in Uzbekistan, 328.

Turf, Hoplia philanthus on, in Britain, 156; Sericesthis pruinosa on, in N. S. Wales, **161**; insects damaging, in U.S.A., 115, 219.

Turkey, Coccinella septempunctata in, 225; forest pests in, 7.

Turkey Red Oil, 257.

Turnip, Hylemyia floralis on, in Britain, 121; Plutella maculipennis on, in Fiji, 290: Myzus persicae on, in S. Rhodesia, 126; Mamestra brassicae on, in Russia, 305; Thrips angusticeps on, in Sweden,

36; pests of, in U.S.A., 178, 191. Turnip Flea-beetle (see Phyllotreta nemo-

rum). Turpentine, against clothes moths and carpet beetles, 163; in mixtures against Coleoptera in timber, 84: attracting Dendroctonus spp., 43.

Two-spotted Collops (see Collops bi-

punctatus).

Two-striped Grasshopper (see Melanoplus bivittatus).

Tychius, measures against, on lucerne in Russia, 305.

Tyloderma fragariae, on strawberry in Ohio, 262.

typographus, Ips.

Tyroglyphus farinae, in cheese and flax seed in New Zealand, 77, 269; technique for studying, 77.

Tyroglyphus siro, auct. (see Tyrolichus

Tyrolichus casei (in cheese), in New Zealand, 77, 130, 269; in Washington, fumigation against, 130, 269, 343; 343; technique for studying, 77.

Tyrophagus putrescentiae (longior), technique for studying, in cheese in New Zealand, 77.

tyrrhea, Gonimbrasia.

U.

Uganda, Phytolyma lata and its parasites on Chlorophora excelsa in, 329; cotton pests in, 88, 127; pests and disease of sweet potato in, 329, 330.

Ulex europaeus, utilisation of Apion ulicis against, in New Zealand, 269.

ulicis, Apion.

ulkei, Rhyncolus (Hexarthrum).

ulmi, Lepidosaphes.

ulmi, auct., Oligonychus (see Paratetranychus pilosus).

Ulmus (see Elm).

Uloborus geniculatus, predacious on Tineola biselliella in Australia, 320.

Ultrawet, 240.

umbrosus, Adoretus.

Unaspis citri (on Citrus), in Argentina, 132; in Australia, 248; in Porto Rico, 243; natural enemies of, 133, 243, 248; bionomics and control of, 133, 248.

undalis, Hellula.

undecimpunctata, Diabrotica.

unicinctus, Stizoides.

unicolor, Stromatium (see S. fulvum).

uniformis, Curculio.

United States of America, cereal pests in, 49, 147; pests of Corylus avellana in, 341; pests of forest and shade tre s in, 31, 43, 53, 71, 173, 238, 270, 310, 313, 314, 352, 353, 379; orchard pests in, 94, 100, 219, 241, 258, 273, 314, 334; pests of small fruits in, 96, 378, 401, 402; pests of ornamental plants in, 1, 2, 38, 43, 71, 96, 100, 378; pests of rubber-producing plants in, 104, 105, 259; pests of vegetable and root crops in, 39, 48, 49, 70, 71, 100, 141, 173, 191, 214, 220, 260, 270, 278, 287; vine pests in, 27, 28, 172; insects and virus diseases of plants in, 1, 2, 38, 39, 70, 71, 172, 173; pests of stored products in, 100, 102, 235, 236, 258, 259, 318, 350; book on household pests in, 318; pests of timber in, 176, 406; grasshoppers and Anabrus simplex in, 193, 259; Pantomerus in, 53, 111, 259; Popillia japonica in, 49, 50, 55, 58, 63, 79, 180, 347, 402; Tachycines asynamorus in, beneficial Arthropods and biological control in, 28, 70, 96, 193, 218, 238, 241, 261, 310, 352, 375; plant pest legislation in, 111, 145, 193, 286; pests intercepted in quarantine in, 100, 223; beneficial insects imported into other countries from, 95, 261, 310. (See also under the various States.)

ununguis, Paratetranychus.

urbana, Ctenolepisma.

urbanus, Byturus.

Uredinella spinulosa, infesting Aspidiotus in Ceylon, 232.

Urena lobata, Aonidiella aurantii on, in Brazil, 204.

urichi, Acropyga (Rhizomyrma).

urozonus, Eupelmus.

Uruguay, Cydia spp. in, 94; parasites of C. molesta in, 95.

Ustilago zeae, 239.

Utah, grasshoppers and Anabrus simplex in, 114, 174; pests of lucerne in, 101, 106, 206; miscellaneous pests in, 101, 106, 281, 353; natural enemies of insects in, 106, 206, 263.

V.

vaccinii, Eriophyes; Mineola.

Vaccinium (see Blueberry and Cranberry). Vaccinium myrtillus (Bilberry), Geometrid on, in Britain, 360.

Vacuum Fumigation, 103, 111, 236, 252, 375.

valens, Dendroctonus.

validirostris, Pissodes.

Vanessa atalanta, parasite of, 281.

vaporariorum, Trialeurodes.

variabilis, Hypera; Pheidole.

varianus, Macremphytus. variegana, Argyroploce.

variegata, Adonia.

variegatus, Faustinus (Euxenus) (see F.

cubae); Zonocerus. variolarius, Euschistus.

varivestis, Epilachna.

vastatrix, Pĥylloxera (see P. vitifoliae).

Vatsol OS, 240.

Venezuela, miscellaneous pests in, 147, 263.

ventralis, Rhizobius.

ventrellus, Dichomeris.

ventricosus, Pediculoides.

verbasci, Anthrenus.

Verbena bonariensis, attracting Pieris rapae in S. Carolina, 348.

Vermont, disease of Gilpinia hercyniae in, 398.

vernata, Paleacrita.

versicolor, Trogoderma.

versicolora, Plagiodera.

versutus, Adoretus.

Verticillium larvarum, sp. n., infesting Limacodid in Ceylon, 232.

Vesiculaphis caricis, on tomato in Hawaii, 251.

vestiticida, Bracon (Microbracon); Triaspis.

vestitus, Anthonomus.

Vetches (Vicia spp.), pests of, in U.S.A., 22, 100, 101, 259, 342; Hypera variabilis reared on, 41.

viburni, Aphis.

Viburnum opulus, Aphids on, in Britain, 382.

Vicia (see Vetches).

Vicia faba (see Broad Beans).

Victoria, Aphid and virus disease of carrot in, 292, 369; *Cydia molesta* on peach in, 322; pests of stored cereals in, 129, 235, 292, 318, 319; Coleoptera in fire-killed timber in, 377; biological control of *Hypericum perforatum* in, 321.

viennensis, Amphitetranychus (see Tetra-

nychus crataegi).

vigintioctopunctata, Epilachna.

Vine, Grape, Hemiberlesia rapax on, in Argentina, 4; pests of, in Australia, 162, 322; Lygus spinolae on, in Austria,

117; Eulecanium persicae on, in Egypt, 228; pests of, in Germany, 13, 14; Euxoa nigricans on, in Holland, Polychrosis botrana on, in Italy, 118; pests of, in Spain, 298; Phylloxera on, in Switzerland, 124; pests of, in U.S.A., 27, 28, 44, 172, 214, 259; Jassids transmitting virus disease of, 172; relation of races of Phylloxera to varieties of, 118, 124.

Vinegar, in bait for fruit-flies, 134.

violacea, Magdalis.

virens, Agromyza.

virescens, Aphthona (see A. euphorbiae); Heliothis.

Ferrisia (Ferrisiana, Pseudovirgata,

coccus).

Virginia, Pseudococcus comstocki on apple in, 109, 208, 209, 258; Scolytid and fungus in pine in, 287; tobacco pests in, 41, 45, 280, 281; pests of vegetables in, 51, 59, 212, 214; beneficial insects and biological control in, 109, 208, 209; Popillia japonica imported into Canada from, 137.

Virginia, West, natural enemies and ·biological control of Pseudococcus comstocki in, 208, 209, 258; Macrosteles divisus and aster-yellows in potato in,

173; termites in, 69.

virginica, Xylocopa. viridana, Tortrix.

viridis, Coccus; Smynthurus.

viridula, Gastroides; Nezara.

Virus Diseases (in plants*), relation of Aleurodids to, 126, 330; relation of Aphids to, 1, 2, 33, 34, 38, 39, 40, 66, 70, 71, 160, 174, 247, 292, 323, 365, 369, 378, 392; Delphacodes striatellus transmitting, 183; relation of Jassids to, 71, 101, 172, 173, 281, 292, 378; relation of mealybugs to, 66, 196; relation of Mesohomotoma tessmanni to, 66; relation of Thrips tabaci to, 70, 174; mechanical transmission of, 1, 70, 174, 247, 292; regulations against, in Nyasaland, 85; book on, 192.

viteana, Polychrosis.

vitellinae, Phyllodecta.

Syrphus vitiensis, Haplogonatopus; corollae.

vitifoliae, Phylloxera.

vitis, Pulvinaria.

vittatus, Collops. vivida, Parasa.

Volcanic Ash, experiments with, against insect pests, 131.

Voria ruralis, parasite of Plusia brassicae in Arizona, 112.

vulgaris, Bracon (Microbracon); Chrvsopa; Melolontha (see M. melolontha).

vulnerator, Pristomerus. vulpinus, Dermestes (see D. maculatus). vuteria, Sesamia.

\mathbf{W} .

wahlbergi, Trichoptilus. wahlbomiana, Cnephasia. walkeri, Eutermes.

Walnut, Cydia pomonella on, in California, 240; C. pomonella on, in France, 355, 356; Paratetranychus pilosus on, in Britain, 233; (kernels), development of Plodia interpunctella on, 231.

Walnut-shell Flour, as carrier for dusts,

239, 283.

Washington, pests of fire-killed Douglas fir in, 40; pests of leguminous crops in, 22, 24, 100, 101, 259; miscellaneous pests in, 21-24, 114; orchard pests in, 22, 23, 57, 108, 258, 274, 281, 284, 285; pests of ornamental plants in, 24, 70, 174; potato pests in, 22, 23, 39, 154, 220, 260; wireworms in, 23, 260; insects and virus diseases of plants in, 39, 70, 174; beneficial insects and biological control in, 21, 22, insecticide situation in, 343.

Water, Hot, uses of, against pests, 35, 72,

120, 163, 260, 375.

Water Hyacinth (see Eichhornia).

Water-cucumber (see Lagernaria vulgaris), Watercress, Phaedon cochleariae on, in Britain, 405.

Watermelon, experiment with Aulacophora foveicollis and, in Greece, 17; Daihinia brevipes on, in Oklahoma, 270; tests of extract of seeds of, as an insecticide, 65.

Watsonia, Aphid transmitting virus to Tritonia from, in U.S.A., 2.

Wattle (see Acacia).

Webbing Clothes Moth (see biselliella).

websteri, Sagaritis.

West Indies, insect pests of food crops in, (See also under the various 313. Islands.)

westwoodi, Dicladocerus.

Wheat, pests of, in E. Africa, 85, 86; Toxoptera graminum on, in Argentina, Anodontonyx noxius on, in Australia, 161; pests of, in Britain, 92, 265, 293, 294, 297, 307; Cephus cinctus on, in Canada, 98; C. pygmaeus on, in Denmark, 382; Zabrus tenebrioides on, in Hungary, 11: cophora argyrogaster on, in New Caledonia, 161; pests of, in Asiatic Russia, 303; pests of, in U.S.A., 23, 101, 103,

^{*} These references are also indexed under the individual plants infected.

151, 237; correlation of infestation by Cecidomyiids with yield of, 293; tests of resistance of varieties of, to Maye-Melanoplus destructor, 103; bivittatus ovipositing in piles of, 153; attacked by Phyllobius pyri, 92; relation of wireworms to, 23, 265, 294; as bait for wireworm, 405; dinitro compound not injurious to, 54.

Wheat (Stored), pests in, and their control, 69, 102, 127, 129, 131, 151, 235, 236, 237, 257, 292, 295, 318, 319, 320, 330, 361, 372, 402; effects of fumigants on, 236, 237, 259, 296; sorption of fumigants by, 128, 236, 237, 264; determination of fumigants in, 128, 264; (rolled), in baits for Anabrus simplex 259 simplex, 259.

Wheat Blossom Midges (see Contarinia tritici and Sitodiplosis mosellana).

Whiteflies, and virus diseases, 126, 330. White-footed Mice (see *Peromyscus*).

White-fringed Beetles (see Pantomorus (Graphognathus)).

whittieri, Quaylea.

Widdringtonia whytei, Cinara thujafolia on, in Nyasaland, 86.

willamettei, Tetranychus.

Willow, Plagiodera versicolora on, U.S.A., 71. (See Salix spp.)

wilsoni, Platypus. Wilt Diseases, in Gilpinia hercyniae, 47, 48, 399; in Lepidoptera, 81, 82, 238,

300, 399. Wind, spread of insects by, 158, 174, 248,

357, 392. Winter Moth (see Operophtera brumata). Wire Gauze, as barrier against Bagrada

hilaris, 293.

Wireworms, keys to, in Germany and Britain, 11, 296; effect of altitude on populations of, in Britain, 405; populations of, in relation to crops, 23, 198, 260, 265, 266, 294, 368; studies on growth, movements and physiology of, 123, 294; measures and experiments against, 23, 55, 56, 104, 257, 260, 337, 405; predacious on Hylemyia floralis, 122.

Wisconsin, pests in cheese factories in, Cercopid on pines in, 107; Macrosiphum onobrychis on peas in, 276, 277, 282; Nematodes infesting Lachnosterna in, 61.

woglumi, Aleurocanthus.

woodi, Acarapis.

Woodlice, tests of insecticides on, 197, 198, 257.

Woodpecker, Hairy (see Dryobates villosus).

Wool, insects infesting, 82, 83, 163, 320,

Woollen Goods (see Textiles).

Woolly Apple Aphis (see Eriosoma lanigerum).

Worcesterberry, Thomasiniana theobaldi not ovipositing on, 308.

W.S.C. Dynamite, Modified, 151.

Wyoming, Anabrus simplex in, 114; new food-plant of Eutettix tenellus in, 281; Epitrix tuberis not occurring in, 220.

\mathbf{X} .

xanthius, Taeniothrips.

xanthodes, Dacus (Notodacus).

xanthomelaena, Galerucella (see G. luteola). Xanthone, spraying with, against Cydia pomonella, 149; dusting with, against Epitrix hirtipennis, 45; causing injury to plants, 46, 149; analysis of spray residues of, 128.

Xanthosoma, Ferrisia virgata on, in

Nigeria, 196.

xanthostigmata, Spilochalcis.

Xenopygus analis (see Philothalpus).

Xestobium rufovillosum, humidity requirements of, in timber in Germany, 18; larval characters of, 167.

Xyleborus camphorae, on Pinus taeda in Mauritius, 371.

Xyleborus dispar, on frost-damaged fruit trees in Holland, 12.

Xyleborus fornicatus fornicatior, effect of manuring on damage to tea by, in Ceylon, 294, 300.

Xylena exoleta, on tobacco and poppy in Denmark, 382.

Xylene, as solvent for benzene hexachloride, 257.

2,4-Xylidene Acetate, tests of toxicity of,

to Lepidopterous larvae, 344. Xylobiops basilare, in timber in U.S.A.,

176. Xylocopa, limewash protecting timber

against, in India, 372.

Xylocopa senior (see Mesotrichia).

Xylocopa virginica, in imported timber in Bermuda, 290.

xylophagorum, Rhoptrocerus (Pachyceras).

Y.

Yam, increasing damage to cotton by Helopeltis in Nigeria, 383; soil fumigant not harmful to, 56.

Yam Bean (see Pachyrhizus erosus).

yawi, Anagrus.

Yeast, in baits, 340, 396; new Psocid reared on, 181.

Yew (see Taxus).

Yezabura (see Anuraphis). ypsilon, Agrotis: Mormidea. INDEX. 503

Z.

Zabrus tenebrioides, bionomics and control of, in Hungary, 11.

Zanthoxylum clava-herculis, tests of extract of, as an insecticide, 65.

zea-mais, Calendra (see Calandra oryzae). Zicrona coerulea, predacious on Leptinotarsa decemlineata in France, 298.

Zinc Arsenate, in dusts and sprays against *Epitrix* on potato, **22**.

Zinc Arsenite, effect of spraying with, against *Epilachna varivestis* on beans, **176.**

Zinc Carbonate, reducing toxicity of derris dust to Musca domestica, 78.

Zinc Chloride, treatment of timber with, against pests, 84, 247, 299, 371.

Zinc Dimethyl Dithiocarbamate, DDT in spray containing, 173.

Zinc Oxide, as a safener for calcium arsenate, 211; increasing toxicity of derris dust to *Musca domestica*, 78; unsuitable for use with nicotine sulphate, 98.

Zinc Sulphate, solution of, against Coleoptera in timber, 84; and lime, spraying with, against *Empoasca fabae*, 317; in mixed sprays, 191, 210, 273, 314, 364. zinckenella, Etiella.

Zizera labradus mangoensis, on beans in Fiji, 290.

ziziphus, Parlatoria.

zizyphi, Aspidiotus combreti.

Zizyphus jujuba, Lasiocampid on, in India, 393.


Zonocerus variegatus, on cotton in Belgian Congo, 367.

Zoysia japonica, Crambus haytiellus on, in Florida, 115.

Zygadenus fremontii, virus diseases in, 38. Zyklon, 255.







NOTICES

Secretaries of Societies and Editors of Journals willing to exchange their publications with those of the Institute are requested to communicate with the Director.

The Annual Subscription, in advance, to the Review, Series A (Agricultural), is 30s. post free; Series B (Medical and Veterinary), 15s. post free. Subscriptions to the current Volume received after 30th June in each year will be charged at the published price, viz.: Series A, 36s.; Series B, 18s. Prices of Back Volumes on application.

Orders and Subscriptions should be sent direct to the Director, Imperial Institute of Entomology, 41, Queen's Gate, London, S.W. 7, or through a bookseller.

